



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:55 AM GMT

PDB ID : 3K6X
Title : M. acetivorans Molybdate-Binding Protein (ModA) in Molybdate-Bound Close Form with 2 Molecules in Asymmetric Unit Forming Beta Barrel
Authors : Chan, S.; Chernishof, I.; Giuroiu, I.; Sawaya, M.R.; Chiang, J.; Gunsalus, R.P.; Arbing, M.A.; Perry, L.J.
Deposited on : 2009-10-09
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1640 (2.28-2.24)
Clashscore	102246	1095 (2.26-2.26)
Ramachandran outliers	100387	1063 (2.26-2.26)
Sidechain outliers	100360	1063 (2.26-2.26)
RSRZ outliers	91569	1647 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	354	 5% 84% 6% 9%
1	B	354	 5% 84% 6% 9%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5137 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Solute-binding protein MA_0280.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	0	0
			2494	1578	395	511	10			
1	B	322	Total	C	N	O	S	0	0	0
			2498	1580	395	513	10			

There are 56 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP Q8TTZ5
A	0	LYS	-	EXPRESSION TAG	UNP Q8TTZ5
A	1	HIS	-	EXPRESSION TAG	UNP Q8TTZ5
A	2	HIS	-	EXPRESSION TAG	UNP Q8TTZ5
A	3	HIS	-	EXPRESSION TAG	UNP Q8TTZ5
A	4	HIS	-	EXPRESSION TAG	UNP Q8TTZ5
A	5	HIS	-	EXPRESSION TAG	UNP Q8TTZ5
A	6	HIS	-	EXPRESSION TAG	UNP Q8TTZ5
A	7	PRO	-	EXPRESSION TAG	UNP Q8TTZ5
A	8	MET	-	EXPRESSION TAG	UNP Q8TTZ5
A	9	SER	-	EXPRESSION TAG	UNP Q8TTZ5
A	10	ASP	-	EXPRESSION TAG	UNP Q8TTZ5
A	11	TYR	-	EXPRESSION TAG	UNP Q8TTZ5
A	12	ASP	-	EXPRESSION TAG	UNP Q8TTZ5
A	13	ILE	-	EXPRESSION TAG	UNP Q8TTZ5
A	14	PRO	-	EXPRESSION TAG	UNP Q8TTZ5
A	15	THR	-	EXPRESSION TAG	UNP Q8TTZ5
A	16	THR	-	EXPRESSION TAG	UNP Q8TTZ5
A	17	GLU	-	EXPRESSION TAG	UNP Q8TTZ5
A	18	ASN	-	EXPRESSION TAG	UNP Q8TTZ5
A	19	LEU	-	EXPRESSION TAG	UNP Q8TTZ5
A	20	TYR	-	EXPRESSION TAG	UNP Q8TTZ5
A	21	PHE	-	EXPRESSION TAG	UNP Q8TTZ5
A	22	GLN	-	EXPRESSION TAG	UNP Q8TTZ5
A	23	GLY	-	EXPRESSION TAG	UNP Q8TTZ5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	ALA	-	EXPRESSION TAG	UNP Q8TTZ5
A	25	MET	-	EXPRESSION TAG	UNP Q8TTZ5
A	26	ALA	-	EXPRESSION TAG	UNP Q8TTZ5
B	-1	MET	-	EXPRESSION TAG	UNP Q8TTZ5
B	0	LYS	-	EXPRESSION TAG	UNP Q8TTZ5
B	1	HIS	-	EXPRESSION TAG	UNP Q8TTZ5
B	2	HIS	-	EXPRESSION TAG	UNP Q8TTZ5
B	3	HIS	-	EXPRESSION TAG	UNP Q8TTZ5
B	4	HIS	-	EXPRESSION TAG	UNP Q8TTZ5
B	5	HIS	-	EXPRESSION TAG	UNP Q8TTZ5
B	6	HIS	-	EXPRESSION TAG	UNP Q8TTZ5
B	7	PRO	-	EXPRESSION TAG	UNP Q8TTZ5
B	8	MET	-	EXPRESSION TAG	UNP Q8TTZ5
B	9	SER	-	EXPRESSION TAG	UNP Q8TTZ5
B	10	ASP	-	EXPRESSION TAG	UNP Q8TTZ5
B	11	TYR	-	EXPRESSION TAG	UNP Q8TTZ5
B	12	ASP	-	EXPRESSION TAG	UNP Q8TTZ5
B	13	ILE	-	EXPRESSION TAG	UNP Q8TTZ5
B	14	PRO	-	EXPRESSION TAG	UNP Q8TTZ5
B	15	THR	-	EXPRESSION TAG	UNP Q8TTZ5
B	16	THR	-	EXPRESSION TAG	UNP Q8TTZ5
B	17	GLU	-	EXPRESSION TAG	UNP Q8TTZ5
B	18	ASN	-	EXPRESSION TAG	UNP Q8TTZ5
B	19	LEU	-	EXPRESSION TAG	UNP Q8TTZ5
B	20	TYR	-	EXPRESSION TAG	UNP Q8TTZ5
B	21	PHE	-	EXPRESSION TAG	UNP Q8TTZ5
B	22	GLN	-	EXPRESSION TAG	UNP Q8TTZ5
B	23	GLY	-	EXPRESSION TAG	UNP Q8TTZ5
B	24	ALA	-	EXPRESSION TAG	UNP Q8TTZ5
B	25	MET	-	EXPRESSION TAG	UNP Q8TTZ5
B	26	ALA	-	EXPRESSION TAG	UNP Q8TTZ5

- Molecule 2 is MOLYBDATE ION (three-letter code: MOO) (formula: MoO₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Mo	O	0	0
			5	1	4		
2	B	1	Total	Mo	O	0	0
			5	1	4		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	47	Total	O	0	0
			47	47		
4	B	48	Total	O	0	0
			48	48		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Solute-binding protein MA_0280



- Molecule 1: Solute-binding protein MA_0280



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	147.19Å 65.59Å 94.77Å 90.00° 123.95° 90.00°	Depositor
Resolution (Å)	57.74 – 2.25 57.78 – 2.25	Depositor EDS
% Data completeness (in resolution range)	87.5 (57.74-2.25) 87.6 (57.78-2.25)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 2.25Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.237 , 0.273 0.238 , 0.271	Depositor DCC
R_{free} test set	1591 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	44.7	Xtriage
Anisotropy	0.150	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	2 of 31302 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5137	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 86.07 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.0488e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MOO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/2546	0.70	1/3474 (0.0%)
1	B	0.43	0/2550	0.69	0/3479
All	All	0.43	0/5096	0.69	1/6953 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	223	LEU	CA-CB-CG	5.03	126.87	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2494	0	2375	11	0
1	B	2498	0	2379	12	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	20	0	0	0	0
3	B	20	0	0	0	0
4	A	47	0	0	0	0
4	B	48	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5137	0	4754	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:123:MET:HG2	1:B:292:ILE:HD11	1.85	0.57
1:B:80:GLN:HB2	1:B:160:ASP:HB2	1.88	0.56
1:A:127:TYR:HA	1:A:234:LEU:HD11	1.90	0.54
1:A:123:MET:HG2	1:A:292:ILE:HD11	1.91	0.53
1:A:59:GLU:HG3	1:A:71:VAL:HB	1.91	0.51
1:B:301:ASN:HD22	1:B:301:ASN:H	1.60	0.50
1:B:42:VAL:O	1:B:304:ASN:ND2	2.45	0.48
1:B:121:ASN:HB2	1:B:294:TYR:HE2	1.79	0.48
1:B:59:GLU:HG3	1:B:71:VAL:HB	1.96	0.47
1:B:162:PRO:HA	1:B:165:TYR:CZ	2.50	0.46
1:A:344:PRO:HD2	1:A:347:LEU:HD12	1.97	0.46
1:A:21:PHE:CD1	1:A:72:GLN:HB3	2.51	0.46
1:B:344:PRO:HD2	1:B:347:LEU:HD12	1.98	0.45
1:B:104:PRO:HA	1:B:108:VAL:HB	1.98	0.44
1:A:22:GLN:HA	1:A:22:GLN:HE21	1.85	0.42
1:B:108:VAL:HA	1:B:109:PRO:HA	1.75	0.42
1:A:251:GLN:HE21	1:A:327:ASN:HB3	1.86	0.41
1:A:108:VAL:HA	1:A:109:PRO:HA	1.83	0.41
1:A:148:ARG:O	1:A:220:LYS:NZ	2.52	0.41
1:A:148:ARG:HA	1:A:149:PRO:HD3	1.90	0.41
1:A:286:VAL:HB	1:B:199:GLU:HG2	2.03	0.40
1:B:79:ALA:HB3	1:B:160:ASP:HA	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/354 (90%)	308 (97%)	10 (3%)	0	100	100
1	B	318/354 (90%)	309 (97%)	8 (2%)	1 (0%)	46	52
All	All	636/708 (90%)	617 (97%)	18 (3%)	1 (0%)	52	61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	67	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/298 (90%)	264 (98%)	5 (2%)	65	75
1	B	270/298 (91%)	267 (99%)	3 (1%)	80	88
All	All	539/596 (90%)	531 (98%)	8 (2%)	72	82

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	TYR
1	A	22	GLN
1	A	138	ASN
1	A	227	GLU
1	A	301	ASN
1	B	20	TYR
1	B	227	GLU
1	B	301	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	GLN
1	A	138	ASN
1	A	301	ASN
1	A	329	GLN
1	B	22	GLN
1	B	65	GLN
1	B	301	ASN
1	B	329	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MOO	A	500	-	2,4,4	6.00	2 (100%)	0,6,6	0.00	-
3	SO4	A	501	-	4,4,4	0.22	0	6,6,6	0.08	0
3	SO4	A	502	-	4,4,4	0.20	0	6,6,6	0.09	0
3	SO4	A	503	-	4,4,4	0.22	0	6,6,6	0.08	0
3	SO4	A	504	-	4,4,4	0.22	0	6,6,6	0.08	0
2	MOO	B	500	-	2,4,4	6.07	2 (100%)	0,6,6	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	501	-	4,4,4	0.25	0	6,6,6	0.08	0
3	SO4	B	502	-	4,4,4	0.22	0	6,6,6	0.10	0
3	SO4	B	503	-	4,4,4	0.22	0	6,6,6	0.09	0
3	SO4	B	504	-	4,4,4	0.21	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MOO	A	500	-	-	0/0/0/0	0/0/0/0
3	SO4	A	501	-	-	0/0/0/0	0/0/0/0
3	SO4	A	502	-	-	0/0/0/0	0/0/0/0
3	SO4	A	503	-	-	0/0/0/0	0/0/0/0
3	SO4	A	504	-	-	0/0/0/0	0/0/0/0
2	MOO	B	500	-	-	0/0/0/0	0/0/0/0
3	SO4	B	501	-	-	0/0/0/0	0/0/0/0
3	SO4	B	502	-	-	0/0/0/0	0/0/0/0
3	SO4	B	503	-	-	0/0/0/0	0/0/0/0
3	SO4	B	504	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	MOO	O1-MO	5.95	1.87	1.73
2	B	500	MOO	O1-MO	5.99	1.87	1.73
2	A	500	MOO	O2-MO	6.05	1.87	1.73
2	B	500	MOO	O2-MO	6.14	1.87	1.73

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	322/354 (90%)	0.68	18 (5%)	28 31	15, 33, 48, 74	0
1	B	322/354 (90%)	0.65	19 (5%)	26 28	13, 34, 48, 97	1 (0%)
All	All	644/708 (90%)	0.66	37 (5%)	27 30	13, 34, 48, 97	1 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	135	ASP	4.9
1	B	22	GLN	4.9
1	B	151	VAL	4.8
1	A	131	SER	4.3
1	A	135	ASP	4.0
1	A	21	PHE	3.8
1	A	237	GLY	3.6
1	B	19	LEU	3.4
1	A	132	LYS	3.4
1	A	341	ASP	3.3
1	B	21	PHE	3.3
1	B	234	LEU	3.3
1	A	236	THR	3.2
1	A	137	ILE	3.2
1	A	140	ASP	3.2
1	B	69	VAL	3.2
1	A	257	VAL	3.2
1	A	22	GLN	3.1
1	A	254	PHE	3.0
1	B	140	ASP	3.0
1	B	230	LEU	2.9
1	A	145	ILE	2.8
1	B	237	GLY	2.8
1	B	127	TYR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	131	SER	2.6
1	A	127	TYR	2.6
1	B	150	ASP	2.4
1	A	347	LEU	2.4
1	B	134	GLY	2.3
1	A	238	GLU	2.3
1	B	235	GLU	2.3
1	A	152	ARG	2.3
1	B	179	ASN	2.2
1	B	341	ASP	2.2
1	B	325	ILE	2.2
1	B	152	ARG	2.1
1	A	322	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	MOO	A	500	5/5	0.99	0.17	0.47	25,26,28,31	0
2	MOO	B	500	5/5	0.99	0.16	0.08	28,30,30,33	0
3	SO4	B	501	5/5	0.88	0.16	-0.77	91,91,91,92	0
3	SO4	A	501	5/5	0.86	0.15	-0.99	82,82,82,82	0
3	SO4	A	502	5/5	0.82	0.16	-	77,77,78,78	0
3	SO4	B	502	5/5	0.91	0.20	-	66,67,67,68	0
3	SO4	B	503	5/5	0.82	0.19	-	116,116,116,116	0
3	SO4	A	503	5/5	0.90	0.12	-	105,106,106,106	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	SO4	A	504	5/5	0.86	0.25	-	104,104,104,104	0
3	SO4	B	504	5/5	0.75	0.30	-	99,99,99,99	0

6.5 Other polymers [i](#)

There are no such residues in this entry.