



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 10:53 PM BST

PDB ID : 2KA3
Title : Structure of EMILIN-1 C1Q-like domain
Authors : Verdone, G.; Corazza, A.; Colebrooke, S.A.; Cicero, D.O.; Eliseo, T.; Boyd, J.; Doliana, R.; Fogolari, F.; Viglino, P.; Colombatti, A.; Campbell, I.D.; Esposito, G.
Deposited on : 2008-10-30

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
SOLUTION NMR




The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	162	
1	B	162	
1	C	162	

2 Ensemble composition and analysis

This entry contains 10 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:846-A:926, A:946-A:995, B:846-B:926, B:946-B:995, C:846-C:926, C:946-C:995 (393)	0.17	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 3, 4, 5, 6, 7, 8, 9, 10
Single-model clusters	2

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 5805 atoms, of which 2886 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called EMILIN-1.

Mol	Chain	Residues	Atoms						Trace
1	A	131	Total	C	H	N	O	S	0
			1935	619	962	164	188	2	
1	B	131	Total	C	H	N	O	S	0
			1935	619	962	164	188	2	
1	C	131	Total	C	H	N	O	S	0
			1935	619	962	164	188	2	

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	834	MET	-	CLONING ARTIFACT	UNP Q9Y6C2
A	835	ARG	-	CLONING ARTIFACT	UNP Q9Y6C2
A	836	GLY	-	CLONING ARTIFACT	UNP Q9Y6C2
A	837	SER	-	CLONING ARTIFACT	UNP Q9Y6C2
A	838	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
A	839	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
A	840	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
A	841	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
A	842	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
A	843	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
A	844	GLY	-	CLONING ARTIFACT	UNP Q9Y6C2
A	845	SER	-	CLONING ARTIFACT	UNP Q9Y6C2
B	834	MET	-	CLONING ARTIFACT	UNP Q9Y6C2
B	835	ARG	-	CLONING ARTIFACT	UNP Q9Y6C2
B	836	GLY	-	CLONING ARTIFACT	UNP Q9Y6C2
B	837	SER	-	CLONING ARTIFACT	UNP Q9Y6C2
B	838	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
B	839	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
B	840	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
B	841	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
B	842	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
B	843	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
B	844	GLY	-	CLONING ARTIFACT	UNP Q9Y6C2
B	845	SER	-	CLONING ARTIFACT	UNP Q9Y6C2
C	834	MET	-	CLONING ARTIFACT	UNP Q9Y6C2
C	835	ARG	-	CLONING ARTIFACT	UNP Q9Y6C2
C	836	GLY	-	CLONING ARTIFACT	UNP Q9Y6C2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	837	SER	-	CLONING ARTIFACT	UNP Q9Y6C2
C	838	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
C	839	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
C	840	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
C	841	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
C	842	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
C	843	HIS	-	EXPRESSION TAG	UNP Q9Y6C2
C	844	GLY	-	CLONING ARTIFACT	UNP Q9Y6C2
C	845	SER	-	CLONING ARTIFACT	UNP Q9Y6C2

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: EMILIN-1



• Molecule 1: EMILIN-1



• Molecule 1: EMILIN-1




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1 (medoid)


• Molecule 1: EMILIN-1



• Molecule 1: EMILIN-1




• Molecule 1: EMILIN-1





4.2.2 Score per residue for model 2

• Molecule 1: EMILIN-1




• Molecule 1: EMILIN-1





• Molecule 1: EMILIN-1

Chain C: 70% 9% 19%



4.2.3 Score per residue for model 3

• Molecule 1: EMILIN-1

Chain A: 70% 10% 19%




• Molecule 1: EMILIN-1

Chain B: 69% 10% 19%




• Molecule 1: EMILIN-1

Chain C: 71% 9% 19%




4.2.4 Score per residue for model 4


• Molecule 1: EMILIN-1



- Molecule 1: EMILIN-1




- Molecule 1: EMILIN-1




4.2.5 Score per residue for model 5

- Molecule 1: EMILIN-1




- Molecule 1: EMILIN-1



- Molecule 1: EMILIN-1


Chain C: 



4.2.6 Score per residue for model 6


- Molecule 1: EMILIN-1

Chain A: 




- Molecule 1: EMILIN-1

Chain B: 



- Molecule 1: EMILIN-1

Chain C: 



4.2.7 Score per residue for model 7

- Molecule 1: EMILIN-1

Chain A: 

- Molecule 1: EMILIN-1



- Molecule 1: EMILIN-1



4.2.8 Score per residue for model 8


- Molecule 1: EMILIN-1



- Molecule 1: EMILIN-1




- Molecule 1: EMILIN-1




4.2.9 Score per residue for model 9


- Molecule 1: EMILIN-1



- Molecule 1: EMILIN-1




- Molecule 1: EMILIN-1




4.2.10 Score per residue for model 10

- Molecule 1: EMILIN-1





• Molecule 1: EMILIN-1



• Molecule 1: EMILIN-1



5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *simulated annealing, restrained energy minimization*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	2.9.9
NAMD	refinement	2.6b2

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.93±0.00	0±0/994 (0.0±0.0%)	1.08±0.01	1±1/1356 (0.1±0.0%)
1	B	0.93±0.00	0±0/994 (0.0±0.0%)	1.08±0.01	1±0/1356 (0.1±0.0%)
1	C	0.93±0.00	0±0/994 (0.0±0.0%)	1.07±0.01	1±1/1356 (0.1±0.0%)
All	All	0.93	0/29820 (0.0%)	1.08	39/40680 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.5±0.7
1	B	0.0±0.0	2.5±0.9
1	C	0.0±0.0	2.8±0.7
All	All	0	68

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	906	GLU	N-CA-CB	5.95	121.31	110.60	3	10
1	A	874	ASN	N-CA-CB	5.92	121.25	110.60	2	1
1	A	906	GLU	N-CA-CB	5.91	121.23	110.60	4	10
1	C	906	GLU	N-CA-CB	5.88	121.18	110.60	2	8
1	B	874	ASN	N-CA-CB	5.84	121.11	110.60	2	1
1	B	913	SER	N-CA-CB	5.32	118.48	110.50	10	1
1	C	913	SER	N-CA-CB	5.23	118.35	110.50	9	3
1	C	892	GLY	N-CA-C	-5.13	100.27	113.10	1	2
1	A	886	PHE	CB-CG-CD2	-5.07	117.25	120.80	10	1
1	A	913	SER	N-CA-CB	5.06	118.09	110.50	10	1
1	A	892	GLY	N-CA-C	-5.04	100.50	113.10	6	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	C	893	ARG	Sidechain	10
1	A	921	ARG	Sidechain	9
1	B	921	ARG	Sidechain	9
1	B	893	ARG	Sidechain	8
1	B	879	TYR	Sidechain	6
1	C	921	ARG	Sidechain	6
1	C	894	TYR	Sidechain	5
1	C	879	TYR	Sidechain	4
1	A	893	ARG	Sidechain	4
1	B	914	ARG	Sidechain	2
1	C	860	ARG	Sidechain	2
1	A	879	TYR	Sidechain	1
1	A	914	ARG	Sidechain	1
1	C	914	ARG	Sidechain	1

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	C	973	962	961	0±0
All	All	29190	28860	28830	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:C:954:LEU:HB2	1:C:956:LEU:HD13	0.43	1.91	10	1
1:C:856:LEU:HD11	1:C:858:LEU:HD23	0.41	1.92	7	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/162 (78%)	94±1 (74±1%)	22±2 (17±2%)	11±1 (9±1%)	2	12
1	B	127/162 (78%)	95±1 (75±1%)	21±1 (16±1%)	12±1 (9±1%)	2	12
1	C	127/162 (78%)	93±1 (73±1%)	23±1 (18±1%)	11±1 (9±1%)	2	13
All	All	3810/4860 (78%)	2820 (74%)	651 (17%)	339 (9%)	2	13

All 43 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	861	SER	10
1	B	858	LEU	10
1	C	906	GLU	10
1	A	919	VAL	10
1	A	906	GLU	10
1	A	907	LYS	10
1	C	858	LEU	10
1	A	967	VAL	10
1	B	874	ASN	10
1	C	861	SER	10
1	C	878	TYR	10
1	B	867	PRO	10
1	B	907	LYS	10
1	A	990	PRO	10
1	B	967	VAL	10
1	B	861	SER	10
1	B	906	GLU	10
1	C	990	PRO	10
1	C	967	VAL	10
1	B	990	PRO	10
1	C	874	ASN	10
1	A	858	LEU	10
1	B	919	VAL	10
1	A	867	PRO	10
1	A	874	ASN	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	C	867	PRO	10
1	B	878	TYR	9
1	C	919	VAL	9
1	A	878	TYR	9
1	B	859	PRO	9
1	A	859	PRO	8
1	C	907	LYS	8
1	C	859	PRO	6
1	B	917	GLN	5
1	A	915	SER	3
1	C	915	SER	2
1	B	915	SER	2
1	A	917	GLN	2
1	C	953	ILE	2
1	C	903	HIS	2
1	A	857	SER	1
1	B	857	SER	1
1	A	953	ILE	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	105/131 (80%)	100±1 (95±1%)	6±1 (5±1%)	33	78
1	B	105/131 (80%)	100±1 (95±1%)	6±1 (5±1%)	33	78
1	C	105/131 (80%)	100±1 (95±1%)	5±1 (5±1%)	38	81
All	All	3150/3930 (80%)	2992 (95%)	158 (5%)	35	79

All 35 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	976	GLU	10
1	A	976	GLU	10
1	B	914	ARG	10
1	A	954	LEU	10
1	C	954	LEU	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	B	954	LEU	10
1	C	914	ARG	10
1	A	914	ARG	10
1	A	886	PHE	8
1	B	960	ASP	6
1	B	886	PHE	6
1	C	862	GLU	5
1	A	960	ASP	5
1	C	886	PHE	5
1	B	894	TYR	4
1	B	966	LEU	4
1	C	976	GLU	4
1	C	966	LEU	4
1	A	966	LEU	4
1	A	894	TYR	3
1	C	960	ASP	2
1	B	913	SER	2
1	C	879	TYR	2
1	C	913	SER	2
1	A	913	SER	2
1	B	858	LEU	1
1	C	906	GLU	1
1	B	975	GLU	1
1	C	975	GLU	1
1	A	975	GLU	1
1	C	957	GLN	1
1	B	866	VAL	1
1	C	989	ASP	1
1	A	866	VAL	1
1	A	858	LEU	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided