



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 11:14 PM BST

PDB ID : 2KAX
Title : Solution structure and dynamics of S100A5 in the apo and Ca²⁺-bound states
Authors : Bertini, I.; Das Gupta, S.; Hu, X.; Karavelas, T.; Luchinat, C.; Parigi, G.; Yuan, J.; Structural Proteomics in Europe (SPINE)
Deposited on : 2008-11-17

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 89%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	92	<div> <div>80%</div> <div>10%</div> <div>10%</div> </div>
1	B	92	<div> <div>83%</div> <div>9%</div> <div>9%</div> </div>

2 Ensemble composition and analysis ⓘ

This entry contains 31 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:2-A:43, A:48-A:88, B:2-B:85 (167)	0.68	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 3, 4, 5, 6, 7, 10, 11, 12, 13, 14, 15, 17, 18, 21, 22, 24, 26, 27, 29, 30, 31
2	2, 28
3	8, 25
4	9, 23
5	16, 20
Single-model clusters	19

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 2998 atoms, of which 1502 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Protein S100-A5.


Mol	Chain	Residues	Atoms						Trace
1	A	92	Total	C	H	N	O	S	0
			1499	473	751	116	151	8	
1	B	92	Total	C	H	N	O	S	0
			1499	473	751	116	151	8	

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Protein S100-A5



- Molecule 1: Protein S100-A5




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.2 Score per residue for model 2

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.3 Score per residue for model 3

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.4 Score per residue for model 4

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.5 Score per residue for model 5

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.6 Score per residue for model 6

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.7 Score per residue for model 7

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.8 Score per residue for model 8

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.9 Score per residue for model 9

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.10 Score per residue for model 10

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.11 Score per residue for model 11

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.12 Score per residue for model 12

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.13 Score per residue for model 13

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.14 Score per residue for model 14

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.15 Score per residue for model 15

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.16 Score per residue for model 16

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5



4.2.17 Score per residue for model 17

- Molecule 1: Protein S100-A5

Chain A: 



- Molecule 1: Protein S100-A5


Chain B: 



4.2.18 Score per residue for model 18

- Molecule 1: Protein S100-A5

Chain A: 



- Molecule 1: Protein S100-A5


Chain B: 



4.2.19 Score per residue for model 19

- Molecule 1: Protein S100-A5

Chain A: 




- Molecule 1: Protein S100-A5

Chain B: 




4.2.20 Score per residue for model 20

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.21 Score per residue for model 21

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.22 Score per residue for model 22

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.23 Score per residue for model 23

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.24 Score per residue for model 24

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.25 Score per residue for model 25

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.26 Score per residue for model 26

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.27 Score per residue for model 27

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.28 Score per residue for model 28

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.29 Score per residue for model 29

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.30 Score per residue for model 30

- Molecule 1: Protein S100-A5




- Molecule 1: Protein S100-A5




4.2.31 Score per residue for model 31

- Molecule 1: Protein S100-A5



- Molecule 1: Protein S100-A5



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 350 calculated structures, 31 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
AMBER	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 16033
Number of chemical shift lists	1
Total number of shifts	2320
Number of shifts mapped to atoms	2320
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	89%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.68±0.01	0±0/686 (0.0±0.0%)	1.10±0.04	2±1/917 (0.2±0.1%)
1	B	0.65±0.01	0±0/683 (0.0±0.0%)	1.10±0.04	1±1/912 (0.1±0.1%)
All	All	0.66	0/42439 (0.0%)	1.10	85/56699 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.7±1.2
1	B	0.0±0.0	1.8±1.2
All	All	0	108

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	22	ARG	NE-CZ-NH1	11.71	126.16	120.30	8	11
1	A	22	ARG	NE-CZ-NH1	11.13	125.86	120.30	21	15
1	A	31	ARG	NE-CZ-NH1	9.85	125.22	120.30	5	16
1	A	22	ARG	NE-CZ-NH2	-9.57	115.52	120.30	21	2
1	B	31	ARG	NE-CZ-NH1	8.16	124.38	120.30	1	17
1	A	22	ARG	CD-NE-CZ	7.49	134.09	123.60	21	3
1	B	22	ARG	CD-NE-CZ	7.35	133.89	123.60	8	4
1	B	72	TYR	CB-CG-CD1	-6.18	117.29	121.00	13	1
1	A	31	ARG	N-CA-CB	-5.94	99.91	110.60	17	1
1	B	22	ARG	NE-CZ-NH2	-5.85	117.38	120.30	8	1
1	B	19	TYR	CB-CG-CD1	-5.73	117.56	121.00	8	1
1	B	21	GLY	C-N-CA	5.73	136.02	121.70	19	1
1	A	31	ARG	CD-NE-CZ	5.66	131.52	123.60	14	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	83	TYR	CB-CG-CD2	-5.63	117.62	121.00	27	1
1	A	20	SER	C-N-CA	5.61	134.08	122.30	8	1
1	A	6	GLU	CA-CB-CG	5.45	125.40	113.40	20	2
1	A	72	TYR	CB-CG-CD2	-5.34	117.79	121.00	20	1
1	B	31	ARG	NE-CZ-NH2	5.33	122.97	120.30	5	1
1	A	75	PHE	CB-CG-CD2	-5.31	117.08	120.80	1	1
1	A	31	ARG	NE-CZ-NH2	-5.17	117.72	120.30	28	2
1	B	48	LYS	N-CA-CB	-5.12	101.38	110.60	21	1
1	A	43	CYS	C-N-CA	5.05	134.33	121.70	15	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	72	TYR	Sidechain	14
1	B	62	ASN	Peptide	10
1	B	22	ARG	Sidechain,Peptide	8
1	A	22	ARG	Sidechain	7
1	B	72	TYR	Sidechain	7
1	A	31	ARG	Sidechain	5
1	A	62	ASN	Peptide	5
1	A	41	GLU	Peptide	4
1	B	31	ARG	Sidechain	4
1	B	36	GLU	Sidechain	3
1	A	19	TYR	Sidechain	3
1	B	84	ASN	Peptide	3
1	B	64	ASP	Peptide	3
1	A	64	ASP	Peptide	3
1	B	23	GLU	Peptide	2
1	B	83	TYR	Sidechain	2
1	A	63	SER	Peptide	2
1	B	48	LYS	Peptide	2
1	A	83	TYR	Sidechain	2
1	B	39	LYS	Peptide	1
1	A	48	LYS	Peptide	1
1	B	45	GLY	Peptide	1
1	B	53	ASP	Peptide	1
1	A	23	GLU	Peptide	1
1	B	60	ASP	Peptide	1
1	B	44	LEU	Peptide	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	B	25	SER	Peptide	1
1	A	60	ASP	Peptide	1
1	A	49	GLU	Peptide	1
1	A	3	THR	Peptide	1
1	B	19	TYR	Sidechain	1
1	B	7	LYS	Peptide	1
1	B	63	SER	Peptide	1
1	A	2	GLU	Peptide	1
1	B	41	GLU	Peptide	1
1	B	43	CYS	Peptide	1
1	A	25	SER	Peptide	1

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	675	685	687	1±1
1	A	676	686	687	1±1
All	All	41881	42501	42594	64

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:34:LEU:HD11	1:A:72:TYR:CD1	0.69	2.23	8	2
1:B:19:TYR:CE1	1:B:37:LEU:HD13	0.64	2.28	13	7
1:B:19:TYR:CD1	1:B:37:LEU:HD13	0.57	2.33	8	5
1:A:19:TYR:CD1	1:A:37:LEU:HD13	0.55	2.36	2	3
1:A:5:LEU:O	1:A:5:LEU:HD13	0.55	2.02	5	1
1:A:48:LYS:HE3	1:A:83:TYR:CD2	0.53	2.37	2	1
1:A:48:LYS:HD3	1:A:48:LYS:H	0.53	1.64	14	1
1:A:19:TYR:CE1	1:A:37:LEU:HD13	0.52	2.38	19	4
1:B:48:LYS:HE3	1:B:83:TYR:CD2	0.49	2.42	14	2
1:A:69:PHE:CD2	1:B:77:THR:HG23	0.48	2.44	23	2
1:A:61:LYS:HE3	1:A:61:LYS:HA	0.48	1.86	19	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:88:LEU:HD11	1:B:27:LEU:HD21	0.48	1.86	27	1
1:B:46:GLU:CD	1:B:46:GLU:H	0.48	2.12	16	1
1:B:27:LEU:HD12	1:B:27:LEU:H	0.48	1.69	20	1
1:B:19:TYR:CD2	1:B:37:LEU:HD13	0.47	2.44	29	1
1:A:34:LEU:HD12	1:A:37:LEU:HD23	0.47	1.87	8	1
1:B:22:ARG:HH21	1:B:33:GLU:CD	0.46	2.13	7	1
1:A:48:LYS:HE2	1:A:86:PHE:CZ	0.46	2.46	22	1
1:B:67:ILE:HG23	1:B:71:GLU:HB2	0.46	1.86	26	1
1:B:46:GLU:H	1:B:46:GLU:CD	0.45	2.14	20	1
1:A:70:LYS:HA	1:B:77:THR:HG21	0.45	1.88	26	1
1:B:19:TYR:CE2	1:B:37:LEU:HD13	0.45	2.46	24	2
1:B:38:ILE:HD13	1:B:75:PHE:CE1	0.45	2.46	21	2
1:B:68:ASP:OD1	1:B:70:LYS:HE3	0.44	2.12	25	1
1:A:19:TYR:CE2	1:A:37:LEU:HD13	0.44	2.48	20	3
1:B:61:LYS:HA	1:B:61:LYS:HE3	0.43	1.91	14	1
1:A:72:TYR:CE1	1:A:76:LEU:HD11	0.43	2.48	11	1
1:B:27:LEU:HD23	1:B:27:LEU:H	0.43	1.73	30	1
1:A:31:ARG:HH11	1:A:67:ILE:CG2	0.43	2.27	29	1
1:A:67:ILE:HG22	1:A:71:GLU:CB	0.42	2.44	16	1
1:A:49:GLU:CD	1:A:49:GLU:H	0.42	2.18	5	1
1:A:31:ARG:CZ	1:A:67:ILE:HG23	0.42	2.44	16	1
1:B:26:LYS:H	1:B:26:LYS:HE3	0.41	1.76	5	1
1:A:76:LEU:HD13	1:B:5:LEU:HD21	0.41	1.92	11	1
1:B:47:MET:O	1:B:48:LYS:HE2	0.41	2.15	22	1
1:A:16:PHE:CE2	1:A:29:LEU:HD22	0.41	2.51	11	1
1:B:5:LEU:HD23	1:B:5:LEU:O	0.41	2.16	12	1
1:B:48:LYS:HE3	1:B:83:TYR:CG	0.41	2.50	4	1
1:B:19:TYR:CE2	1:B:37:LEU:HD21	0.40	2.51	23	1
1:A:77:THR:HG21	1:B:70:LYS:HA	0.40	1.92	8	1
1:A:5:LEU:HD12	1:B:42:LEU:HD12	0.40	1.93	5	1
1:B:2:GLU:N	1:B:2:GLU:CD	0.40	2.75	28	1

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	83/92 (90%)	71±2 (86±2%)	10±2 (12±2%)	2±1 (3±1%)	11	47
1	B	84/92 (91%)	71±3 (85±3%)	10±3 (12±3%)	3±1 (3±2%)	8	38
All	All	5177/5704 (91%)	4411 (85%)	610 (12%)	156 (3%)	9	42

All 46 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	50	SER	11
1	B	44	LEU	9
1	B	46	GLU	8
1	A	50	SER	7
1	A	63	SER	7
1	A	2	GLU	6
1	A	83	TYR	6
1	B	45	GLY	5
1	A	43	CYS	5
1	B	47	MET	5
1	A	23	GLU	5
1	B	63	SER	5
1	B	43	CYS	4
1	A	61	LYS	4
1	B	2	GLU	4
1	B	49	GLU	4
1	B	51	SER	4
1	B	22	ARG	3
1	B	26	LYS	3
1	B	4	PRO	3
1	B	24	GLY	3
1	A	69	PHE	3
1	B	83	TYR	3
1	A	49	GLU	3
1	B	69	PHE	3
1	A	84	ASN	3
1	B	85	ASP	3
1	B	25	SER	2
1	A	88	LEU	2
1	A	51	SER	2
1	A	20	SER	2
1	A	24	GLY	2
1	A	22	ARG	2
1	A	25	SER	2
1	B	20	SER	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	21	GLY	1
1	B	61	LYS	1
1	A	62	ASN	1
1	A	87	PHE	1
1	B	3	THR	1
1	A	86	PHE	1
1	B	21	GLY	1
1	B	48	LYS	1
1	A	3	THR	1
1	B	82	ALA	1
1	A	85	ASP	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	79/87 (91%)	67±3 (85±3%)	12±3 (15±3%)	8	47
1	B	79/87 (91%)	68±3 (86±4%)	11±3 (14±4%)	8	49
All	All	4898/5394 (91%)	4202 (86%)	696 (14%)	8	48

All 119 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	42	LEU	23
1	A	61	LYS	18
1	B	61	LYS	17
1	B	46	GLU	16
1	B	57	LYS	16
1	A	3	THR	15
1	A	49	GLU	15
1	A	48	LYS	15
1	B	68	ASP	14
1	B	31	ARG	14
1	B	48	LYS	14
1	A	60	ASP	13
1	A	79	LEU	12

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	B	42	LEU	12
1	A	65	GLN	12
1	B	29	LEU	11
1	B	3	THR	11
1	B	35	LYS	11
1	B	22	ARG	11
1	A	39	LYS	10
1	B	79	LEU	10
1	A	7	LYS	10
1	A	35	LYS	10
1	A	31	ARG	9
1	A	18	LYS	9
1	B	49	GLU	9
1	A	88	LEU	9
1	B	26	LYS	9
1	B	39	LYS	9
1	A	63	SER	9
1	B	69	PHE	9
1	B	65	GLN	9
1	B	67	ILE	9
1	B	7	LYS	9
1	A	81	MET	8
1	A	68	ASP	8
1	A	41	GLU	8
1	B	41	GLU	8
1	A	57	LYS	8
1	A	26	LYS	8
1	A	52	ILE	8
1	B	6	GLU	7
1	B	18	LYS	7
1	A	27	LEU	7
1	B	60	ASP	7
1	A	29	LEU	7
1	B	44	LEU	7
1	A	2	GLU	6
1	B	23	GLU	6
1	A	53	ASP	6
1	A	32	LYS	6
1	A	6	GLU	6
1	A	67	ILE	6
1	B	32	LYS	5
1	A	85	ASP	5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	22	ARG	5
1	A	62	ASN	5
1	A	70	LYS	5
1	A	30	SER	5
1	B	10	THR	5
1	A	58	SER	4
1	A	50	SER	4
1	B	30	SER	4
1	B	47	MET	4
1	B	33	GLU	4
1	B	52	ILE	4
1	A	51	SER	3
1	A	37	LEU	3
1	B	53	ASP	3
1	B	63	SER	3
1	B	85	ASP	3
1	B	2	GLU	3
1	A	10	THR	3
1	B	27	LEU	3
1	A	23	GLU	3
1	A	69	PHE	3
1	B	66	GLU	3
1	B	40	LYS	3
1	A	43	CYS	3
1	B	64	ASP	3
1	B	36	GLU	2
1	B	56	MET	2
1	A	71	GLU	2
1	B	15	THR	2
1	A	9	LEU	2
1	A	83	TYR	2
1	B	58	SER	2
1	B	17	HIS	2
1	A	11	THR	2
1	A	33	GLU	2
1	A	56	MET	2
1	A	5	LEU	2
1	A	25	SER	2
1	B	50	SER	1
1	B	5	LEU	1
1	A	84	ASN	1
1	A	64	ASP	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	B	20	SER	1
1	B	12	MET	1
1	B	84	ASN	1
1	B	62	ASN	1
1	B	28	THR	1
1	B	51	SER	1
1	A	34	LEU	1
1	B	81	MET	1
1	B	43	CYS	1
1	A	66	GLU	1
1	A	87	PHE	1
1	A	36	GLU	1
1	B	83	TYR	1
1	A	80	CYS	1
1	B	70	LYS	1
1	B	14	THR	1
1	B	25	SER	1
1	A	75	PHE	1
1	B	37	LEU	1
1	A	54	ASP	1
1	A	17	HIS	1
1	A	40	LYS	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 89% for the well-defined parts and 89% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 16033

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	2320
Number of shifts mapped to atoms	2320
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	184	-2.52 ± 0.16	Should be applied
$^{13}\text{C}_\beta$	178	-1.47 ± 0.06	Should be applied
$^{13}\text{C}'$	174	1.33 ± 0.14	Should be applied
^{15}N	178	0.44 ± 0.25	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 89%, i.e. 1885 atoms were assigned a chemical shift out of a possible 2110. 0 out of 28 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	819/831 (99%)	330/332 (99%)	326/334 (98%)	163/165 (99%)
Sidechain	1010/1143 (88%)	631/671 (94%)	375/432 (87%)	4/40 (10%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	56/136 (41%)	56/72 (78%)	0/60 (0%)	0/4 (0%)
Overall	1885/2110 (89%)	1017/1075 (95%)	701/826 (85%)	167/209 (80%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 89%, i.e. 2080 atoms were assigned a chemical shift out of a possible 2330. 0 out of 30 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	898/916 (98%)	362/366 (99%)	358/368 (97%)	178/182 (98%)
Sidechain	1118/1260 (89%)	700/740 (95%)	412/476 (87%)	6/44 (14%)
Aromatic	64/154 (42%)	64/82 (78%)	0/68 (0%)	0/4 (0%)
Overall	2080/2330 (89%)	1126/1188 (95%)	770/912 (84%)	184/230 (80%)


7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The images below report *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:

