



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 11, 2017 – 06:47 AM EST

PDB ID : 5KA6  
Title : HIV-1 gp41 variant Q552R and L555M resistance mutations  
Authors : Bhardwaj, A.; Khasnis, M.D.; Halkidis, K.; Root, M.J.  
Deposited on : 2016-06-01  
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	91	<div> <div>3%</div> <div> <div></div> <div>78%</div> <div>9%</div> <div>13%</div> </div> </div>
1	B	91	<div> <div>5%</div> <div> <div></div> <div>73%</div> <div>14%</div> <div>13%</div> </div> </div>
1	C	91	<div> <div>3%</div> <div> <div></div> <div>68%</div> <div>16%</div> <div>15%</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2035 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transmembrane protein gp41.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	79	Total	C	N	O	S	0	0	0
			655	408	120	124	3			
1	B	79	Total	C	N	O	S	0	0	0
			651	405	120	124	2			
1	C	77	Total	C	N	O	S	0	0	0
			633	395	115	121	2			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	542	MET	-	initiating methionine	UNP P04578
A	552	ARG	GLN	engineered mutation	UNP P04578
A	555	MET	LEU	engineered mutation	UNP P04578
A	620	GLY	-	linker	UNP P04578
A	621	GLY	-	linker	UNP P04578
A	622	ARG	-	linker	UNP P04578
A	623	GLY	-	linker	UNP P04578
A	624	GLY	-	linker	UNP P04578
A	662	GLY	-	expression tag	UNP P04578
A	663	GLY	-	expression tag	UNP P04578
A	664	HIS	-	expression tag	UNP P04578
A	665	HIS	-	expression tag	UNP P04578
A	666	HIS	-	expression tag	UNP P04578
A	667	HIS	-	expression tag	UNP P04578
A	668	HIS	-	expression tag	UNP P04578
A	669	HIS	-	expression tag	UNP P04578
B	542	MET	-	initiating methionine	UNP P04578
B	552	ARG	GLN	engineered mutation	UNP P04578
B	555	MET	LEU	engineered mutation	UNP P04578
B	620	GLY	-	linker	UNP P04578
B	621	GLY	-	linker	UNP P04578
B	622	ARG	-	linker	UNP P04578
B	623	GLY	-	linker	UNP P04578

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	624	GLY	-	linker	UNP P04578
B	662	GLY	-	expression tag	UNP P04578
B	663	GLY	-	expression tag	UNP P04578
B	664	HIS	-	expression tag	UNP P04578
B	665	HIS	-	expression tag	UNP P04578
B	666	HIS	-	expression tag	UNP P04578
B	667	HIS	-	expression tag	UNP P04578
B	668	HIS	-	expression tag	UNP P04578
B	669	HIS	-	expression tag	UNP P04578
C	542	MET	-	initiating methionine	UNP P04578
C	552	ARG	GLN	engineered mutation	UNP P04578
C	555	MET	LEU	engineered mutation	UNP P04578
C	620	GLY	-	linker	UNP P04578
C	621	GLY	-	linker	UNP P04578
C	622	ARG	-	linker	UNP P04578
C	623	GLY	-	linker	UNP P04578
C	624	GLY	-	linker	UNP P04578
C	662	GLY	-	expression tag	UNP P04578
C	663	GLY	-	expression tag	UNP P04578
C	664	HIS	-	expression tag	UNP P04578
C	665	HIS	-	expression tag	UNP P04578
C	666	HIS	-	expression tag	UNP P04578
C	667	HIS	-	expression tag	UNP P04578
C	668	HIS	-	expression tag	UNP P04578
C	669	HIS	-	expression tag	UNP P04578



- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	39	Total O 39 39	0	0
2	B	30	Total O 30 30	0	0
2	C	27	Total O 27 27	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transmembrane protein gp41



- Molecule 1: Transmembrane protein gp41



- Molecule 1: Transmembrane protein gp41



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	43.59Å 43.59Å 113.05Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.81 – 1.85 35.81 – 1.85	Depositor EDS
% Data completeness (in resolution range)	96.9 (35.81-1.85) 83.0 (35.81-1.85)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.54 (at 1.85Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.231 , 0.278 0.214 , 0.261	Depositor DCC
$R_{free}$ test set	1692 reflections (9.75%)	DCC
Wilson B-factor (Å <sup>2</sup> )	26.6	Xtriage
Anisotropy	0.251	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 50.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l 0.073 for h,-h-k,-l 0.039 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2035	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.28	0/663	0.38	0/893
1	B	0.25	0/659	0.38	0/888
1	C	0.32	0/640	0.43	0/863
All	All	0.28	0/1962	0.40	0/2644

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	655	0	649	8	0
1	B	651	0	643	13	0
1	C	633	0	630	14	0
2	A	39	0	0	4	0
2	B	30	0	0	7	0
2	C	27	0	0	3	0
All	All	2035	0	1922	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 28 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:544:LEU:HG	1:C:545:LEU:HD22	1.59	0.84
1:B:630:GLU:OE2	2:B:701:HOH:O	1.99	0.81
1:C:648:GLU:OE1	2:C:701:HOH:O	1.99	0.80
1:A:550:GLN:OE1	2:A:701:HOH:O	2.03	0.74
1:A:542:MET:N	2:A:705:HOH:O	2.26	0.69

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	75/91 (82%)	75 (100%)	0	0	100	100
1	B	75/91 (82%)	75 (100%)	0	0	100	100
1	C	73/91 (80%)	73 (100%)	0	0	100	100
All	All	223/273 (82%)	223 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	72/79 (91%)	72 (100%)	0	100	100
1	B	71/79 (90%)	71 (100%)	0	100	100
1	C	69/79 (87%)	69 (100%)	0	100	100

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	212/237 (90%)	212 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	550	GLN
1	C	577	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	79/91 (86%)	0.08	3 (3%) 44 41	24, 36, 81, 92	0
1	B	79/91 (86%)	0.16	5 (6%) 23 22	22, 40, 82, 98	0
1	C	77/91 (84%)	0.15	3 (3%) 43 40	25, 42, 75, 101	0
All	All	235/273 (86%)	0.13	11 (4%) 35 33	22, 40, 81, 101	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	663	GLY	11.2
1	C	582	ALA	5.3
1	C	544	LEU	5.3
1	B	624	GLY	3.9
1	A	542	MET	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.