



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 2, 2017 – 04:40 PM EST

PDB ID : 5KDJ
Title : ZmpB metallopeptidase from *Clostridium perfringens*
Authors : Noach, I.; Ficko-Blean, E.; Stuart, C.; Boraston, A.B.
Deposited on : 2016-06-08
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028442
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	674	<div> <div>10%</div> <div>69%</div> <div>5%</div> <div>26%</div> </div>
1	B	674	<div> <div>%</div> <div>69%</div> <div>.</div> <div>26%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	B	1101	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8827 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called F5/8 type C domain protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	501	Total	C	N	O	S	0	0	0
			4019	2552	675	782	10			
1	A	498	Total	C	N	O	S	0	3	0
			4019	2555	672	782	10			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	411	MET	-	initiating methionine	UNP A0A0H2YN38
B	412	GLY	-	expression tag	UNP A0A0H2YN38
B	413	SER	-	expression tag	UNP A0A0H2YN38
B	414	SER	-	expression tag	UNP A0A0H2YN38
B	415	HIS	-	expression tag	UNP A0A0H2YN38
B	416	HIS	-	expression tag	UNP A0A0H2YN38
B	417	HIS	-	expression tag	UNP A0A0H2YN38
B	418	HIS	-	expression tag	UNP A0A0H2YN38
B	419	HIS	-	expression tag	UNP A0A0H2YN38
B	420	HIS	-	expression tag	UNP A0A0H2YN38
B	421	SER	-	expression tag	UNP A0A0H2YN38
B	422	SER	-	expression tag	UNP A0A0H2YN38
B	423	GLY	-	expression tag	UNP A0A0H2YN38
B	424	LEU	-	expression tag	UNP A0A0H2YN38
B	425	VAL	-	expression tag	UNP A0A0H2YN38
B	426	PRO	-	expression tag	UNP A0A0H2YN38
B	427	ARG	-	expression tag	UNP A0A0H2YN38
B	428	GLY	-	expression tag	UNP A0A0H2YN38
B	429	SER	-	expression tag	UNP A0A0H2YN38
B	430	HIS	-	expression tag	UNP A0A0H2YN38
B	431	MET	-	expression tag	UNP A0A0H2YN38
B	432	ALA	-	expression tag	UNP A0A0H2YN38
B	433	SER	-	expression tag	UNP A0A0H2YN38
A	411	MET	-	initiating methionine	UNP A0A0H2YN38
A	412	GLY	-	expression tag	UNP A0A0H2YN38

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	413	SER	-	expression tag	UNP A0A0H2YN38
A	414	SER	-	expression tag	UNP A0A0H2YN38
A	415	HIS	-	expression tag	UNP A0A0H2YN38
A	416	HIS	-	expression tag	UNP A0A0H2YN38
A	417	HIS	-	expression tag	UNP A0A0H2YN38
A	418	HIS	-	expression tag	UNP A0A0H2YN38
A	419	HIS	-	expression tag	UNP A0A0H2YN38
A	420	HIS	-	expression tag	UNP A0A0H2YN38
A	421	SER	-	expression tag	UNP A0A0H2YN38
A	422	SER	-	expression tag	UNP A0A0H2YN38
A	423	GLY	-	expression tag	UNP A0A0H2YN38
A	424	LEU	-	expression tag	UNP A0A0H2YN38
A	425	VAL	-	expression tag	UNP A0A0H2YN38
A	426	PRO	-	expression tag	UNP A0A0H2YN38
A	427	ARG	-	expression tag	UNP A0A0H2YN38
A	428	GLY	-	expression tag	UNP A0A0H2YN38
A	429	SER	-	expression tag	UNP A0A0H2YN38
A	430	HIS	-	expression tag	UNP A0A0H2YN38
A	431	MET	-	expression tag	UNP A0A0H2YN38
A	432	ALA	-	expression tag	UNP A0A0H2YN38
A	433	SER	-	expression tag	UNP A0A0H2YN38

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		
4	A	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	472	Total	O	0	2
			474	474		
5	A	300	Total	O	0	0
			300	300		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: F5/8 type C domain protein



- Molecule 1: F5/8 type C domain protein



GLU	VAL	LEU	TYR	SER	LEU	GLU	ASP	SER	SER	LYS	GLY	ASN	GLU	TYR	THR	LYS	THR	SER	SER	LYS	VAL	ASN	VAL	VAL	SER	ARG	LYS
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.62Å 95.80Å 187.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.82 – 2.15 29.82 – 2.15	Depositor EDS
% Data completeness (in resolution range)	96.2 (29.82-2.15) 96.2 (29.82-2.15)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.61 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.191 , 0.229 0.191 , 0.229	Depositor DCC
R_{free} test set	3170 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.380	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8827	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	3/4118 (0.1%)	0.73	4/5568 (0.1%)
1	B	0.65	1/4109 (0.0%)	0.73	4/5554 (0.1%)
All	All	0.64	4/8227 (0.0%)	0.73	8/11122 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	673	TRP	CD2-CE2	5.87	1.48	1.41
1	A	752	TRP	CD2-CE2	5.51	1.48	1.41
1	B	797	TRP	CD2-CE2	5.41	1.47	1.41
1	A	699	TRP	CD2-CE2	5.02	1.47	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	689[A]	ASP	CB-CG-OD1	10.22	127.50	118.30
1	A	689[B]	ASP	CB-CG-OD1	10.22	127.50	118.30
1	B	529	ARG	NE-CZ-NH1	7.88	124.24	120.30
1	B	594	ASP	CB-CG-OD1	6.14	123.83	118.30
1	A	594	ASP	CB-CG-OD1	6.11	123.80	118.30
1	B	529	ARG	NE-CZ-NH2	-6.01	117.30	120.30
1	B	696	MET	CA-CB-CG	-5.49	103.96	113.30
1	A	743	ASP	CB-CG-OD1	5.04	122.84	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4019	0	3901	20	0
1	B	4019	0	3895	16	0
2	A	6	0	8	3	0
2	B	6	0	8	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	300	0	0	3	0
5	B	474	0	0	5	0
All	All	8827	0	7812	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:666:GLN:HE22	2:A:1101:GOL:H31	1.32	0.93
1:A:826:GLN:HE22	1:A:858:ARG:HH11	1.32	0.75
1:A:756:HIS:HD1	1:A:775:ASN:HD21	1.33	0.74
1:B:756:HIS:HD1	1:B:775:ASN:HD21	1.37	0.72
1:B:496:ASP:N	5:B:1201:HOH:O	2.23	0.71
1:A:826:GLN:NE2	1:A:858:ARG:HH11	1.97	0.63
1:B:808:ASP:OD1	1:B:958:ARG:NH1	2.37	0.57
1:A:826:GLN:NE2	1:A:862:TYR:OH	2.41	0.54
1:A:666:GLN:NE2	2:A:1101:GOL:H31	2.12	0.54
1:A:808:ASP:OD1	1:A:958:ARG:NH1	2.41	0.53
1:A:618:LYS:NZ	5:A:1206:HOH:O	2.42	0.53
1:B:893:LYS:NZ	5:B:1204:HOH:O	2.36	0.52
1:A:666:GLN:HE22	2:A:1101:GOL:C3	2.16	0.51
1:A:496:ASP:N	5:A:1207:HOH:O	2.45	0.49
1:B:705:LYS:HE2	5:B:1617:HOH:O	2.13	0.48
1:A:942:PHE:HZ	1:A:957:ILE:HD11	1.78	0.47
1:B:756:HIS:CE1	1:B:774:ASN:HD21	2.33	0.46
1:A:622:PHE:CG	1:A:681:PRO:HB2	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:756:HIS:HD1	1:B:775:ASN:ND2	2.09	0.45
1:B:942:PHE:HZ	1:B:957:ILE:HD11	1.81	0.45
1:B:622:PHE:CG	1:B:681:PRO:HB2	2.52	0.44
1:B:912:LEU:O	1:B:992:ARG:HD3	2.18	0.44
1:A:552:LYS:O	1:A:593:GLY:HA3	2.19	0.43
1:A:828:ALA:O	1:A:832:GLN:HG3	2.18	0.43
1:B:815:LEU:HA	1:B:884:HIS:O	2.19	0.43
1:A:815:LEU:HA	1:A:884:HIS:O	2.18	0.42
1:A:756:HIS:HD1	1:A:775:ASN:ND2	2.09	0.42
1:B:529:ARG:HD3	5:B:1293:HOH:O	2.20	0.42
1:B:676:LYS:HD2	5:B:1536:HOH:O	2.20	0.42
1:B:818:LYS:HE2	1:B:818:LYS:HB3	1.93	0.41
1:A:756:HIS:CE1	1:A:774:ASN:HD21	2.38	0.41
1:A:676:LYS:HE3	1:A:676:LYS:HB2	1.92	0.40
1:B:828:ALA:N	1:B:829:PRO:HD2	2.36	0.40
1:B:713:PHE:HE1	1:B:715:ILE:HG12	1.86	0.40
1:A:754:VAL:HG23	5:A:1291:HOH:O	2.20	0.40
1:A:895:ASP:O	1:A:898:LYS:HG2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	495/674 (73%)	482 (97%)	13 (3%)	0	100	100
1	B	495/674 (73%)	482 (97%)	13 (3%)	0	100	100
All	All	990/1348 (73%)	964 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	435/591 (74%)	428 (98%)	7 (2%)	70	76
1	B	433/591 (73%)	424 (98%)	9 (2%)	61	65
All	All	868/1182 (73%)	852 (98%)	16 (2%)	66	71

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	496	ASP
1	B	594	ASP
1	B	691	TYR
1	B	699	TRP
1	B	795	ASN
1	B	901	LYS
1	B	946	ASP
1	B	984	VAL
1	B	1003	ASN
1	A	587	LYS
1	A	594	ASP
1	A	691	TYR
1	A	699	TRP
1	A	767	ARG
1	A	795	ASN
1	A	1003	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	775	ASN
1	B	924	ASN
1	A	775	ASN
1	A	795	ASN
1	A	826	GLN
1	A	924	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	1101	-	5,5,5	0.31	0	5,5,5	0.39	0
2	GOL	B	1101	-	5,5,5	0.24	0	5,5,5	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1101	-	-	0/4/4/4	0/0/0/0
2	GOL	B	1101	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1101	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	498/674 (73%)	0.62	66 (13%) 4 7	8, 23, 59, 72	0
1	B	501/674 (74%)	0.01	9 (1%) 71 79	8, 15, 32, 49	0
All	All	999/1348 (74%)	0.31	75 (7%) 17 23	8, 18, 51, 72	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	724	GLY	9.4
1	A	725	GLY	8.4
1	B	724	GLY	7.8
1	A	961	GLY	6.3
1	A	1004	ALA	5.9
1	A	959	ARG	5.4
1	A	1003	ASN	5.2
1	A	960	ASP	5.1
1	A	726	ALA	5.1
1	B	725	GLY	4.5
1	A	944	VAL	4.4
1	A	986	VAL	4.2
1	A	931	THR	4.1
1	A	929	VAL	4.0
1	A	984	VAL	3.9
1	A	1000	ILE	3.9
1	A	938	ILE	3.8
1	A	943	SER	3.7
1	A	921	PHE	3.7
1	A	919	ASP	3.7
1	A	932	SER	3.7
1	A	942	PHE	3.5
1	A	922	THR	3.5
1	A	923	GLU	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	962	LYS	3.4
1	A	924	ASN	3.4
1	A	917	LYS	3.3
1	A	892	VAL	3.3
1	A	925	ALA	3.2
1	B	727	PHE	3.2
1	A	926	LYS	3.2
1	A	998	ASN	3.1
1	A	894	GLU	3.0
1	A	928	SER	2.9
1	A	1002	VAL	2.8
1	A	927	VAL	2.7
1	B	1003	ASN	2.7
1	A	581	GLU	2.7
1	B	984	VAL	2.7
1	A	985	TYR	2.6
1	A	889	ASP	2.6
1	A	940	LEU	2.6
1	A	930	SER	2.6
1	A	723	SER	2.6
1	B	1004	ALA	2.5
1	A	891	LYS	2.5
1	B	933	GLY	2.5
1	A	996	THR	2.4
1	A	568	PRO	2.4
1	A	819	ALA	2.4
1	A	900	PRO	2.4
1	A	903	ASP	2.3
1	A	945	ASP	2.3
1	A	727	PHE	2.3
1	A	997	LEU	2.3
1	A	946	ASP	2.2
1	A	975	THR	2.2
1	A	948	ASN	2.2
1	A	886	ILE	2.2
1	A	662	LEU	2.2
1	A	890	ASP	2.2
1	A	854	GLY	2.2
1	A	897	ALA	2.2
1	A	812	ASN	2.2
1	A	587	LYS	2.2
1	B	581	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	937	ASN	2.1
1	A	818	LYS	2.1
1	A	893	LYS	2.1
1	A	878	THR	2.1
1	A	920	GLY	2.1
1	B	662	LEU	2.0
1	A	856	LYS	2.0
1	A	898	LYS	2.0
1	A	877	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GOL	B	1101	6/6	0.87	0.18	4.58	34,38,41,45	0
2	GOL	A	1101	6/6	0.95	0.14	1.55	35,36,37,38	0
4	ZN	A	1102	1/1	0.98	0.06	-	39,39,39,39	0
3	NA	B	1102	1/1	0.97	0.29	-	27,27,27,27	0
4	ZN	B	1103	1/1	0.99	0.04	-	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.