



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2016 – 06:27 PM EDT

PDB ID : 5KFY
Title : Human DNA polymerase eta-DNA ternary complex: reaction first with 1 mM Mn²⁺ for 1800s then with 5 mM Mn²⁺ for 60s at 4 degree
Authors : Gao, Y.; Yang, W.
Deposited on : 2016-06-12
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027790
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027790


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	435	<div> <div>10%</div> <div>87%</div> <div>12%</div> <div>.</div> </div>
2	T	12	<div> <div>8%</div> <div>83%</div> <div>17%</div> </div>
3	P	8	<div> <div>75%</div> <div>25%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	503	-	-	-	X
5	GOL	A	504	-	-	-	X
7	EDO	A	506	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase eta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	430	Total	C	N	O	S	0	19	0
			3433	2152	615	638	28			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q9Y253
A	-1	PRO	-	expression tag	UNP Q9Y253
A	0	HIS	-	expression tag	UNP Q9Y253

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*AP*TP*TP*AP*TP*GP*AP*CP*GP*CP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	12	Total	C	N	O	P	0	2	1
			252	119	41	79	13			

- Molecule 3 is a DNA chain called DNA (5'-D(*AP*GP*CP*GP*TP*CP*AP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	8	Total	C	N	O	P	0	1	0
			166	79	30	49	8			

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mn	0	1
			2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (three-letter code: DTP) (formula: $C_{10}H_{16}N_5O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			30	10	5	12	3		

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	380	Total	O	0	0
			380	380		
8	T	36	Total	O	0	0
			36	36		
8	P	23	Total	O	0	0
			23	23		

4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	98.59 Å 98.59 Å 82.34 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.76 – 1.70 19.76 – 1.70	Depositor EDS
% Data completeness (in resolution range)	93.5 (19.76-1.70) 93.6 (19.76-1.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 1.70 Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.202 , 0.228 0.202 , 0.228	Depositor DCC
R_{free} test set	3628 reflections (7.76%)	DCC
Wilson B-factor (Å ²)	15.2	Xtriage
Anisotropy	0.061	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.055 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4338	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, DTP, EDO, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/3573	0.43	0/4819
2	T	0.51	0/296	0.98	0/456
3	P	0.62	0/201	1.06	0/306
All	All	0.34	0/4070	0.55	0/5581

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3433	0	3503	40	0
2	T	252	0	134	2	0
3	P	166	0	85	2	0
4	A	2	0	0	0	0
5	A	12	0	16	0	0
6	A	30	0	12	1	0
7	A	4	0	6	2	0
8	A	380	0	0	11	1
8	P	23	0	0	0	0
8	T	36	0	0	1	1
All	All	4338	0	3756	43	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:ILE:HA	1:A:188:LEU:HD22	1.74	0.70
1:A:322[B]:SER:HB2	1:A:423[B]:PHE:HD1	1.59	0.68
1:A:382:ARG:NH1	8:A:605:HOH:O	2.31	0.64
1:A:377:ARG:NH1	8:A:608:HOH:O	2.33	0.61
1:A:132:LEU:O	1:A:135:GLN:HG3	2.01	0.60
1:A:13:ASP:OD1	1:A:231[A]:LYS:NZ	2.35	0.58
1:A:81:ARG:NH2	8:A:603:HOH:O	2.28	0.58
1:A:227:CYS:HB3	7:A:506:EDO:H12	1.84	0.58
1:A:369:SER:HB3	1:A:423[A]:PHE:HB3	1.88	0.56
1:A:224[A]:LYS:HD2	7:A:506:EDO:H11	1.91	0.53
1:A:106:PHE:CG	1:A:200[A]:MET:HG2	2.48	0.49
1:A:256[A]:ARG:NH1	8:A:611:HOH:O	2.37	0.49
1:A:365:GLN:OE1	1:A:428:LYS:NZ	2.37	0.49
1:A:61:ARG:NH1	8:A:607:HOH:O	2.34	0.49
1:A:127:GLU:O	1:A:130:GLN:HG2	2.13	0.48
1:A:369:SER:HB3	1:A:423[B]:PHE:HB3	1.93	0.48
1:A:130:GLN:O	1:A:133:GLN:HG2	2.14	0.47
1:A:109:ILE:HD11	1:A:314:GLN:HG3	1.95	0.47
1:A:182:ASN:OD1	1:A:184:THR:HG22	2.15	0.47
1:A:179:GLN:OE1	1:A:179:GLN:HA	2.13	0.47
1:A:253:ARG:HD3	8:A:921:HOH:O	2.15	0.46
1:A:116:GLU:OE1	1:A:224[A]:LYS:NZ	2.45	0.45
1:A:96:SER:O	1:A:100:MET:HG2	2.16	0.45
1:A:256[A]:ARG:HD2	8:A:865:HOH:O	2.17	0.44
1:A:423[B]:PHE:CZ	1:A:425:CYS:HB2	2.53	0.44
2:T:2:DA:N3	8:T:104:HOH:O	2.36	0.44
1:A:180:ILE:HG22	1:A:188:LEU:HD13	1.99	0.43
1:A:0:HIS:HB3	8:A:730:HOH:O	2.16	0.43
1:A:256[B]:ARG:HE	3:P:7:DA:H4'	1.83	0.43
6:A:505:DTP:H8	6:A:505:DTP:O5'	2.19	0.43
1:A:371:ARG:HB3	1:A:421:MET:HB3	2.00	0.42
1:A:375:ASP:OD2	1:A:377:ARG:NE	2.49	0.42
1:A:367:VAL:HG22	1:A:384[B]:CYS:SG	2.59	0.42
1:A:311:LYS:HE2	8:A:634:HOH:O	2.18	0.42
1:A:128:ARG:HD3	1:A:131:LYS:HE3	2.02	0.41
1:A:230[A]:ASN:O	1:A:234:ARG:HB2	2.20	0.41
1:A:317:LYS:HE3	1:A:317:LYS:HB3	1.64	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:ARG:NH2	8:A:610:HOH:O	2.37	0.41
1:A:404:LYS:HA	1:A:415:TRP:HH2	1.86	0.41
1:A:224[A]:LYS:HE2	3:P:8[A]:DT:OP2	2.21	0.41
1:A:103[B]:MET:HG2	1:A:200[B]:MET:SD	2.61	0.41
1:A:177[A]:SER:OG	8:A:601:HOH:O	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:683:HOH:O	8:T:115:HOH:O[6_555]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	446/435 (102%)	438 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	388/372 (104%)	381 (98%)	7 (2%)	66	49

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115	ASP
1	A	181	ASP
1	A	231[A]	LYS
1	A	231[B]	LYS
1	A	372	VAL
1	A	423[A]	PHE
1	A	423[B]	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	A	503	-	5,5,5	0.34	0	5,5,5	0.24	0
5	GOL	A	504	-	5,5,5	0.35	0	5,5,5	0.25	0
6	DTP	A	505	4	25,32,32	1.27	4 (16%)	26,50,50	2.45	1 (3%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	EDO	A	506	-	3,3,3	0.38	0	2,2,2	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	503	-	-	0/4/4/4	0/0/0/0
5	GOL	A	504	-	-	0/4/4/4	0/0/0/0
6	DTP	A	505	4	-	0/18/34/34	0/3/3/3
7	EDO	A	506	-	-	0/1/1/1	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	505	DTP	C3'-C4'	-2.83	1.44	1.53
6	A	505	DTP	C2'-C3'	-2.42	1.46	1.52
6	A	505	DTP	C6-N6	2.32	1.43	1.34
6	A	505	DTP	C2-N3	2.61	1.36	1.32

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	505	DTP	N3-C2-N1	-11.70	119.68	128.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	505	DTP	1	0
7	A	506	EDO	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	430/435 (98%)	0.57	42 (9%) 10 10	8, 19, 56, 79	0
2	T	12/12 (100%)	0.12	1 (8%) 14 16	15, 21, 44, 63	0
3	P	8/8 (100%)	-0.23	0 100 100	14, 24, 29, 32	0
All	All	450/455 (98%)	0.54	43 (9%) 10 11	8, 19, 56, 79	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	160	THR	13.0
1	A	154	PRO	11.7
1	A	161	VAL	10.1
1	A	0	HIS	8.6
1	A	411	ILE	8.5
1	A	412	GLN	8.0
1	A	378	LEU	6.8
1	A	153	GLY	6.4
1	A	-1	PRO	6.0
1	A	410	GLY	5.7
1	A	181	ASP	5.6
1	A	180	ILE	5.3
1	A	376	LYS	5.1
1	A	413	THR	4.3
1	A	2	ALA	4.0
1	A	162	GLN	3.9
1	A	408	THR	3.8
1	A	179	GLN	3.8
1	A	172	PHE	3.6
1	A	130	GLN	3.6
1	A	184	THR	3.5
1	A	183	LEU	3.5
1	A	133	GLN	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	132	LEU	3.5
1	A	131	LYS	3.1
1	A	1	MET	3.0
1	A	417	PRO	3.0
1	A	379	SER	2.9
1	A	136	PRO	2.8
1	A	405	ASN	2.7
2	T	3[A]	DT	2.6
1	A	423[A]	PHE	2.5
1	A	374	GLY	2.5
1	A	182	ASN	2.4
1	A	-2	GLY	2.4
1	A	134	GLY	2.4
1	A	328	LYS	2.4
1	A	416	SER	2.3
1	A	163	LYS	2.3
1	A	381	LEU	2.2
1	A	135	GLN	2.2
1	A	380	SER	2.1
1	A	406	CYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	GOL	A	503	6/6	0.87	0.27	15.51	16,24,29,29	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	GOL	A	504	6/6	0.59	0.32	8.90	27,34,35,40	0
7	EDO	A	506	4/4	0.92	0.24	3.62	8,9,15,16	4
6	DTP	A	505	30/30	0.97	0.07	-0.85	6,10,14,15	0
4	MN	A	502[B]	1/1	1.00	0.04	-1.62	10,10,10,10	0
4	MN	A	501	1/1	1.00	0.05	-2.55	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.