



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 14, 2016 – 07:59 PM EDT

PDB ID : 5KG7  
Title : Human DNA polymerase eta-DNA ternary complex: reaction first with 1 mM Mn<sup>2+</sup> for 1800s then with 10 mM Zn<sup>2+</sup> for 60s  
Authors : Gao, Y.; Yang, W.  
Deposited on : 2016-06-12  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027790  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027790


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1609 (1.76-1.76)
Clashscore	102246	1730 (1.76-1.76)
Ramachandran outliers	100387	1711 (1.76-1.76)
Sidechain outliers	100360	1711 (1.76-1.76)
RSRZ outliers	91569	1610 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	435	 8% 87% 11% ..
2	T	12	 17% 50% 50%
3	P	9	 89% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	503	-	-	-	X
5	GOL	A	504	-	-	-	X
5	GOL	A	505	-	-	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4379 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase eta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	430	Total	C	N	O	S	0	16	0
			3420	2137	622	632	29			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q9Y253
A	-1	PRO	-	expression tag	UNP Q9Y253
A	0	HIS	-	expression tag	UNP Q9Y253

- Molecule 2 is a DNA chain called DNA (5'-D(\*CP\*AP\*TP\*TP\*AP\*TP\*GP\*AP\*CP\*GP\*CP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	12	Total	C	N	O	P	0	0	1
			226	108	39	68	11			


- Molecule 3 is a DNA chain called DNA (5'-D(\*AP\*GP\*CP\*GP\*TP\*CP\*AP\*TP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	9	Total	C	N	O	P	0	4	0
			208	99	37	62	10			

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	P	1	Total	Mn	0	1
			1	1		
4	A	2	Total	Mn	0	1
			2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (three-letter code: DTP) (formula:  $C_{10}H_{16}N_5O_{12}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	1
			30	10	5	12	3		

- Molecule 7 is DIPHOSPHATE (three-letter code: DPO) (formula:  $O_7P_2$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	P	0	1
			9	7	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	400	Total	O	0	4
			401	401		
8	T	36	Total	O	0	0
			36	36		
8	P	28	Total	O	0	0
			28	28		



- Molecule 1: DNA polymerase eta



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.22Å 98.22Å 82.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.96 – 1.75 19.96 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.6 (19.96-1.75) 99.6 (19.96-1.75)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.84 (at 1.74Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.186 , 0.228 0.186 , 0.228	Depositor DCC
$R_{free}$ test set	3489 reflections (7.72%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.3	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 46.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.059 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4379	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.45% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: DPO, GOL, DTP, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	2/3551 (0.1%)	0.64	2/4787 (0.0%)
2	T	0.83	0/252	1.00	0/388
3	P	3.29	2/272 (0.7%)	1.49	2/419 (0.5%)
All	All	1.04	4/4075 (0.1%)	0.77	4/5594 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	8[A]	DT	O3'-P	37.00	2.05	1.61
3	P	8[B]	DT	O3'-P	37.00	2.05	1.61
1	A	312	PRO	N-CD	5.76	1.55	1.47
1	A	14	MET	C-O	-5.08	1.13	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	8[A]	DT	P-O3'-C3'	-15.92	100.60	119.70
3	P	8[B]	DT	P-O3'-C3'	-15.92	100.60	119.70
1	A	61[A]	ARG	CA-CB-CG	8.95	133.09	113.40
1	A	61[B]	ARG	CA-CB-CG	8.95	133.09	113.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3420	0	3488	38	1
2	T	226	0	126	6	0
3	P	208	0	99	2	0
4	A	2	0	0	0	0
4	P	1	0	0	0	0
5	A	18	0	24	1	0
6	A	30	0	11	1	0
7	A	9	0	0	0	0
8	A	401	0	0	6	2
8	P	28	0	0	1	0
8	T	36	0	0	2	0
All	All	4379	0	3748	46	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:P:8[B]:DT:H2"	8:P:205:HOH:O	1.53	1.06
1:A:378:LEU:HD22	1:A:378:LEU:H	1.46	0.79
1:A:382:ARG:NH2	8:A:601:HOH:O	2.23	0.72
2:T:10:DG:N7	8:T:103:HOH:O	2.25	0.69
1:A:375:ASP:OD1	1:A:377:ARG:HG3	1.97	0.65
1:A:378:LEU:CD2	1:A:378:LEU:H	2.10	0.64
1:A:376:LYS:HG3	1:A:377:ARG:N	2.15	0.61
1:A:323:LYS:NZ	8:A:604:HOH:O	2.33	0.61
1:A:127:GLU:O	1:A:130:GLN:HG2	2.00	0.61
1:A:160:THR:HG23	1:A:162:GLN:HG2	1.82	0.61
1:A:130:GLN:O	1:A:133:GLN:NE2	2.35	0.59
1:A:61[B]:ARG:NH1	2:T:3:DT:O4	2.37	0.58
1:A:376:LYS:CG	1:A:377:ARG:N	2.67	0.57
1:A:322[A]:SER:HB3	1:A:423:PHE:CD2	2.42	0.55
1:A:372:VAL:HG11	1:A:417:PRO:HG2	1.90	0.54
2:T:5:DA:N1	3:P:8[B]:DT:N3	2.53	0.53
1:A:169:GLN:OE1	1:A:173:GLN:HG3	2.10	0.52
5:A:503:GOL:H32	8:A:648:HOH:O	2.12	0.50
1:A:180:ILE:HA	1:A:188:LEU:HD22	1.94	0.49
1:A:84:ARG:NH2	1:A:350:GLU:OE1	2.46	0.48
1:A:372:VAL:CG1	1:A:417:PRO:HG2	2.44	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:T:9:DC:OP2	8:T:101:HOH:O	2.20	0.47
1:A:367:VAL:HG22	1:A:384[B]:CYS:SG	2.54	0.47
1:A:136:PRO:HA	1:A:180:ILE:HG21	1.98	0.46
1:A:401:THR:HA	1:A:404:LYS:CD	2.45	0.46
1:A:106:PHE:CG	1:A:200[B]:MET:HG2	2.50	0.46
1:A:362:VAL:HG22	1:A:432:SER:HB3	1.97	0.46
8:A:616:HOH:O	2:T:2:DA:H2''	2.15	0.45
1:A:356:ARG:NH2	8:A:619:HOH:O	2.47	0.45
1:A:369:SER:HB2	1:A:382:ARG:HD3	1.99	0.45
1:A:376:LYS:HG3	1:A:377:ARG:CG	2.47	0.44
1:A:369:SER:HB3	1:A:423:PHE:HB3	1.99	0.44
1:A:319:ILE:HG23	1:A:351[B]:ARG:CZ	2.49	0.43
1:A:375:ASP:OD1	1:A:376:LYS:N	2.52	0.42
1:A:405:ASN:N	1:A:405:ASN:OD1	2.44	0.42
1:A:389:ARG:NH1	8:A:632:HOH:O	2.52	0.42
1:A:375:ASP:OD1	1:A:376:LYS:HG2	2.19	0.42
2:T:3:DT:H2'	2:T:4:DT:H71	2.01	0.42
1:A:333:THR:OG1	1:A:336:GLN:HB2	2.19	0.41
1:A:132:LEU:O	1:A:135:GLN:HB2	2.20	0.41
1:A:48:ILE:HA	1:A:61[B]:ARG:HG3	2.03	0.41
1:A:367:VAL:HG11	1:A:382:ARG:HD2	2.03	0.41
1:A:178:LEU:HA	1:A:178:LEU:HD23	1.93	0.41
6:A:506[A]:DTP:O5'	6:A:506[A]:DTP:H8	2.21	0.40
1:A:311:LYS:HD3	1:A:313[A]:ARG:HH22	1.87	0.40
1:A:180:ILE:H	1:A:180:ILE:HG13	1.59	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:684:HOH:O	8:A:897:HOH:O[5_554]	2.12	0.08
8:A:722:HOH:O	8:A:769:HOH:O[5_664]	2.13	0.07
1:A:75:LEU:O	1:A:288[B]:SER:OG[6_555]	2.18	0.02

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/435 (102%)	436 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	384/372 (103%)	375 (98%)	9 (2%)	58	33

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	GLN
1	A	41	SER
1	A	115	ASP
1	A	160	THR
1	A	377	ARG
1	A	378	LEU
1	A	382	ARG
1	A	402	VAL
1	A	404	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GOL	A	503	-	5,5,5	0.49	0	5,5,5	0.41	0
5	GOL	A	504	-	5,5,5	0.28	0	5,5,5	0.30	0
5	GOL	A	505	-	5,5,5	0.25	0	5,5,5	0.22	0
6	DTP	A	506[A]	4	25,32,32	1.21	4 (16%)	26,50,50	2.45	1 (3%)
7	DPO	A	507[B]	4	8,8,8	1.02	1 (12%)	12,13,13	1.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	503	-	-	0/4/4/4	0/0/0/0
5	GOL	A	504	-	-	0/4/4/4	0/0/0/0
5	GOL	A	505	-	-	0/4/4/4	0/0/0/0
6	DTP	A	506[A]	4	-	0/18/34/34	0/3/3/3
7	DPO	A	507[B]	4	-	0/6/6/6	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	506[A]	DTP	C3'-C4'	-2.49	1.45	1.53
6	A	506[A]	DTP	C2'-C3'	-2.13	1.47	1.52
7	A	507[B]	DPO	P1-O4	-2.04	1.56	1.62
6	A	506[A]	DTP	C6-N6	2.30	1.43	1.34
6	A	506[A]	DTP	C2-N3	2.81	1.37	1.32

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	506[A]	DTP	N3-C2-N1	-11.84	119.57	128.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	503	GOL	1	0
6	A	506[A]	DTP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	430/435 (98%)	0.16	35 (8%) 15 18	7, 16, 49, 80	0
2	T	12/12 (100%)	0.29	2 (16%) 2 4	15, 21, 56, 65	0
3	P	9/9 (100%)	-0.30	0 100 100	12, 20, 29, 30	1 (11%)
All	All	451/456 (98%)	0.15	37 (8%) 14 18	7, 17, 50, 80	1 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	411	ILE	8.7
1	A	161	VAL	6.6
1	A	160	THR	5.7
1	A	376	LYS	5.1
1	A	180	ILE	5.1
1	A	410	GLY	5.0
1	A	181	ASP	5.0
1	A	1	MET	4.5
1	A	2	ALA	4.3
1	A	153	GLY	4.2
1	A	-2	GLY	4.2
1	A	412	GLN	4.0
1	A	183	LEU	4.0
1	A	378	LEU	3.8
2	T	3	DT	3.7
1	A	408	THR	3.7
1	A	182	ASN	3.6
1	A	133	GLN	3.6
1	A	132	LEU	3.5
1	A	-1	PRO	3.4
1	A	134	GLY	3.0
1	A	154	PRO	2.9
1	A	184	THR	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	179	GLN	2.7
1	A	328	LYS	2.7
1	A	130	GLN	2.7
1	A	135	GLN	2.6
1	A	413	THR	2.6
1	A	327	GLY	2.5
1	A	172	PHE	2.5
2	T	1	DC	2.5
1	A	380	SER	2.4
1	A	0	HIS	2.3
1	A	131	LYS	2.3
1	A	373	GLN	2.2
1	A	417	PRO	2.1
1	A	382	ARG	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	A	505	6/6	0.76	0.33	14.80	25,32,41,44	0
5	GOL	A	504	6/6	0.88	0.20	13.27	15,29,35,41	0
5	GOL	A	503	6/6	0.92	0.22	13.00	13,22,30,33	0
6	DTP	A	506[A]	30/30	0.98	0.05	-0.80	8,10,14,14	0
7	DPO	A	507[B]	9/9	0.99	0.05	-1.31	9,9,11,11	9
4	MN	A	501	1/1	1.00	0.03	-1.56	9,9,9,9	0
4	MN	A	502[B]	1/1	1.00	0.03	-2.25	9,9,9,9	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	MN	P	101[B]	1/1	0.93	0.23	-	14,14,14,14	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.