



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:02 AM GMT

PDB ID : 3KIO  
Title : mouse RNase H2 complex  
Authors : Shaban, N.; Harvey, S.; Perrino, F.W.; Hollis, T.  
Deposited on : 2009-11-02  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	 3% 41% 32% 11% • 15%
2	B	332	 4% 22% 17% • 58%
3	C	166	 2% 28% 25% 13% • 32%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3854 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonuclease H2 subunit A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	Se	0	0	0
			1986	1261	331	384	5	5			

- Molecule 2 is a protein called Ribonuclease H2 subunit B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	140	Total	C	N	O	S	Se	0	0	0
			983	636	170	173	1	3			

There are 25 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-23	MSE	-	EXPRESSION TAG	UNP Q80ZV0
B	-22	GLY	-	EXPRESSION TAG	UNP Q80ZV0
B	-21	SER	-	EXPRESSION TAG	UNP Q80ZV0
B	-20	SER	-	EXPRESSION TAG	UNP Q80ZV0
B	-19	HIS	-	EXPRESSION TAG	UNP Q80ZV0
B	-18	HIS	-	EXPRESSION TAG	UNP Q80ZV0
B	-17	HIS	-	EXPRESSION TAG	UNP Q80ZV0
B	-16	HIS	-	EXPRESSION TAG	UNP Q80ZV0
B	-15	HIS	-	EXPRESSION TAG	UNP Q80ZV0
B	-14	HIS	-	EXPRESSION TAG	UNP Q80ZV0
B	-13	SER	-	EXPRESSION TAG	UNP Q80ZV0
B	-12	GLN	-	EXPRESSION TAG	UNP Q80ZV0
B	-11	ASP	-	EXPRESSION TAG	UNP Q80ZV0
B	-10	PRO	-	EXPRESSION TAG	UNP Q80ZV0
B	-9	ASN	-	EXPRESSION TAG	UNP Q80ZV0
B	-8	SER	-	EXPRESSION TAG	UNP Q80ZV0
B	-7	LEU	-	EXPRESSION TAG	UNP Q80ZV0
B	-6	GLU	-	EXPRESSION TAG	UNP Q80ZV0
B	-5	VAL	-	EXPRESSION TAG	UNP Q80ZV0
B	-4	LEU	-	EXPRESSION TAG	UNP Q80ZV0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	PHE	-	EXPRESSION TAG	UNP Q80ZV0
B	-2	GLN	-	EXPRESSION TAG	UNP Q80ZV0
B	-1	GLY	-	EXPRESSION TAG	UNP Q80ZV0
B	0	PRO	-	EXPRESSION TAG	UNP Q80ZV0
B	239	PRO	THR	VARIANT	UNP Q80ZV0

- Molecule 3 is a protein called Ribonuclease H2 subunit C.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	113	Total	C	N	O	S	Se	0	0	0
			825	521	149	153	1	1			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	9	GLU	ASP	VARIANT	UNP Q9CQ18

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	42	Total	O	0	0
			42	42		
4	B	6	Total	O	0	0
			6	6		
4	C	12	Total	O	0	0
			12	12		



Chain C:

Amino Acid	Percentage
ASP	2%
ALA	
LYS	
VAL	
HIS	
ARG	
LEU	
GLY	
TRP	
PRO	
SER	
LEU	
ALA	
ALA	
ALA	
ILE	
HIS	
ALA	
GLN	
VAL	
PRO	
GLU	
ASP	
G78	
G81	
F82	
V83	
P84	
T86	
E87	
F88	
K89	
G90	
E91	
G92	
L93	
I94	
G95	
K96	
L97	
A103	
E104	
D105	
K106	
A107	
D108	
E109	
A110	
Q111	
E112	
P113	
L114	
E115	
L116	
ASP	
PHE	
ASP	
ARG	
LEU	
ILE	
GLY	
ALA	
THR	
GLY	
SER	
PHE	
SER	
HIS	
PHE	
THR	
L136	
V137	
G138	
L139	
E140	
T141	
V142	
P143	
G144	
F145	
P63	
P66	
G67	
L68	
P69	
G70	
V75	
P76	
F77	

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	279.29Å 40.42Å 67.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.62 – 2.90 24.62 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.3 (24.62-2.90) 99.4 (24.62-2.90)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.90 (at 2.89Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.4_113)	Depositor
R, $R_{free}$	0.234 , 0.305 0.247 , 0.320	Depositor DCC
$R_{free}$ test set	924 reflections (5.20%)	DCC
Wilson B-factor (Å <sup>2</sup> )	49.9	Xtriage
Anisotropy	0.011	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 70.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 17780 reflections	Xtriage
$F_o, F_c$ correlation	0.86	EDS
Total number of atoms	3854	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.61	2/2025 (0.1%)	0.76	8/2751 (0.3%)
2	B	0.43	0/997	0.58	0/1354
3	C	0.47	0/842	0.77	1/1140 (0.1%)
All	All	0.54	2/3864 (0.1%)	0.72	9/5245 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	112	TYR	CD2-CE2	-6.29	1.29	1.39
1	A	112	TYR	CE2-CZ	-5.45	1.31	1.38

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	64	LYS	N-CA-C	7.73	131.86	111.00
1	A	64	LYS	N-CA-CB	-7.54	97.02	110.60
3	C	58	GLY	N-CA-C	6.76	130.00	113.10
1	A	111	LYS	CB-CA-C	-6.35	97.69	110.40
1	A	88	ASP	CB-CA-C	5.83	122.07	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	92	ASP	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1986	0	1927	103	0
2	B	983	0	866	61	0
3	C	825	0	795	73	0
4	A	42	0	0	3	0
4	B	6	0	0	0	0
4	C	12	0	0	1	0
All	All	3854	0	3588	221	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 221 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:222:ARG:HH11	1:A:222:ARG:HG2	1.06	1.20
2:B:93:PRO:HD3	2:B:270:GLY:HA2	1.41	1.02
1:A:27:GLU:HG3	1:A:28:PRO:HD2	1.45	0.96
1:A:2:ASP:HA	1:A:5:GLU:CG	2.01	0.91
1:A:27:GLU:HG3	1:A:28:PRO:CD	2.08	0.84

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/301 (84%)	229 (91%)	20 (8%)	3 (1%)	16	48
2	B	132/332 (40%)	97 (74%)	21 (16%)	14 (11%)	0	1
3	C	109/166 (66%)	81 (74%)	16 (15%)	12 (11%)	0	1
All	All	493/799 (62%)	407 (83%)	57 (12%)	29 (6%)	2	6

5 of 29 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	11	ALA
2	B	46	GLY
2	B	92	ASP
2	B	156	SER
2	B	158	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/252 (85%)	171 (80%)	44 (20%)	1	4
2	B	82/286 (29%)	73 (89%)	9 (11%)	8	23
3	C	80/125 (64%)	56 (70%)	24 (30%)	0	1
All	All	377/663 (57%)	300 (80%)	77 (20%)	1	4

5 of 77 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	213	ASN
1	A	257	ASP
3	C	112	GLU
1	A	214	ASP
1	A	236	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	113	ASN
1	A	135	ASN
2	B	148	ASN
2	B	170	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	251/301 (83%)	-0.17	8 (3%) 51 43	19, 36, 114, 151	0
2	B	137/332 (41%)	0.49	13 (9%) 10 6	30, 106, 149, 159	0
3	C	112/166 (67%)	-0.11	4 (3%) 46 38	19, 47, 130, 155	0
All	All	500/799 (62%)	0.02	25 (5%) 32 26	19, 46, 136, 159	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	45	SER	6.1
1	A	74	ASN	4.5
1	A	62	ALA	4.4
1	A	10	ASN	3.8
3	C	137	TRP	3.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.