



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:10 PM GMT

PDB ID : 4KRP
Title : Nanobody/VHH domain 9G8 in complex with the extracellular region of EGFR
Authors : Ferguson, K.M.; Schmitz, K.R.
Deposited on : 2013-05-16
Resolution : 2.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2676 (2.84-2.80)
Clashscore	102246	3124 (2.84-2.80)
Ramachandran outliers	100387	3072 (2.84-2.80)
Sidechain outliers	100360	3074 (2.84-2.80)
RSRZ outliers	91569	2690 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	625	<div> <div>11%</div> <div> <div></div> <div>78%</div> <div>7%</div> <div>15%</div> </div> </div>
2	B	136	<div> <div>15%</div> <div> <div></div> <div>80%</div> <div>• •</div> <div>15%</div> </div> </div>
3	C	211	<div> <div></div> <div> <div></div> <div>90%</div> <div>9%</div> </div> </div>
4	D	220	<div> <div></div> <div> <div></div> <div>89%</div> <div>8%</div> <div>•</div> </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14877 atoms, of which 7017 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Epidermal growth factor receptor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	534	Total	C	H	N	O	S	0	0	0
			7001	2341	3252	651	712	45			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	614	THR	-	EXPRESSION TAG	UNP P00533
A	620	HIS	-	EXPRESSION TAG	UNP P00533
A	621	HIS	-	EXPRESSION TAG	UNP P00533
A	622	HIS	-	EXPRESSION TAG	UNP P00533
A	623	HIS	-	EXPRESSION TAG	UNP P00533
A	624	HIS	-	EXPRESSION TAG	UNP P00533
A	625	HIS	-	EXPRESSION TAG	UNP P00533

- Molecule 2 is a protein called Nanobody/VHH domain 9G8.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	116	Total	C	H	N	O	S	0	0	0
			1473	516	655	135	163	4			

- Molecule 3 is a protein called Cetuximab light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	211	Total	C	H	N	O	S	0	0	0
			3099	992	1506	270	327	4			

- Molecule 4 is a protein called Cetuximab heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	D	213	Total	C	H	N	O	S	0	0	0
			3079	1007	1494	259	314	5			

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	2	Total	C	H	N	O	0	0
			55	16	27	2	10		
5	A	2	Total	C	H	N	O	0	0
			55	16	27	2	10		

- Molecule 6 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
6	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
6	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
6	D	1	Total	C	H	N	O	0	0
			28	8	14	1	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	O	0	0
			1	1		
7	C	1	Total	O	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	1	Total	O	0	0
			1	1		



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.37Å 95.84Å 129.48Å 90.00° 99.95° 90.00°	Depositor
Resolution (Å)	44.46 – 2.82 44.46 – 2.82	Depositor EDS
% Data completeness (in resolution range)	98.0 (44.46-2.82) 98.1 (44.46-2.82)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.27 (at 2.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.215 , 0.263 0.218 , 0.266	Depositor DCC
R_{free} test set	1890 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	51.9	Xtriage
Anisotropy	0.182	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 51.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 37834 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	14877	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/3825	0.43	0/5230
2	B	0.26	0/838	0.39	0/1146
3	C	0.36	0/1627	0.52	0/2218
4	D	0.36	0/1627	0.52	0/2233
All	All	0.30	0/7917	0.47	0/10827

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3749	3252	3319	23	0
2	B	818	655	655	4	0
3	C	1593	1506	1503	13	0
4	D	1585	1494	1492	9	0
5	A	56	54	50	0	0
6	A	42	42	39	1	0
6	D	14	14	13	0	0
7	A	1	0	0	0	0
7	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	D	1	0	0	0	0
All	All	7860	7017	7071	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:337:ASN:HD22	6:A:703:NAG:H83	1.60	0.67
1:A:376:GLU:OE1	1:A:403:ARG:NH1	2.28	0.66
1:A:455:LYS:O	2:B:27:ARG:NH2	2.31	0.64
1:A:403:ARG:NH2	2:B:100:TYR:O	2.31	0.64
2:B:54:SER:O	2:B:104:SER:OG	2.18	0.61
3:C:80:SER:HA	3:C:106:LEU:HD22	1.83	0.61
4:D:32:TYR:CZ	4:D:101:TYR:CE1	2.90	0.59
1:A:14:LEU:HG	1:A:89:TYR:OH	2.02	0.58
1:A:245:LEU:HD13	1:A:577:GLY:O	2.05	0.56
3:C:164:THR:HG22	3:C:174:SER:H	1.72	0.55
1:A:28:GLN:HG3	1:A:29:ARG:N	2.23	0.53
1:A:276:VAL:N	1:A:284:VAL:O	2.40	0.53
2:B:101:GLN:OE1	2:B:107:TYR:OH	2.26	0.51
3:C:13:VAL:HG13	3:C:17:GLU:HB2	1.94	0.50
3:C:158:ASN:OD1	3:C:158:ASN:N	2.46	0.49
4:D:6:GLN:O	4:D:111:GLN:NE2	2.46	0.49
1:A:588:ASP:O	1:A:591:HIS:N	2.48	0.47
1:A:407:LYS:NZ	1:A:434:ASP:OD2	2.39	0.47
1:A:447:TYR:O	1:A:449:ASN:N	2.43	0.47
3:C:140:TYR:CG	3:C:141:PRO:HA	2.50	0.46
1:A:365:PRO:HB3	1:A:387:PRO:HG3	1.97	0.46
1:A:41:LEU:HB3	1:A:65:VAL:HG22	1.99	0.45
1:A:272:PRO:HG2	1:A:275:TYR:CG	2.51	0.45
4:D:47:TRP:HZ2	4:D:50:VAL:HG23	1.82	0.44
3:C:44:PRO:HG2	4:D:109:TRP:CE2	2.53	0.44
4:D:72:ASP:OD1	4:D:74:SER:OG	2.28	0.44
1:A:584:TRP:NE1	1:A:601:THR:O	2.50	0.44
3:C:19:VAL:HG23	3:C:78:VAL:CG2	2.48	0.44
1:A:443:LYS:HE3	3:C:94:TRP:CZ2	2.53	0.43
1:A:268:VAL:HG12	1:A:270:LYS:H	1.83	0.43
1:A:208:CYS:N	1:A:216:CYS:SG	2.91	0.43
4:D:207:LYS:N	4:D:208:PRO:HD2	2.34	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:124:GLN:O	3:C:127:SER:OG	2.32	0.42
3:C:37:GLN:O	3:C:45:ARG:N	2.51	0.42
4:D:4:LEU:HD12	4:D:22:CYS:SG	2.59	0.41
1:A:326:SER:OG	1:A:327:ILE:N	2.52	0.41
4:D:105:GLU:HG2	4:D:105:GLU:O	2.21	0.41
3:C:108:ARG:HG3	3:C:109:THR:O	2.21	0.41
1:A:246:TYR:OH	1:A:563:ASP:OD2	2.27	0.41
3:C:13:VAL:HG11	3:C:78:VAL:HG21	2.02	0.41
4:D:105:GLU:N	4:D:105:GLU:OE1	2.43	0.41
1:A:427:ARG:HA	1:A:492:TRP:CD1	2.56	0.41
1:A:272:PRO:HG2	1:A:275:TYR:CD2	2.56	0.40
3:C:19:VAL:CG2	3:C:78:VAL:HG21	2.51	0.40
1:A:97:VAL:HB	1:A:126:PHE:CE2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	530/625 (85%)	478 (90%)	51 (10%)	1 (0%)	52	83
2	B	112/136 (82%)	96 (86%)	16 (14%)	0	100	100
3	C	209/211 (99%)	204 (98%)	4 (2%)	1 (0%)	34	68
4	D	209/220 (95%)	198 (95%)	11 (5%)	0	100	100
All	All	1060/1192 (89%)	976 (92%)	82 (8%)	2 (0%)	52	83

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	214	ALA
3	C	138	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/546 (67%)	363 (99%)	5 (1%)	74	93
2	B	66/110 (60%)	64 (97%)	2 (3%)	48	81
3	C	179/188 (95%)	177 (99%)	2 (1%)	80	95
4	D	176/190 (93%)	173 (98%)	3 (2%)	68	91
All	All	789/1034 (76%)	777 (98%)	12 (2%)	72	93

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	266	THR
1	A	278	THR
1	A	295	GLU
1	A	505	VAL
1	A	604	CYS
2	B	27	ARG
2	B	53	TRP
3	C	106	LEU
3	C	181	LEU
4	D	25	SER
4	D	51	ILE
4	D	134	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	480	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	NAG	A	701	1,5	14,14,15	0.69	1 (7%)	15,19,21	0.56	0
5	NAG	A	702	5	14,14,15	0.54	0	15,19,21	1.05	0
5	NAG	A	705	1,5	14,14,15	0.53	0	15,19,21	1.06	0
5	NAG	A	706	5	14,14,15	0.55	0	15,19,21	1.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	701	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	702	5	-	0/6/23/26	0/1/1/1
5	NAG	A	705	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	706	5	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	701	NAG	O5-C1	-2.27	1.39	1.43

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	NAG	A	703	1	14,14,15	0.29	0	15,19,21	0.51	0
6	NAG	A	704	1	14,14,15	0.53	0	15,19,21	1.03	0
6	NAG	A	707	1	14,14,15	0.27	0	15,19,21	0.61	0
6	NAG	D	301	4	14,14,15	0.52	0	15,19,21	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	703	1	-	0/6/23/26	0/1/1/1
6	NAG	A	704	1	-	0/6/23/26	0/1/1/1
6	NAG	A	707	1	-	0/6/23/26	0/1/1/1
6	NAG	D	301	4	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	703	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	534/625 (85%)	0.56	69 (12%) 5 2	26, 85, 150, 163	0
2	B	116/136 (85%)	0.79	20 (17%) 2 1	59, 94, 124, 137	0
3	C	211/211 (100%)	-0.27	0 100 100	18, 35, 54, 70	0
4	D	213/220 (96%)	-0.26	0 100 100	18, 34, 53, 84	0
All	All	1074/1192 (90%)	0.26	89 (8%) 14 7	18, 52, 142, 163	0

All (89) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	103	ALA	9.2
1	A	126	PHE	9.0
1	A	116	LEU	7.4
1	A	82	ILE	6.8
1	A	107	GLY	6.4
1	A	219	PRO	6.2
1	A	101	TYR	6.1
1	A	124	VAL	6.1
1	A	80	LEU	5.9
1	A	54	PHE	5.6
1	A	129	ASN	5.3
2	B	89	GLU	5.3
1	A	130	PRO	5.2
1	A	234	ALA	5.1
1	A	104	ASN	5.1
1	A	114	ARG	4.9
1	A	119	ILE	4.8
2	B	88	PRO	4.8
1	A	91	ASN	4.7
1	A	115	ASN	4.6
1	A	24	PHE	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	94	TYR	4.4
2	B	91	THR	4.3
1	A	81	GLN	4.2
1	A	97	VAL	4.1
1	A	117	GLN	4.0
1	A	222	SER	4.0
1	A	95	LEU	4.0
2	B	55	SER	3.8
1	A	36	VAL	3.8
2	B	42	GLY	3.8
1	A	217	THR	3.7
1	A	66	LEU	3.7
1	A	49	ASN	3.6
1	A	612	CYS	3.6
1	A	602	TYR	3.6
1	A	92	SER	3.5
1	A	96	ALA	3.5
1	A	43	ILE	3.5
1	A	94	ALA	3.5
1	A	83	ILE	3.5
1	A	127	SER	3.4
1	A	214	ALA	3.3
1	A	58	ILE	3.2
1	A	38	LEU	3.2
1	A	611	GLY	3.2
2	B	75	ALA	3.1
1	A	123	ALA	3.1
1	A	213	ALA	3.0
1	A	106	THR	3.0
1	A	125	ARG	3.0
1	A	67	ILE	2.9
2	B	22	CYS	2.9
2	B	71	SER	2.9
1	A	108	LEU	2.9
1	A	607	PRO	2.9
2	B	80	TYR	2.9
1	A	55	LEU	2.8
1	A	37	VAL	2.8
1	A	225	LEU	2.8
1	A	89	TYR	2.7
1	A	100	ASN	2.7
1	A	113	MET	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	102	ASP	2.7
1	A	587	ALA	2.7
1	A	215	GLY	2.6
1	A	71	THR	2.6
1	A	35	GLU	2.5
1	A	286	ALA	2.5
1	A	274	ASN	2.5
1	A	586	TYR	2.5
2	B	90	ASP	2.5
2	B	86	LEU	2.4
2	B	9	GLY	2.4
2	B	8	GLY	2.4
1	A	211	GLN	2.3
2	B	123	VAL	2.3
1	A	34	CYS	2.3
1	A	41	LEU	2.2
2	B	74	ASN	2.2
2	B	36	TRP	2.2
2	B	73	ASP	2.1
1	A	574	GLY	2.1
1	A	610	GLU	2.1
1	A	120	LEU	2.1
2	B	121	THR	2.1
1	A	79	ASN	2.1
1	A	88	TYR	2.1
2	B	92	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	NAG	A	702	14/15	0.74	0.24	0.86	66,92,111,114	0
5	NAG	A	701	14/15	0.93	0.14	-0.45	53,79,96,104	0
5	NAG	A	705	14/15	0.92	0.13	-2.66	43,58,70,77	0
5	NAG	A	706	14/15	0.81	0.21	-	69,89,105,118	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	NAG	D	301	14/15	0.78	0.15	-	64,83,97,112	0
6	NAG	A	707	14/15	0.84	0.19	-	60,85,102,106	0
6	NAG	A	704	14/15	0.53	0.22	-	71,97,116,121	0
6	NAG	A	703	14/15	0.89	0.17	-	59,95,116,120	0

6.5 Other polymers [i](#)

There are no such residues in this entry.