



Full wwPDB NMR Structure Validation Report ⓘ

Apr 27, 2016 – 12:32 AM BST

PDB ID : 2KY2
Title : The Structure of RNA Internal Loops with Tandem AG Pairs:
5'UAGG/3'GGAU
Authors : Hammond, N.B.; Kennedy, S.D.; Turner, D.H.
Deposited on : 2010-05-13

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR



The overall completeness of chemical shifts assignment is 57%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
RNA backbone	3027	600

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	9	 <div>33%67%</div>
1	B	9	 <div>56%44%</div>

2 Ensemble composition and analysis ⓘ

This entry contains 10 models. This entry does not contain protein, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 586 atoms, of which 200 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'.

Mol	Chain	Residues	Atoms						Trace
1	A	9	Total	C	H	N	O	P	0
			293	87	100	38	60	8	
1	B	9	Total	C	H	N	O	P	0
			293	87	100	38	60	8	

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'



- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'



- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'



4.2.2 Score per residue for model 2

- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain A:  44% 56%



- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'


Chain B:  44% 56%



4.2.3 Score per residue for model 3

- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain A:  11% 89%



- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'


Chain B:  44% 56%



4.2.4 Score per residue for model 4


- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain A:  56% 44%




- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain B:  44% 56%




4.2.5 Score per residue for model 5

- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain A:  33% 67%



- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain B:  33% 67%



4.2.6 Score per residue for model 6

- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain A:  44% 56%



- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'


Chain B:  44% 56%



4.2.7 Score per residue for model 7

- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain A:  22% 78%



- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain B:  44% 56%



4.2.8 Score per residue for model 8

- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain A:  44% 56%



- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'


Chain B:  33% 67%



4.2.9 Score per residue for model 9

- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain A:  33% 67%




- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain B:  56% 44%



4.2.10 Score per residue for model 10

- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain A:  33% 67%



- Molecule 1: 5'-R(*GP*GP*UP*AP*GP*GP*CP*CP*A)-3'

Chain B:  33% 67%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 40 calculated structures, 10 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	1.2
AMBER	refinement	9

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 16953
Number of chemical shift lists	1
Total number of shifts	216
Number of shifts mapped to atoms	216
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	57%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.41±0.01	0±0/216 (0.0±0.0%)	2.19±0.02	12±1/336 (3.6±0.3%)
1	B	1.40±0.01	0±0/216 (0.0±0.0%)	2.18±0.02	12±1/336 (3.6±0.3%)
All	All	1.41	0/4320 (0.0%)	2.19	240/6720 (3.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	B	0.0±0.0	0.1±0.3
All	All	0	1

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	9	A	N1-C6-N6	-9.44	112.94	118.60	5	10
1	B	18	A	N1-C6-N6	-9.16	113.10	118.60	5	10
1	A	4	A	N1-C6-N6	-8.20	113.68	118.60	8	10
1	B	13	A	N1-C6-N6	-8.05	113.77	118.60	9	10
1	A	9	A	C5-C6-N1	7.81	121.61	117.70	9	10
1	B	18	A	C5-C6-N1	7.63	121.52	117.70	5	10
1	B	13	A	C5-C6-N1	7.60	121.50	117.70	5	10
1	B	17	C	N3-C2-O2	-7.59	116.59	121.90	5	10
1	A	4	A	C5-C6-N1	7.56	121.48	117.70	6	10
1	A	7	C	N3-C2-O2	-7.50	116.65	121.90	3	10
1	B	16	C	N3-C2-O2	-7.45	116.69	121.90	4	10
1	A	8	C	N3-C2-O2	-7.32	116.78	121.90	1	10
1	B	18	A	C4-C5-C6	-6.84	113.58	117.00	7	10
1	A	9	A	C4-C5-C6	-6.75	113.62	117.00	5	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	13	A	C4-C5-C6	-6.31	113.84	117.00	5	10
1	A	4	A	C4-C5-C6	-6.22	113.89	117.00	2	10
1	B	16	C	N1-C2-O2	6.11	122.57	118.90	4	9
1	A	8	C	N1-C2-O2	6.05	122.53	118.90	1	9
1	B	17	C	N1-C2-O2	6.00	122.50	118.90	5	10
1	A	6	G	N1-C6-O6	-5.86	116.38	119.90	5	6
1	B	15	G	N1-C6-O6	-5.83	116.40	119.90	1	4
1	A	7	C	N1-C2-O2	5.81	122.38	118.90	3	8
1	A	8	C	O4'-C1'-N1	5.74	112.79	108.20	10	6
1	B	17	C	O4'-C1'-N1	5.55	112.64	108.20	5	6
1	B	14	G	N1-C6-O6	-5.42	116.65	119.90	7	3
1	A	2	G	N1-C6-O6	-5.41	116.66	119.90	9	3
1	A	3	U	O4'-C1'-N1	5.36	112.48	108.20	1	6
1	B	11	G	N1-C6-O6	-5.34	116.70	119.90	1	3
1	B	12	U	O4'-C1'-N1	5.20	112.36	108.20	1	3
1	A	5	G	N1-C6-O6	-5.18	116.79	119.90	3	2
1	B	10	G	N1-C6-O6	-5.18	116.79	119.90	5	1
1	B	16	C	O4'-C1'-N1	5.18	112.34	108.20	8	1

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	B	11	G	Sidechain	1

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	3860	2000	2000	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	8/9 (89%)	0±0 (0±0%)	0±0 (0±0%)	0.72±0.03
1	B	8/9 (89%)	0±0 (0±0%)	0±0 (0±0%)	0.73±0.03
All	All	160/180 (89%)	0 (0%)	0 (0%)	0.72

The overall RNA backbone suiteness is 0.72.

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 57% for the well-defined parts and 57% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 16953

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	216
Number of shifts mapped to atoms	216
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 57%, i.e. 196 atoms were assigned a chemical shift out of a possible 346. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Sidechain	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	196/346 (57%)	160/202 (79%)	36/118 (31%)	0/26 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 57%, i.e. 196 atoms were assigned a chemical shift out of a possible 346. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Sidechain	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	196/346 (57%)	160/202 (79%)	36/118 (31%)	0/26 (0%)

7.1.4 Statistically unusual chemical shifts ⓘ

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots ⓘ

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned_chem_shift_list_1). RCI is only applicable to proteins.