



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:15 PM GMT

PDB ID : 4KY4
Title : Crystal structure of non-classical TS inhibitor 2 in complex with Toxoplasma gondii TS-DHFR
Authors : Sharma, H.; Anderson, K.S.
Deposited on : 2013-05-28
Resolution : 2.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	610	<div> <div>2%</div> <div>63% 16% • 20%</div> </div>
1	B	610	<div> <div>3%</div> <div>61% 17% • 20%</div> </div>
1	C	610	<div> <div>3%</div> <div>64% 15% • 20%</div> </div>
1	D	610	<div> <div>3%</div> <div>62% 15% • 20%</div> </div>
1	E	610	<div> <div>3%</div> <div>64% 15% • 20%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	610	
1	G	610	
1	H	610	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	UMP	D	701	-	-	-	X
2	UMP	F	701	-	-	X	-
3	04J	B	702	-	-	-	X
3	04J	D	702	-	-	-	X
3	04J	H	702	-	-	X	-
5	1UE	A	704	-	-	X	-
5	1UE	B	704	-	-	X	X
5	1UE	C	704	-	-	X	-
5	1UE	D	704	-	-	X	-
5	1UE	E	704	-	-	-	X
5	1UE	F	704	-	-	X	X
5	1UE	G	704	-	-	X	X
5	1UE	H	704	-	-	-	X

2 Entry composition


There are 5 unique types of molecules in this entry. The entry contains 32492 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional dihydrofolate reductase-thymidylate synthase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	489	Total	C	N	O	S	0	1	0
			3931	2522	680	704	25			
1	B	491	Total	C	N	O	S	0	1	0
			3948	2531	682	710	25			
1	C	489	Total	C	N	O	S	0	1	0
			3931	2522	680	704	25			
1	D	491	Total	C	N	O	S	0	1	0
			3948	2531	682	710	25			
1	E	489	Total	C	N	O	S	0	1	0
			3931	2522	680	704	25			
1	F	491	Total	C	N	O	S	0	1	0
			3948	2531	682	710	25			
1	G	489	Total	C	N	O	S	0	1	0
			3931	2522	680	704	25			
1	H	491	Total	C	N	O	S	0	1	0
			3948	2531	682	710	25			

- Molecule 2 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula: C₉H₁₃N₂O₈P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	B	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	C	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	D	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	E	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	F	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	G	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	H	1	Total	C	N	O	P	0	0
			20	9	2	8	1		

- Molecule 3 is AMINOPTERIN (three-letter code: 04J) (formula: C₁₉H₂₀N₈O₅).




- Molecule 4 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
4	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
4	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
4	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
4	E	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
4	F	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
4	G	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
4	H	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 5 is 2-AMINO-5-(PHENYLSULFANYL)-3,9-DIHYDRO-4H-PYRIMIDO[4,5-B]IN DOL-4-ONE (three-letter code: 1UE) (formula: C₁₆H₁₂N₄OS).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			22	16	4	1	1		
5	B	1	Total	C	N	O	S	0	0
			22	16	4	1	1		
5	C	1	Total	C	N	O	S	0	0
			22	16	4	1	1		
5	D	1	Total	C	N	O	S	0	0
			22	16	4	1	1		
5	E	1	Total	C	N	O	S	0	0
			22	16	4	1	1		
5	F	1	Total	C	N	O	S	0	0
			22	16	4	1	1		
5	G	1	Total	C	N	O	S	0	0
			22	16	4	1	1		
5	H	1	Total	C	N	O	S	0	0
			22	16	4	1	1		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional dihydrofolate reductase-thymidylate synthase




- Molecule 1: Bifunctional dihydrofolate reductase-thymidylate synthase






- Molecule 1: Bifunctional dihydrofolate reductase-thymidylate synthase




- Molecule 1: Bifunctional dihydrofolate reductase-thymidylate synthase








• Molecule 1: Bifunctional dihydrofolate reductase-thymidylate synthase



• Molecule 1: Bifunctional dihydrofolate reductase-thymidylate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	53.54Å 145.12Å 176.16Å 89.97° 89.95° 89.89°	Depositor
Resolution (Å)	48.28 – 2.79 48.37 – 2.79	Depositor EDS
% Data completeness (in resolution range)	97.8 (48.28-2.79) 97.5 (48.37-2.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.220 , 0.268 0.221 , 0.264	Depositor DCC
R_{free} test set	6476 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	73.2	Xtriage
Anisotropy	0.414	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 37.2	EDS
Estimated twinning fraction	0.427 for h,-k,-l 0.427 for -h,k,-l 0.439 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 128608 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	32492	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, 1UE, UMP, 04J

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.59	0/4026	0.76	3/5450 (0.1%)
1	B	0.67	2/4043 (0.0%)	0.85	4/5473 (0.1%)
1	C	0.58	0/4026	0.76	1/5450 (0.0%)
1	D	0.66	2/4043 (0.0%)	0.84	3/5473 (0.1%)
1	E	0.58	0/4026	0.75	2/5450 (0.0%)
1	F	0.65	2/4043 (0.0%)	0.84	4/5473 (0.1%)
1	G	0.65	3/4026 (0.1%)	0.82	2/5450 (0.0%)
1	H	0.58	0/4043	0.75	1/5473 (0.0%)
All	All	0.62	9/32276 (0.0%)	0.80	20/43692 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	153	GLY	N-CA	6.70	1.56	1.46
1	D	153	GLY	N-CA	6.26	1.55	1.46
1	G	25	TRP	CG-CD1	5.66	1.44	1.36
1	B	25	TRP	CG-CD1	5.46	1.44	1.36
1	G	153	GLY	N-CA	5.45	1.54	1.46

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	9	VAL	CB-CA-C	-7.17	97.77	111.40
1	D	9	VAL	CB-CA-C	-6.95	98.20	111.40
1	F	358	ARG	NE-CZ-NH1	6.72	123.66	120.30
1	F	9	VAL	CB-CA-C	-6.58	98.89	111.40
1	H	470	ARG	NE-CZ-NH1	6.43	123.52	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	155	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3931	0	3900	78	0
1	B	3948	0	3910	79	0
1	C	3931	0	3900	78	0
1	D	3948	0	3910	75	0
1	E	3931	0	3900	70	0
1	F	3948	0	3910	80	0
1	G	3931	0	3900	71	0
1	H	3948	0	3910	70	0
2	A	20	0	11	3	0
2	B	20	0	11	2	0
2	C	20	0	11	5	0
2	D	20	0	11	2	0
2	E	20	0	11	2	0
2	F	20	0	11	7	0
2	G	20	0	11	2	0
2	H	20	0	11	5	0
3	A	32	0	18	5	0
3	B	32	0	18	1	0
3	C	32	0	18	2	0
3	D	32	0	18	3	0
3	E	32	0	18	2	0
3	F	32	0	18	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	32	0	18	2	0
3	H	32	0	18	14	0
4	A	48	0	26	4	0
4	B	48	0	26	2	0
4	C	48	0	26	2	0
4	D	48	0	26	4	0
4	E	48	0	26	3	0
4	F	48	0	26	4	0
4	G	48	0	26	3	0
4	H	48	0	26	15	0
5	A	22	0	12	7	0
5	B	22	0	12	9	0
5	C	22	0	12	7	0
5	D	22	0	12	7	0
5	E	22	0	12	4	0
5	F	22	0	12	7	0
5	G	22	0	12	7	0
5	H	22	0	12	4	0
All	All	32492	0	31776	604	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 604 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:516:LEU:HG	5:A:704:1UE:C6'	1.62	1.28
1:C:516:LEU:HG	5:C:704:1UE:C6'	1.70	1.21
4:D:703:NDP:H8A	4:D:703:NDP:H52A	1.25	1.14
1:C:516:LEU:HG	5:C:704:1UE:H11	1.18	1.13
1:A:516:LEU:HG	5:A:704:1UE:H11	1.18	1.11

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	476/610 (78%)	448 (94%)	23 (5%)	5 (1%)	17	50
1	B	478/610 (78%)	454 (95%)	21 (4%)	3 (1%)	30	65
1	C	476/610 (78%)	444 (93%)	28 (6%)	4 (1%)	24	58
1	D	478/610 (78%)	451 (94%)	22 (5%)	5 (1%)	19	52
1	E	476/610 (78%)	448 (94%)	24 (5%)	4 (1%)	24	58
1	F	478/610 (78%)	456 (95%)	18 (4%)	4 (1%)	24	58
1	G	476/610 (78%)	450 (94%)	22 (5%)	4 (1%)	24	58
1	H	478/610 (78%)	451 (94%)	23 (5%)	4 (1%)	24	58
All	All	3816/4880 (78%)	3602 (94%)	181 (5%)	33 (1%)	21	55

5 of 33 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	139	GLU
1	A	286	SER
1	A	429	TYR
1	B	190	ASP
1	B	429	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	429/525 (82%)	405 (94%)	24 (6%)	26	59
1	B	431/525 (82%)	389 (90%)	42 (10%)	10	29
1	C	429/525 (82%)	401 (94%)	28 (6%)	21	52
1	D	431/525 (82%)	387 (90%)	44 (10%)	9	26
1	E	429/525 (82%)	404 (94%)	25 (6%)	25	57
1	F	431/525 (82%)	385 (89%)	46 (11%)	8	24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	429/525 (82%)	384 (90%)	45 (10%)	8	24
1	H	431/525 (82%)	407 (94%)	24 (6%)	26	59
All	All	3440/4200 (82%)	3162 (92%)	278 (8%)	15	39

5 of 278 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	395	LEU
1	E	376	LYS
1	H	37	ARG
1	D	401	LYS
1	E	41	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	168	HIS
1	F	168	HIS
1	H	450	GLN
1	E	450	GLN
1	F	416	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

32 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UMP	A	701	-	16,21,21	1.52	4 (25%)	23,31,31	2.01	5 (21%)
3	04J	A	702	-	26,34,34	2.00	4 (15%)	29,47,47	1.73	6 (20%)
4	NDP	A	703	-	42,52,52	1.76	7 (16%)	55,80,80	2.19	5 (9%)
5	1UE	A	704	-	24,25,25	2.48	9 (37%)	25,36,36	1.77	4 (16%)
2	UMP	B	701	-	16,21,21	1.61	5 (31%)	23,31,31	1.86	6 (26%)
3	04J	B	702	-	26,34,34	1.95	3 (11%)	29,47,47	1.42	4 (13%)
4	NDP	B	703	-	42,52,52	1.72	7 (16%)	55,80,80	2.37	9 (16%)
5	1UE	B	704	-	24,25,25	2.46	9 (37%)	25,36,36	1.86	4 (16%)
2	UMP	C	701	-	16,21,21	1.52	4 (25%)	23,31,31	2.03	5 (21%)
3	04J	C	702	-	26,34,34	1.96	3 (11%)	29,47,47	1.68	6 (20%)
4	NDP	C	703	-	42,52,52	1.73	7 (16%)	55,80,80	2.28	7 (12%)
5	1UE	C	704	-	24,25,25	2.38	9 (37%)	25,36,36	1.76	5 (20%)
2	UMP	D	701	-	16,21,21	1.70	6 (37%)	23,31,31	1.88	6 (26%)
3	04J	D	702	-	26,34,34	1.99	4 (15%)	29,47,47	1.42	4 (13%)
4	NDP	D	703	-	42,52,52	1.71	11 (26%)	55,80,80	2.29	7 (12%)
5	1UE	D	704	-	24,25,25	2.36	9 (37%)	25,36,36	1.80	5 (20%)
2	UMP	E	701	-	16,21,21	1.70	5 (31%)	23,31,31	1.84	5 (21%)
3	04J	E	702	-	26,34,34	1.91	3 (11%)	29,47,47	1.52	5 (17%)
4	NDP	E	703	-	42,52,52	1.76	7 (16%)	55,80,80	2.22	3 (5%)
5	1UE	E	704	-	24,25,25	2.29	9 (37%)	25,36,36	1.78	4 (16%)
2	UMP	F	701	-	16,21,21	1.61	5 (31%)	23,31,31	1.61	2 (8%)
3	04J	F	702	-	26,34,34	1.99	4 (15%)	29,47,47	1.45	5 (17%)
4	NDP	F	703	-	42,52,52	1.75	13 (30%)	55,80,80	2.30	9 (16%)
5	1UE	F	704	-	24,25,25	2.44	10 (41%)	25,36,36	1.91	4 (16%)
2	UMP	G	701	-	16,21,21	1.67	5 (31%)	23,31,31	1.85	5 (21%)
3	04J	G	702	-	26,34,34	1.96	4 (15%)	29,47,47	1.51	5 (17%)
4	NDP	G	703	-	42,52,52	1.73	11 (26%)	55,80,80	2.27	7 (12%)
5	1UE	G	704	-	24,25,25	2.55	9 (37%)	25,36,36	1.88	4 (16%)
2	UMP	H	701	-	16,21,21	1.53	4 (25%)	23,31,31	2.14	5 (21%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	04J	H	702	-	26,34,34	1.90	4 (15%)	29,47,47	1.60	5 (17%)
4	NDP	H	703	-	42,52,52	1.69	7 (16%)	55,80,80	2.33	8 (14%)
5	1UE	H	704	-	24,25,25	2.31	10 (41%)	25,36,36	1.83	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UMP	A	701	-	-	0/6/22/22	0/2/2/2
3	04J	A	702	-	-	0/16/22/22	0/3/3/3
4	NDP	A	703	-	-	0/30/77/77	0/5/5/5
5	1UE	A	704	-	-	0/4/4/4	0/4/4/4
2	UMP	B	701	-	-	0/6/22/22	0/2/2/2
3	04J	B	702	-	-	0/16/22/22	0/3/3/3
4	NDP	B	703	-	-	0/30/77/77	0/5/5/5
5	1UE	B	704	-	-	0/4/4/4	0/4/4/4
2	UMP	C	701	-	-	0/6/22/22	0/2/2/2
3	04J	C	702	-	-	0/16/22/22	0/3/3/3
4	NDP	C	703	-	-	0/30/77/77	0/5/5/5
5	1UE	C	704	-	-	0/4/4/4	0/4/4/4
2	UMP	D	701	-	-	0/6/22/22	0/2/2/2
3	04J	D	702	-	-	0/16/22/22	0/3/3/3
4	NDP	D	703	-	-	0/30/77/77	0/5/5/5
5	1UE	D	704	-	-	0/4/4/4	0/4/4/4
2	UMP	E	701	-	-	0/6/22/22	0/2/2/2
3	04J	E	702	-	-	0/16/22/22	0/3/3/3
4	NDP	E	703	-	-	0/30/77/77	0/5/5/5
5	1UE	E	704	-	-	0/4/4/4	0/4/4/4
2	UMP	F	701	-	-	0/6/22/22	0/2/2/2
3	04J	F	702	-	-	0/16/22/22	0/3/3/3
4	NDP	F	703	-	-	0/30/77/77	0/5/5/5
5	1UE	F	704	-	-	0/4/4/4	0/4/4/4
2	UMP	G	701	-	-	0/6/22/22	0/2/2/2
3	04J	G	702	-	-	0/16/22/22	0/3/3/3
4	NDP	G	703	-	-	0/30/77/77	0/5/5/5
5	1UE	G	704	-	-	0/4/4/4	0/4/4/4
2	UMP	H	701	-	-	0/6/22/22	0/2/2/2
3	04J	H	702	-	-	0/16/22/22	0/3/3/3
4	NDP	H	703	-	-	0/30/77/77	0/5/5/5
5	1UE	H	704	-	-	0/4/4/4	0/4/4/4

The worst 5 of 211 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	704	1UE	C2'-S1	-6.37	1.65	1.77
5	A	704	1UE	C2'-S1	-6.23	1.65	1.77
5	B	704	1UE	C2'-S1	-6.02	1.66	1.77
3	A	702	04J	C11-C	-5.71	1.38	1.50
3	A	702	04J	C9-C6	-5.44	1.40	1.51

The worst 5 of 168 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	703	NDP	N3A-C2A-N1A	-14.36	117.90	128.89
4	B	703	NDP	N3A-C2A-N1A	-14.30	117.95	128.89
4	C	703	NDP	N3A-C2A-N1A	-14.12	118.08	128.89
4	F	703	NDP	N3A-C2A-N1A	-14.02	118.16	128.89
4	D	703	NDP	N3A-C2A-N1A	-13.99	118.18	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

32 monomers are involved in 142 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	UMP	3	0
3	A	702	04J	5	0
4	A	703	NDP	4	0
5	A	704	1UE	7	0
2	B	701	UMP	2	0
3	B	702	04J	1	0
4	B	703	NDP	2	0
5	B	704	1UE	9	0
2	C	701	UMP	5	0
3	C	702	04J	2	0
4	C	703	NDP	2	0
5	C	704	1UE	7	0
2	D	701	UMP	2	0
3	D	702	04J	3	0
4	D	703	NDP	4	0
5	D	704	1UE	7	0
2	E	701	UMP	2	0
3	E	702	04J	2	0
4	E	703	NDP	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	704	1UE	4	0
2	F	701	UMP	7	0
3	F	702	04J	2	0
4	F	703	NDP	4	0
5	F	704	1UE	7	0
2	G	701	UMP	2	0
3	G	702	04J	2	0
4	G	703	NDP	3	0
5	G	704	1UE	7	0
2	H	701	UMP	5	0
3	H	702	04J	14	0
4	H	703	NDP	15	0
5	H	704	1UE	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	489/610 (80%)	0.34	15 (3%)	52	40	51, 71, 125, 144	0
1	B	491/610 (80%)	0.44	18 (3%)	45	33	48, 69, 118, 180	0
1	C	489/610 (80%)	0.32	16 (3%)	50	38	51, 71, 125, 148	0
1	D	491/610 (80%)	0.41	20 (4%)	41	29	50, 70, 119, 180	0
1	E	489/610 (80%)	0.31	20 (4%)	41	29	53, 72, 125, 146	0
1	F	491/610 (80%)	0.38	22 (4%)	37	26	48, 70, 120, 178	0
1	G	489/610 (80%)	0.37	21 (4%)	39	27	50, 70, 117, 170	0
1	H	491/610 (80%)	0.31	22 (4%)	37	26	53, 72, 126, 144	0
All	All	3920/4880 (80%)	0.36	154 (3%)	43	31	48, 71, 124, 180	0

The worst 5 of 154 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	300	GLU	11.3
1	B	610	VAL	9.1
1	F	300	GLU	7.9
1	D	300	GLU	7.6
1	B	308	LYS	6.9

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	1UE	F	704	22/22	0.69	0.51	4.08	127,153,164,169	0
5	1UE	E	704	22/22	0.73	0.39	3.59	115,130,146,151	0
2	UMP	D	701	20/20	0.88	0.29	2.51	114,125,138,148	0
5	1UE	G	704	22/22	0.71	0.42	2.50	120,150,162,169	0
5	1UE	H	704	22/22	0.78	0.34	2.39	117,133,144,149	0
3	04J	B	702	32/32	0.87	0.32	2.22	96,128,152,156	0
3	04J	D	702	32/32	0.86	0.31	2.19	98,122,153,155	0
5	1UE	B	704	22/22	0.79	0.41	2.11	112,144,163,169	0
5	1UE	D	704	22/22	0.77	0.36	1.96	110,153,164,166	0
3	04J	E	702	32/32	0.90	0.39	1.95	89,134,162,165	0
2	UMP	H	701	20/20	0.87	0.24	1.82	107,132,138,138	0
5	1UE	C	704	22/22	0.80	0.28	1.64	106,130,152,161	0
3	04J	C	702	32/32	0.89	0.39	1.54	101,142,157,165	0
3	04J	A	702	32/32	0.89	0.33	1.42	95,139,161,161	0
2	UMP	G	701	20/20	0.83	0.25	1.36	110,127,134,138	0
3	04J	F	702	32/32	0.87	0.28	1.33	95,122,152,157	0
2	UMP	B	701	20/20	0.90	0.25	1.18	106,122,134,136	0
5	1UE	A	704	22/22	0.77	0.26	1.14	110,125,139,154	0
3	04J	G	702	32/32	0.88	0.26	1.04	94,122,156,162	0
2	UMP	F	701	20/20	0.88	0.23	1.00	113,125,142,151	0
2	UMP	E	701	20/20	0.93	0.24	0.94	95,120,129,130	0
3	04J	H	702	32/32	0.90	0.32	0.92	98,136,154,156	0
4	NDP	B	703	48/48	0.94	0.23	0.84	81,120,132,139	0
2	UMP	A	701	20/20	0.91	0.21	0.44	103,124,135,135	0
4	NDP	E	703	48/48	0.93	0.21	0.28	112,149,190,194	0
2	UMP	C	701	20/20	0.90	0.18	0.13	93,116,131,135	0
4	NDP	G	703	48/48	0.94	0.19	0.13	91,123,130,133	0
4	NDP	A	703	48/48	0.94	0.20	0.09	118,149,185,188	0
4	NDP	F	703	48/48	0.94	0.19	0.06	87,120,132,135	0
4	NDP	D	703	48/48	0.93	0.19	-0.00	93,117,133,137	0
4	NDP	H	703	48/48	0.91	0.18	-0.11	116,151,181,187	0
4	NDP	C	703	48/48	0.93	0.18	-0.14	117,144,185,186	0

6.5 Other polymers [i](#)

There are no such residues in this entry.