



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 26, 2016 – 11:03 AM EDT

PDB ID : 5L0E
Title : Crystal Structure of Autotaxin and Compound 1
Authors : Durbin, J.D.
Deposited on : 2016-07-27
Resolution : 3.06 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1191 (3.10-3.02)
Clashscore	102246	1303 (3.10-3.02)
Ramachandran outliers	100387	1254 (3.10-3.02)
Sidechain outliers	100360	1254 (3.10-3.02)
RSRZ outliers	91569	1197 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	871	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>5%</div> <div>10%</div> </div> </div>
1	B	871	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>6%</div> <div>10%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MAN	A	909	-	-	-	X
4	MAN	B	905	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 12992 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ectonucleotide pyrophosphatase/phosphodiesterase family member 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	783	Total	C	N	O	S	0	0	0
			6319	4001	1088	1181	49			
1	B	782	Total	C	N	O	S	0	0	0
			6310	3995	1086	1180	49			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	591	THR	ARG	engineered mutation	UNP Q64610
A	592	GLU	LYS	engineered mutation	UNP Q64610
A	863	ALA	-	expression tag	UNP Q64610
A	864	ALA	-	expression tag	UNP Q64610
A	865	ALA	-	expression tag	UNP Q64610
A	866	HIS	-	expression tag	UNP Q64610
A	867	HIS	-	expression tag	UNP Q64610
A	868	HIS	-	expression tag	UNP Q64610
A	869	HIS	-	expression tag	UNP Q64610
A	870	HIS	-	expression tag	UNP Q64610
A	871	HIS	-	expression tag	UNP Q64610
B	591	THR	ARG	engineered mutation	UNP Q64610
B	592	GLU	LYS	engineered mutation	UNP Q64610
B	863	ALA	-	expression tag	UNP Q64610
B	864	ALA	-	expression tag	UNP Q64610
B	865	ALA	-	expression tag	UNP Q64610
B	866	HIS	-	expression tag	UNP Q64610
B	867	HIS	-	expression tag	UNP Q64610
B	868	HIS	-	expression tag	UNP Q64610
B	869	HIS	-	expression tag	UNP Q64610
B	870	HIS	-	expression tag	UNP Q64610
B	871	HIS	-	expression tag	UNP Q64610

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is BETA-D-MANNOSE (three-letter code: BMA) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	B	1	Total	C	O	0	0
			11	6	5		
4	B	1	Total	C	O	0	0
			11	6	5		
4	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is 6-(3-{2-[(2,3-dihydro-1H-inden-2-yl)amino]-7,8-dihydropyrido[4,3-d]pyrimidin-6(5H)-yl}propanoyl)-1,3-benzoxazol-2(3H)-one (three-letter code: 6ZN) (formula: C₂₆H₂₅N₅O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			34	26	5	3		
5	B	1	Total	C	N	O	0	0
			34	26	5	3		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total 1	Cl 1	0	0
6	A	1	Total 1	Cl 1	0	0

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	3	Total 3	Zn 3	0	0
7	A	3	Total 3	Zn 3	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	40	Total 40	O 40	0	0
8	B	42	Total 42	O 42	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	184.18Å 96.84Å 136.92Å 90.00° 112.82° 90.00°	Depositor
Resolution (Å)	19.98 – 3.06 19.98 – 3.06	Depositor EDS
% Data completeness (in resolution range)	95.0 (19.98-3.06) 95.0 (19.98-3.06)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.42 (at 3.04Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.183 , 0.211 0.197 , 0.228	Depositor DCC
R_{free} test set	983 reflections (2.47%)	DCC
Wilson B-factor (Å ²)	48.7	Xtriage
Anisotropy	0.626	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 65.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	12992	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BMA, NAG, CL, 6ZN, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/6497	0.68	0/8814
1	B	0.51	0/6488	0.69	0/8803
All	All	0.51	0/12985	0.68	0/17617

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6319	0	6049	20	0
1	B	6310	0	6036	21	0
2	A	42	0	37	0	0
2	B	42	0	37	0	0
3	A	11	0	8	0	0
3	B	11	0	9	0	0
4	A	66	0	56	0	0
4	B	33	0	28	0	0
5	A	34	0	0	1	0
5	B	34	0	0	0	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	1	0	0	0	0
7	A	3	0	0	0	0
7	B	3	0	0	0	0
8	A	40	0	0	0	0
8	B	42	0	0	1	0
All	All	12992	0	12260	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 40 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:163:PRO:HB3	1:A:350:CYS:O	1.87	0.75
1:B:77:ASN:OD1	1:B:272:THR:HG21	1.99	0.62
1:A:243:LEU:HD22	5:A:911:6ZN:C7	2.31	0.61
1:A:171:ASP:O	1:A:310:PRO:HD2	2.03	0.58
1:B:171:ASP:O	1:B:310:PRO:HD2	2.05	0.56

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	777/871 (89%)	745 (96%)	31 (4%)	1 (0%)	56	88
1	B	776/871 (89%)	743 (96%)	33 (4%)	0	100	100
All	All	1553/1742 (89%)	1488 (96%)	64 (4%)	1 (0%)	56	88

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	590	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	710/786 (90%)	699 (98%)	11 (2%)	72	90
1	B	709/786 (90%)	695 (98%)	14 (2%)	63	87
All	All	1419/1572 (90%)	1394 (98%)	25 (2%)	66	88

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	124	LEU
1	B	162	ARG
1	B	659	SER
1	B	161	VAL
1	B	183	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	118	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 27 ligands modelled in this entry, 8 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	901	1,2	14,14,15	0.35	0	15,19,21	0.63	0
2	NAG	A	902	3,2	14,14,15	0.31	0	15,19,21	0.65	0
3	BMA	A	903	2,4	11,11,12	0.75	0	15,15,17	1.24	1 (6%)
4	MAN	A	904	3,4	11,11,12	0.47	0	15,15,17	0.92	2 (13%)
4	MAN	A	905	4	11,11,12	0.44	0	15,15,17	0.87	1 (6%)
4	MAN	A	906	4	11,11,12	0.40	0	15,15,17	0.77	1 (6%)
4	MAN	A	907	3	11,11,12	0.37	0	15,15,17	0.68	1 (6%)
4	MAN	A	908	4	11,11,12	0.35	0	15,15,17	0.85	1 (6%)
4	MAN	A	909	4	11,11,12	0.34	0	15,15,17	0.64	0
2	NAG	A	910	1	14,14,15	0.31	0	15,19,21	0.54	0
5	6ZN	A	911	7	35,39,39	1.53	4 (11%)	42,56,56	2.13	16 (38%)
2	NAG	B	901	1,2	14,14,15	0.38	0	15,19,21	0.66	0
2	NAG	B	902	3,2	14,14,15	0.28	0	15,19,21	0.67	0
3	BMA	B	903	2,4	11,11,12	0.78	0	15,15,17	1.17	0
4	MAN	B	904	3,4	11,11,12	0.43	0	15,15,17	0.95	1 (6%)
4	MAN	B	905	4	11,11,12	0.43	0	15,15,17	0.85	1 (6%)
4	MAN	B	906	4	11,11,12	0.38	0	15,15,17	0.81	1 (6%)
2	NAG	B	907	1	14,14,15	0.28	0	15,19,21	0.51	0
5	6ZN	B	908	7	35,39,39	1.40	3 (8%)	42,56,56	2.30	10 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	901	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	902	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	903	2,4	-	0/2/19/22	0/1/1/1
4	MAN	A	904	3,4	-	0/2/19/22	0/1/1/1
4	MAN	A	905	4	-	0/2/19/22	0/1/1/1
4	MAN	A	906	4	-	0/2/19/22	0/1/1/1
4	MAN	A	907	3	-	0/2/19/22	0/1/1/1
4	MAN	A	908	4	-	0/2/19/22	0/1/1/1
4	MAN	A	909	4	-	0/2/19/22	0/1/1/1
2	NAG	A	910	1	-	0/6/23/26	0/1/1/1
5	6ZN	A	911	7	-	0/13/30/30	0/5/6/6
2	NAG	B	901	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	902	3,2	-	0/6/23/26	0/1/1/1
3	BMA	B	903	2,4	-	0/2/19/22	0/1/1/1
4	MAN	B	904	3,4	-	0/2/19/22	0/1/1/1
4	MAN	B	905	4	-	0/2/19/22	0/1/1/1
4	MAN	B	906	4	-	0/2/19/22	0/1/1/1
2	NAG	B	907	1	-	0/6/23/26	0/1/1/1
5	6ZN	B	908	7	-	0/13/30/30	0/5/6/6

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	911	6ZN	C20-C11	-2.23	1.47	1.50
5	A	911	6ZN	C6-C13	-2.08	1.38	1.41
5	B	908	6ZN	C6-C13	-2.07	1.38	1.41
5	A	911	6ZN	O32-C17	2.78	1.22	1.19
5	B	908	6ZN	O32-C17	2.79	1.22	1.19

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	908	6ZN	C16-N30-C24	-8.13	114.46	123.51
5	A	911	6ZN	C16-N30-C24	-6.34	116.45	123.51
5	A	911	6ZN	C26-C25-C18	-4.39	108.01	112.79
5	B	908	6ZN	C11-C20-C24	-4.35	98.98	103.00
5	B	908	6ZN	N30-C16-N27	-4.34	110.89	116.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	911	6ZN	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	783/871 (89%)	-0.29	21 (2%) 58 32	15, 47, 85, 105	0
1	B	782/871 (89%)	-0.28	14 (1%) 71 47	18, 46, 81, 113	0
All	All	1565/1742 (89%)	-0.28	35 (2%) 65 40	15, 46, 83, 113	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	589	GLY	5.7
1	B	52	THR	4.7
1	B	396	SER	4.2
1	A	54	THR	4.2
1	B	590	SER	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	MAN	B	905	11/12	0.84	0.33	4.20	96,99,100,101	0
4	MAN	A	909	11/12	0.86	0.31	2.50	95,97,100,101	0
2	NAG	A	910	14/15	0.83	0.30	1.64	65,71,75,76	0
2	NAG	B	907	14/15	0.82	0.34	1.13	82,86,90,90	0
4	MAN	A	905	11/12	0.84	0.24	1.10	68,73,75,76	0
2	NAG	A	901	14/15	0.96	0.18	0.36	39,46,53,54	0
5	6ZN	A	911	34/34	0.95	0.17	0.22	27,47,61,61	0
5	6ZN	B	908	34/34	0.97	0.14	-0.52	29,36,56,59	0
2	NAG	B	901	14/15	0.97	0.14	-0.53	32,37,44,53	0
6	CL	B	912	1/1	0.98	0.15	-0.78	64,64,64,64	0
6	CL	A	912	1/1	0.94	0.10	-2.05	51,51,51,51	0
7	ZN	A	914	1/1	0.99	0.09	-2.18	30,30,30,30	0
7	ZN	B	910	1/1	0.97	0.07	-3.48	89,89,89,89	0
7	ZN	B	911	1/1	0.98	0.07	-3.54	74,74,74,74	0
7	ZN	A	915	1/1	0.99	0.03	-3.66	72,72,72,72	0
7	ZN	A	913	1/1	0.98	0.04	-6.68	61,61,61,61	1
7	ZN	B	909	1/1	1.00	0.07	-6.84	35,35,35,35	0
4	MAN	A	904	11/12	0.92	0.17	-	68,71,76,81	0
4	MAN	A	908	11/12	0.85	0.32	-	88,95,98,98	0
4	MAN	B	904	11/12	0.90	0.27	-	81,86,89,93	0
4	MAN	A	907	11/12	0.85	0.41	-	82,83,84,85	0
3	BMA	B	903	11/12	0.83	0.35	-	86,89,90,91	0
2	NAG	A	902	14/15	0.94	0.25	-	60,64,70,74	0
4	MAN	A	906	11/12	0.92	0.15	-	61,65,67,67	0
4	MAN	B	906	11/12	0.87	0.33	-	88,91,92,94	0
3	BMA	A	903	11/12	0.91	0.32	-	75,77,78,80	0
2	NAG	B	902	14/15	0.88	0.25	-	55,64,73,81	0

6.5 Other polymers [i](#)

There are no such residues in this entry.