



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 26, 2016 – 10:52 AM EDT

PDB ID : 5L0K
Title : Crystal Structure of Autotaxin and Compound PF-8380
Authors : Durbin, J.D.
Deposited on : 2016-07-27
Resolution : 2.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3050 (2.78-2.70)
Clashscore	102246	3424 (2.78-2.70)
Ramachandran outliers	100387	3367 (2.78-2.70)
Sidechain outliers	100360	3368 (2.78-2.70)
RSRZ outliers	91569	3055 (2.78-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	809	<div> <div>2%</div> <div> <div></div> <div>91%</div> <div>5%</div> <div>.</div> </div> </div>
1	B	809	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>7%</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	EPE	B	908	-	-	-	X
7	CL	B	909	-	-	-	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 12625 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ectonucleotide pyrophosphatase/phosphodiesterase family member 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	777	Total	C	N	O	S	0	0	0
			6174	3919	1051	1155	49			
1	B	777	Total	C	N	O	S	0	0	0
			6171	3923	1044	1155	49			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	53	ALA	ASN	engineered mutation	UNP Q64610
A	398	ALA	ASN	engineered mutation	UNP Q64610
A	410	ALA	ASN	engineered mutation	UNP Q64610
A	591	THR	ARG	engineered mutation	UNP Q64610
A	592	GLU	LYS	engineered mutation	UNP Q64610
A	806	ALA	ASN	engineered mutation	UNP Q64610
B	53	ALA	ASN	engineered mutation	UNP Q64610
B	398	ALA	ASN	engineered mutation	UNP Q64610
B	410	ALA	ASN	engineered mutation	UNP Q64610
B	591	THR	ARG	engineered mutation	UNP Q64610
B	592	GLU	LYS	engineered mutation	UNP Q64610
B	806	ALA	ASN	engineered mutation	UNP Q64610

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Zn	0	0
			2	2		
3	A	2	Total	Zn	0	0
			2	2		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is (3,5-dichlorophenyl)methyl 4-[3-oxo-3-(2-oxo-2,3-dihydro-1,3-benzoxazol-6-yl)propyl]piperazine-1-carboxylate (three-letter code: 6ZO) (formula: C₂₂H₂₁Cl₂N₃O₅).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Cl	N	O	0	0
			32	22	2	3	5		
5	B	1	Total	C	Cl	N	O	0	0
			32	22	2	3	5		

- Molecule 6 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			12	6	2	3	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	B	1	Total	C	N	O	S	0	0
			12	6	2	3	1		

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	1	Total	Cl	0	0
			1	1		
7	A	1	Total	Cl	0	0
			1	1		

- Molecule 8 is BETA-D-MANNOSE (three-letter code: BMA) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			11	6	5		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	65	Total	O	0	0
			65	65		
9	B	52	Total	O	0	0
			52	52		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ectonucleotide pyrophosphatase/phosphodiesterase family member 2



- Molecule 1: Ectonucleotide pyrophosphatase/phosphodiesterase family member 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	95.92Å 121.42Å 156.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	96.23 – 2.73 48.74 – 2.73	Depositor EDS
% Data completeness (in resolution range)	98.3 (96.23-2.73) 98.4 (48.74-2.73)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.02 (at 2.73Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.226 , 0.270 0.226 , 0.267	Depositor DCC
R_{free} test set	826 reflections (1.74%)	DCC
Wilson B-factor (Å ²)	35.7	Xtriage
Anisotropy	0.481	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	12625	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.41 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.6846e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BMA, 6ZO, NAG, CL, CA, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/6352	0.57	0/8636
1	B	0.42	0/6348	0.56	0/8633
All	All	0.42	0/12700	0.56	0/17269

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6174	0	5810	27	0
1	B	6171	0	5812	34	0
2	A	28	0	25	0	0
2	B	28	0	24	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	32	0	0	0	0
5	B	32	0	0	0	0
6	A	12	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	12	0	12	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
8	B	11	0	10	0	0
9	A	65	0	0	0	0
9	B	52	0	0	0	0
All	All	12625	0	11705	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:372:LEU:HD22	1:B:384:LEU:HD22	1.65	0.78
1:B:78:LEU:HD12	1:B:272:THR:HG21	1.64	0.76
1:B:546:GLU:HB3	1:B:602:ALA:HB1	1.67	0.75
1:A:635:ILE:CD1	1:A:685:LEU:HD13	2.17	0.74
1:A:635:ILE:HD13	1:A:685:LEU:HD13	1.70	0.72
1:A:711:VAL:HG13	1:A:815:VAL:HG13	1.77	0.67
1:A:166:ILE:HD12	1:A:351:VAL:HG11	1.84	0.59
1:B:206:PRO:HB3	1:B:389:LEU:HD22	1.87	0.56
1:B:78:LEU:HD11	1:B:274:PHE:HB2	1.88	0.56
1:A:206:PRO:HB3	1:A:389:LEU:HD22	1.88	0.55
1:A:682:PRO:HG3	1:A:685:LEU:HD12	1.89	0.55
1:B:641:VAL:HG22	1:B:685:LEU:CD2	2.39	0.53
1:A:635:ILE:HD11	1:A:685:LEU:HD13	1.91	0.53
1:A:78:LEU:HD12	1:A:272:THR:HG21	1.91	0.53
1:B:394:ALA:HB2	1:B:402:TYR:CZ	2.44	0.51
1:A:493:THR:HG21	1:A:515:LEU:HD23	1.92	0.50
1:A:166:ILE:CD1	1:A:351:VAL:HG11	2.41	0.50
1:A:259:LEU:HD22	1:A:512:MET:CE	2.42	0.49
1:B:724:ALA:HA	1:B:730:VAL:HG13	1.94	0.49
1:B:394:ALA:HB2	1:B:402:TYR:CE2	2.48	0.49
1:B:711:VAL:HG13	1:B:815:VAL:HG13	1.93	0.49
1:B:535:ARG:HG2	1:B:536:THR:HG23	1.94	0.48
1:A:793:PHE:CD1	1:A:795:LEU:HD13	2.49	0.48
1:B:493:THR:HG21	1:B:515:LEU:HD23	1.96	0.48
1:B:259:LEU:HD22	1:B:512:MET:CE	2.44	0.47
1:A:526:THR:CG2	1:A:849:LEU:HD22	2.43	0.47
1:B:144:VAL:HG21	1:B:186:PRO:HB2	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:682:PRO:CG	1:A:685:LEU:HD12	2.45	0.47
1:B:823:THR:HG23	1:B:855:LEU:HD11	1.97	0.46
1:B:328:LEU:HA	1:B:331:ILE:HD12	1.96	0.46
1:B:711:VAL:CG1	1:B:795:LEU:HD21	2.45	0.46
1:B:388:THR:HG22	1:B:476:PHE:CZ	2.51	0.46
1:B:372:LEU:CD2	1:B:384:LEU:HD22	2.43	0.45
1:B:568:CYS:SG	1:B:569:ASP:N	2.90	0.45
1:A:682:PRO:HB3	1:A:716:GLN:HB3	2.00	0.44
1:A:313:SER:HB2	1:A:324:MET:HE1	1.99	0.43
1:A:715:PHE:O	1:A:720:VAL:HG23	2.18	0.43
1:B:641:VAL:HG22	1:B:685:LEU:HD23	2.00	0.43
1:A:210:PHE:HB3	1:A:227:ILE:HG21	2.00	0.43
1:B:830:GLU:OE2	1:B:852:LYS:NZ	2.51	0.43
1:B:648:LEU:HD21	1:B:693:TYR:CE1	2.53	0.43
1:B:794:ILE:HD13	1:B:832:LEU:CD1	2.48	0.43
1:B:166:ILE:HD12	1:B:351:VAL:HG11	2.00	0.43
1:B:641:VAL:HG22	1:B:685:LEU:HD21	2.01	0.43
1:B:675:MET:HE3	1:B:763:VAL:HG13	2.01	0.43
1:A:555:ILE:HG23	1:A:656:VAL:CG2	2.49	0.42
1:A:259:LEU:HD22	1:A:512:MET:HE3	2.02	0.42
1:A:431:ARG:HD3	1:A:823:THR:O	2.19	0.42
1:B:526:THR:CG2	1:B:849:LEU:HD22	2.49	0.42
1:A:309:GLN:HG3	1:A:310:PRO:HA	2.01	0.42
1:B:711:VAL:HG11	1:B:795:LEU:HD21	2.02	0.42
1:A:222:PRO:HA	1:A:225:HIS:CE1	2.55	0.42
1:A:708:PHE:CE1	1:A:795:LEU:HD23	2.55	0.42
1:B:209:THR:HG22	1:B:213:LEU:HD12	2.02	0.42
1:B:686:SER:HB2	1:B:692:LYS:HZ3	1.85	0.42
1:A:268:VAL:HG21	1:A:518:LEU:HD21	2.01	0.42
1:B:408:ILE:O	1:B:412:THR:HG23	2.20	0.42
1:A:268:VAL:CG2	1:A:518:LEU:HD21	2.51	0.41
1:B:394:ALA:HB2	1:B:402:TYR:CE1	2.56	0.41
1:A:388:THR:HG22	1:A:476:PHE:CZ	2.56	0.41
1:B:682:PRO:HB3	1:B:716:GLN:HB3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	771/809 (95%)	747 (97%)	23 (3%)	1 (0%)	56	83
1	B	769/809 (95%)	740 (96%)	28 (4%)	1 (0%)	56	83
All	All	1540/1618 (95%)	1487 (97%)	51 (3%)	2 (0%)	56	83

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	435	ALA
1	B	435	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	676/731 (92%)	672 (99%)	4 (1%)	90	97
1	B	677/731 (93%)	674 (100%)	3 (0%)	93	98
All	All	1353/1462 (92%)	1346 (100%)	7 (0%)	92	97

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	309	GLN
1	A	757	GLU
1	A	785	ASP
1	A	795	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	667	LEU
1	B	785	ASP
1	B	800	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	481	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 8 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	901	1,2	14,14,15	0.58	0	15,19,21	0.90	1 (6%)
2	NAG	A	902	2	14,14,15	0.52	0	15,19,21	1.42	3 (20%)
5	6ZO	A	906	3	31,35,35	1.11	2 (6%)	40,49,49	2.48	6 (15%)
6	EPE	A	907	-	12,12,15	0.93	1 (8%)	13,16,20	2.20	3 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	901	1,2	14,14,15	0.55	0	15,19,21	0.75	0
2	NAG	B	902	8,2	14,14,15	0.52	0	15,19,21	1.27	2 (13%)
8	BMA	B	903	2	11,11,12	0.52	0	15,15,17	1.32	2 (13%)
5	6ZO	B	907	3	31,35,35	1.16	2 (6%)	40,49,49	2.47	6 (15%)
6	EPE	B	908	-	12,12,15	0.92	1 (8%)	13,16,20	2.03	4 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	901	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	902	2	-	0/6/23/26	0/1/1/1
5	6ZO	A	906	3	-	0/18/28/28	0/3/4/4
6	EPE	A	907	-	-	0/6/14/19	0/1/1/1
2	NAG	B	901	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	902	8,2	-	0/6/23/26	0/1/1/1
8	BMA	B	903	2	-	0/2/19/22	0/1/1/1
5	6ZO	B	907	3	-	0/18/28/28	0/3/4/4
6	EPE	B	908	-	-	0/6/14/19	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	907	6ZO	C27-C28	-2.45	1.37	1.41
5	A	906	6ZO	C27-C28	-2.00	1.38	1.41
5	A	906	6ZO	C9-N11	2.65	1.39	1.35
5	B	907	6ZO	C9-N11	2.69	1.39	1.35
6	B	908	EPE	C10-S	2.74	1.81	1.77
6	A	907	EPE	C10-S	2.85	1.81	1.77

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	907	6ZO	O8-C9-O10	-3.69	118.56	124.82
5	A	906	6ZO	O8-C9-O10	-3.54	118.82	124.82
5	A	906	6ZO	O10-C9-N11	-3.50	118.90	124.39
5	B	907	6ZO	O10-C9-N11	-3.21	119.35	124.39
5	B	907	6ZO	C25-C27-C28	-2.86	117.71	120.86
8	B	903	BMA	O5-C1-C2	-2.79	106.44	110.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	906	6ZO	C25-C27-C28	-2.75	117.83	120.86
5	A	906	6ZO	O8-C7-C1	2.06	114.62	109.39
2	B	902	NAG	C1-O5-C5	2.08	115.19	112.14
6	B	908	EPE	O3S-S-C10	2.10	109.36	104.99
2	A	902	NAG	C2-N2-C7	2.24	126.03	123.11
5	B	907	6ZO	C4-C6-C5	2.25	119.95	117.27
2	A	901	NAG	C1-O5-C5	2.26	115.46	112.14
5	A	906	6ZO	C4-C6-C5	2.27	119.98	117.27
5	B	907	6ZO	O8-C7-C1	2.33	115.31	109.39
6	B	908	EPE	C5-N4-C3	2.38	118.40	110.45
2	A	902	NAG	O5-C5-C4	2.47	114.23	110.13
8	B	903	BMA	C1-O5-C5	2.54	115.87	112.14
6	A	907	EPE	C5-N4-C3	2.55	118.97	110.45
2	B	902	NAG	C2-N2-C7	2.61	126.50	123.11
6	B	908	EPE	O1S-S-C10	2.91	108.93	106.87
6	A	907	EPE	O2S-S-C10	2.99	108.98	106.87
2	A	902	NAG	C1-O5-C5	3.00	116.56	112.14
6	B	908	EPE	O2S-S-C10	4.99	110.40	106.87
6	A	907	EPE	O1S-S-C10	6.18	111.23	106.87
5	B	907	6ZO	O8-C9-N11	13.16	122.09	111.61
5	A	906	6ZO	O8-C9-N11	13.40	122.28	111.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	777/809 (96%)	0.14	18 (2%) 64 66	23, 36, 56, 69	0
1	B	777/809 (96%)	0.18	16 (2%) 67 69	24, 40, 60, 76	0
All	All	1554/1618 (96%)	0.16	34 (2%) 65 68	23, 38, 59, 76	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	537	ASN	4.5
1	B	71	PRO	4.1
1	B	460	VAL	4.0
1	A	469	PHE	3.5
1	B	560	SER	3.5
1	B	459	ASP	3.4
1	B	161	VAL	3.4
1	A	537	ASN	2.9
1	A	594	ARG	2.7
1	B	158	ALA	2.7
1	A	107	CYS	2.7
1	A	397	ILE	2.6
1	A	53	ALA	2.4
1	A	396	SER	2.4
1	B	547	VAL	2.4
1	A	125	SER	2.4
1	A	134	GLN	2.3
1	B	625	PHE	2.3
1	B	562	PHE	2.3
1	A	619	SER	2.3
1	B	556	MET	2.2
1	A	536	THR	2.2
1	A	400	SER	2.1
1	B	136	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	412	THR	2.1
1	B	400	SER	2.1
1	B	417	ASP	2.1
1	A	693	TYR	2.1
1	A	468	CYS	2.1
1	A	591	THR	2.1
1	A	242	HIS	2.0
1	B	536	THR	2.0
1	A	70	PRO	2.0
1	A	401	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	EPE	B	908	12/15	0.89	0.36	2.38	60,67,74,74	0
7	CL	B	909	1/1	0.97	0.24	2.20	50,50,50,50	0
6	EPE	A	907	12/15	0.96	0.27	1.91	49,60,69,70	0
7	CL	A	908	1/1	0.96	0.19	1.55	34,34,34,34	0
5	6ZO	B	907	32/32	0.90	0.23	1.42	47,59,62,63	0
5	6ZO	A	906	32/32	0.92	0.21	1.31	51,54,58,60	0
2	NAG	A	901	14/15	0.94	0.19	1.17	31,35,39,47	0
2	NAG	B	901	14/15	0.94	0.15	-0.22	41,44,46,50	0
3	ZN	A	903	1/1	0.99	0.12	-1.61	37,37,37,37	0
4	CA	B	906	1/1	0.98	0.07	-2.22	39,39,39,39	0
4	CA	A	905	1/1	0.98	0.09	-2.31	33,33,33,33	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ZN	B	904	1/1	0.99	0.13	-2.88	37,37,37,37	0
3	ZN	B	905	1/1	0.95	0.06	-3.72	63,63,63,63	0
3	ZN	A	904	1/1	0.96	0.05	-5.47	54,54,54,54	0
2	NAG	B	902	14/15	0.88	0.29	-	56,61,65,72	0
8	BMA	B	903	11/12	0.75	0.34	-	78,82,84,84	0
2	NAG	A	902	14/15	0.82	0.33	-	53,58,60,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.