



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 26, 2016 – 05:08 PM EDT

PDB ID : 5L25  
Title : Crystal Structure of Arabidopsis thaliana Bor1  
Authors : Thurtle-Schmidt, B.H.; Stroud, R.M.  
Deposited on : 2016-07-31  
Resolution : 4.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 4.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1020 (4.62-3.60)
Clashscore	102246	1119 (4.62-3.60)
Ramachandran outliers	100387	1065 (4.62-3.60)
Sidechain outliers	100360	1051 (4.62-3.60)
RSRZ outliers	91569	1024 (4.62-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	652	<div> <div style="width: 44%; background-color: green;"></div> <div style="width: 17%; background-color: yellow;"></div> <div style="width: 39%; background-color: grey;"></div> </div> <div>44% 17% . 39%</div>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2873 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Boron transporter 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	399	Total	C	N	O	S	0	0	0
			2873	1897	461	495	20			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	646	ALA	-	expression tag	UNP Q8VYR7
A	647	ALA	-	expression tag	UNP Q8VYR7
A	648	ALA	-	expression tag	UNP Q8VYR7
A	649	LEU	-	expression tag	UNP Q8VYR7
A	650	VAL	-	expression tag	UNP Q8VYR7
A	651	PRO	-	expression tag	UNP Q8VYR7
A	652	ARG	-	expression tag	UNP Q8VYR7



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 4 <sub>2</sub> 2 <sub>1</sub> 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	184.56 Å   184.56 Å   89.83 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	20.07 – 4.11 20.07 – 4.11	Depositor EDS
% Data completeness (in resolution range)	94.8 (20.07-4.11) 95.8 (20.07-4.11)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.80 (at 4.07 Å)	Xtriage
Refinement program	REFMAC 5.8.0151	Depositor
R, $R_{free}$	0.359   ,   0.391 0.360   ,   0.393	Depositor DCC
$R_{free}$ test set	565 reflections (4.99%)	DCC
Wilson B-factor (Å <sup>2</sup> )	223.8	Xtriage
Anisotropy	0.296	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.17 , 218.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	2873	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	258.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.59	1/2935 (0.0%)	0.68	5/4005 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	471	SER	CB-OG	5.49	1.49	1.42

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	194	PRO	N-CA-CB	8.28	113.23	103.30
1	A	539	PRO	N-CA-CB	6.45	111.05	103.30
1	A	254	PRO	N-CA-CB	5.88	110.36	103.30
1	A	182	PRO	N-CA-CB	5.75	110.20	103.30
1	A	482	PRO	N-CA-CB	5.05	109.36	103.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2873	0	2721	95	0
All	All	2873	0	2721	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 17.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:553:TYR:CE1	1:A:577:LEU:HD12	1.46	1.50
1:A:553:TYR:OH	1:A:574:ILE:N	1.62	1.32
1:A:553:TYR:CD1	1:A:577:LEU:CD1	2.35	1.09
1:A:553:TYR:OH	1:A:573:MET:C	1.91	1.09
1:A:553:TYR:CD1	1:A:577:LEU:HD12	1.88	1.09
1:A:553:TYR:CE1	1:A:577:LEU:CD1	2.36	1.08
1:A:553:TYR:CD1	1:A:577:LEU:HD13	2.08	0.89
1:A:546:PHE:CD1	1:A:549:PHE:HD1	1.90	0.89
1:A:553:TYR:OH	1:A:574:ILE:CA	2.20	0.89
1:A:546:PHE:CD1	1:A:549:PHE:CD1	2.64	0.86
1:A:553:TYR:HE1	1:A:577:LEU:HD12	1.05	0.85
1:A:169:GLN:O	1:A:172:ILE:HG12	1.78	0.83
1:A:480:ALA:O	1:A:484:LEU:CB	2.26	0.83
1:A:553:TYR:HD1	1:A:577:LEU:CD1	1.92	0.82
1:A:546:PHE:O	1:A:548:LEU:N	2.15	0.79
1:A:553:TYR:HH	1:A:574:ILE:N	1.81	0.78
1:A:553:TYR:HD1	1:A:577:LEU:HD13	1.46	0.78
1:A:93:LEU:HD23	1:A:94:GLY:N	2.01	0.75
1:A:54:PHE:HB3	1:A:100:VAL:HG21	1.69	0.75
1:A:34:PHE:HA	1:A:37:LEU:HD12	1.73	0.70
1:A:553:TYR:OH	1:A:574:ILE:HA	1.90	0.70
1:A:91:LEU:O	1:A:93:LEU:N	2.25	0.70
1:A:169:GLN:O	1:A:172:ILE:CG1	2.40	0.69
1:A:574:ILE:HG23	1:A:578:ILE:HD12	1.75	0.67
1:A:201:GLY:O	1:A:205:LEU:HB2	1.97	0.65
1:A:78:CYS:C	1:A:93:LEU:HD11	2.16	0.65
1:A:56:GLU:HA	1:A:59:GLU:HB2	1.78	0.65
1:A:291:ILE:O	1:A:294:ILE:HG22	1.97	0.64
1:A:546:PHE:O	1:A:547:THR:C	2.36	0.64
1:A:57:GLN:HE21	1:A:496:PHE:HE1	1.47	0.63
1:A:560:LEU:HB3	1:A:566:ALA:HB1	1.78	0.63
1:A:552:THR:O	1:A:556:ILE:HG13	1.98	0.63
1:A:577:LEU:O	1:A:581:ARG:HG3	1.98	0.62
1:A:214:THR:HG1	1:A:247:TRP:HE1	1.43	0.62
1:A:553:TYR:CE2	1:A:578:ILE:HD11	2.36	0.61
1:A:163:ILE:O	1:A:167:PHE:HB2	2.01	0.60
1:A:174:GLY:HA3	1:A:493:TRP:CD2	2.36	0.60
1:A:357:VAL:HG12	1:A:359:PRO:HD2	1.83	0.60

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:477:CYS:HA	1:A:480:ALA:HB3	1.84	0.60
1:A:58:LEU:HD22	1:A:66:LEU:HD23	1.86	0.58
1:A:79:GLY:HA2	1:A:93:LEU:HD13	1.86	0.57
1:A:158:LEU:HD23	1:A:553:TYR:HD2	1.70	0.57
1:A:241:PRO:O	1:A:244:VAL:HB	2.04	0.57
1:A:200:ASN:HA	1:A:203:PHE:HB2	1.87	0.56
1:A:553:TYR:HE1	1:A:577:LEU:CD1	1.96	0.55
1:A:546:PHE:CZ	1:A:549:PHE:HB2	2.42	0.55
1:A:546:PHE:CE1	1:A:549:PHE:HB2	2.42	0.55
1:A:78:CYS:CB	1:A:93:LEU:HD11	2.39	0.52
1:A:137:PHE:CD2	1:A:337:LEU:HD11	2.44	0.52
1:A:337:LEU:O	1:A:339:LEU:N	2.43	0.52
1:A:546:PHE:C	1:A:548:LEU:N	2.60	0.52
1:A:93:LEU:HD23	1:A:93:LEU:C	2.31	0.51
1:A:569:MET:O	1:A:573:MET:HG2	2.10	0.50
1:A:79:GLY:N	1:A:93:LEU:HD11	2.28	0.49
1:A:79:GLY:CA	1:A:93:LEU:HD13	2.43	0.49
1:A:78:CYS:HB2	1:A:93:LEU:HD11	1.93	0.48
1:A:195:SER:HA	1:A:199:ALA:HB3	1.96	0.48
1:A:581:ARG:O	1:A:585:LEU:HG	2.13	0.48
1:A:174:GLY:HA3	1:A:493:TRP:CE3	2.49	0.48
1:A:38:ALA:HB3	1:A:39:PRO:HD3	1.96	0.48
1:A:123:ALA:O	1:A:126:GLY:N	2.48	0.47
1:A:504:LEU:HB2	1:A:505:PRO:HD3	1.97	0.47
1:A:546:PHE:CE1	1:A:549:PHE:CD1	3.03	0.47
1:A:221:ALA:HB3	1:A:236:ALA:HB2	1.97	0.46
1:A:172:ILE:HG13	1:A:173:LYS:N	2.30	0.46
1:A:169:GLN:HA	1:A:172:ILE:HG12	1.98	0.46
1:A:510:TRP:O	1:A:513:ILE:HG22	2.16	0.46
1:A:492:LEU:O	1:A:496:PHE:HD2	1.99	0.45
1:A:546:PHE:O	1:A:549:PHE:N	2.34	0.45
1:A:55:GLY:HA2	1:A:71:THR:OG1	2.17	0.45
1:A:211:LEU:HD21	1:A:495:TYR:CD1	2.52	0.44
1:A:492:LEU:HA	1:A:495:TYR:HD2	1.81	0.44
1:A:491:VAL:HG12	1:A:495:TYR:CE2	2.52	0.44
1:A:492:LEU:O	1:A:495:TYR:HB2	2.17	0.44
1:A:557:CYS:O	1:A:561:THR:HG23	2.18	0.43
1:A:79:GLY:N	1:A:93:LEU:CD1	2.81	0.43
1:A:308:TYR:CE1	1:A:355:ASN:HB3	2.54	0.43
1:A:58:LEU:HD21	1:A:100:VAL:HG13	2.01	0.43
1:A:233:SER:HA	1:A:236:ALA:HB3	2.01	0.43

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:304:ILE:HG23	1:A:308:TYR:CD2	2.54	0.43
1:A:307:LEU:HD12	1:A:571:PRO:HG3	2.01	0.43
1:A:495:TYR:O	1:A:496:PHE:C	2.57	0.43
1:A:50:PRO:HA	1:A:53:SER:HB3	2.01	0.43
1:A:139:LEU:HB2	1:A:309:TYR:OH	2.19	0.42
1:A:304:ILE:HG23	1:A:308:TYR:CE2	2.54	0.42
1:A:492:LEU:HD23	1:A:492:LEU:HA	1.86	0.42
1:A:127:TRP:HB3	1:A:294:ILE:HD11	2.02	0.42
1:A:514:LEU:HA	1:A:514:LEU:HD23	1.88	0.42
1:A:495:TYR:O	1:A:498:PHE:N	2.53	0.41
1:A:466:SER:HA	1:A:469:LEU:HD12	2.01	0.41
1:A:206:VAL:HA	1:A:209:PHE:CZ	2.56	0.41
1:A:300:PRO:O	1:A:304:ILE:HB	2.21	0.41
1:A:209:PHE:CD1	1:A:210:GLY:N	2.89	0.40
1:A:339:LEU:O	1:A:343:THR:HG23	2.21	0.40
1:A:224:TRP:O	1:A:232:ARG:NH1	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	383/652 (59%)	334 (87%)	40 (10%)	9 (2%)	8 50

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	92	ILE
1	A	203	PHE
1	A	338	LEU
1	A	535	VAL
1	A	82	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	117	GLY
1	A	369	ALA
1	A	193	LEU
1	A	194	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	265/553 (48%)	260 (98%)	5 (2%)	65 86

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	SER
1	A	127	TRP
1	A	167	PHE
1	A	562	TRP
1	A	583	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	399/652 (61%)	-0.78	1 (0%) 94 92	129, 249, 382, 500	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	142	CYS	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.