



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 27, 2016 – 12:56 AM BST

PDB ID : 2L3J
Title : The solution structure of the ADAR2 dsRBM-RNA complex reveals a sequence-specific read out of the minor groove
Authors : Stefl, R.; Oberstrass, F.C.; Allain, F.H.-T.
Deposited on : 2010-09-14

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958
RNA backbone	3027	600

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	236	
2	B	71	

2 Ensemble composition and analysis ⓘ

This entry contains 19 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:6-A:72 (67)	0.27	10
2	A:163-A:181, A:188-A:225 (57)	0.15	10

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 1 single-model cluster was found.

Cluster number	Models
1	2, 5, 6, 8, 15, 16, 19
2	3, 10, 12, 13
3	4, 9, 14
4	7, 17
5	11, 18
Single-model clusters	1

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5520 atoms, of which 2218 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Double-stranded RNA-specific editase 1.

Mol	Chain	Residues	Atoms						Trace
1	A	236	Total	C	H	N	O	S	0
			3251	1144	1456	307	338	6	

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	229	LEU	-	EXPRESSION TAG	UNP P51400
A	230	GLU	-	EXPRESSION TAG	UNP P51400
A	231	HIS	-	EXPRESSION TAG	UNP P51400
A	232	HIS	-	EXPRESSION TAG	UNP P51400
A	233	HIS	-	EXPRESSION TAG	UNP P51400
A	234	HIS	-	EXPRESSION TAG	UNP P51400
A	235	HIS	-	EXPRESSION TAG	UNP P51400
A	236	HIS	-	EXPRESSION TAG	UNP P51400

- Molecule 2 is a RNA chain called RNA (71-MER).


Mol	Chain	Residues	Atoms						Trace
2	B	71	Total	C	H	N	O	P	0
			2269	676	762	266	495	70	

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Double-stranded RNA-specific editase 1




- Molecule 2: RNA (71-MER)




4.2 Residue scores for the representative (medoid) model from the NMR ensemble



The representative model is number 10. Colouring as in section 4.1 above.

- Molecule 1: Double-stranded RNA-specific editase 1





● Molecule 2: RNA (71-MER)



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 40 calculated structures, 19 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
AMBER	refinement	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.53±0.01	0±0/977 (0.0±0.0%)	0.81±0.01	0±0/1314 (0.0±0.0%)
2	B	1.01±0.01	0±0/1685 (0.0±0.0%)	1.50±0.01	4±1/2623 (0.2±0.0%)
All	All	0.86	0/50578 (0.0%)	1.31	83/74803 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
2	B	0.0±0.0	0.3±0.5
All	All	0	5

There are no bond-length outliers.

5 of 23 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	274	A	N9-C1'-C2'	7.16	123.31	114.00	13	5
2	B	286	G	O4'-C1'-N9	6.81	113.65	108.20	7	18
2	B	259	U	O4'-C1'-N1	6.55	113.44	108.20	9	8
1	A	69	PHE	CB-CG-CD2	-5.97	116.62	120.80	16	1
2	B	241	U	O4'-C1'-N1	5.87	112.89	108.20	16	11

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
2	B	271	C	Sidechain	4
2	B	273	A	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	957	829	994	9±2
2	B	1507	762	762	2±2
All	All	46816	30229	33364	197

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

5 of 50 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:20:LEU:HD22	1:A:21:GLN:N	0.63	2.09	16	19
2:B:273:A:C2	2:B:274:A:C4	0.63	2.86	8	6
2:B:273:A:N3	2:B:274:A:C8	0.62	2.67	5	6
1:A:24:LEU:HD23	1:A:38:MET:HG2	0.57	1.76	14	11
1:A:42:VAL:HG11	1:A:69:PHE:CE2	0.56	2.36	2	5

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	124/236 (53%)	107±2 (86±2%)	13±2 (11±2%)	4±1 (3±1%)	9	43
All	All	2356/4484 (53%)	2032 (86%)	256 (11%)	68 (3%)	9	43

5 of 12 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	180	SER	16
1	A	53	THR	15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	31	VAL	13
1	A	43	ASN	7
1	A	19	GLY	6

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	103/200 (52%)	94±1 (91±1%)	9±1 (9±1%)	18	63
All	All	1957/3800 (52%)	1788 (91%)	169 (9%)	18	63

5 of 21 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	20	LEU	19
1	A	174	LEU	19
1	A	191	VAL	19
1	A	170	LEU	18
1	A	37	VAL	14

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
2	B	70/71 (99%)	2±1 (3±1%)	2±1 (2±1%)	0.72±0.01
All	All	1330/1349 (99%)	42 (3%)	32 (2%)	0.72

The overall RNA backbone suiteness is 0.72.

All unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
2	B	272	U	19
2	B	273	A	19
2	B	270	G	2
2	B	271	C	2

All unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
2	B	272	U	18
2	B	271	C	12
2	B	270	G	2

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided