



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:11 AM GMT

PDB ID : 3L34
Title : The crystal structure of a two-component sensor domain (2nd form) from *Pseudomonas aeruginosa* PA01
Authors : Tan, K.; Chhor, G.; Buck, K.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2009-12-16
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	130	
1	B	130	
1	C	130	
1	D	130	
1	E	130	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	130	<div><div></div><div>8%</div><div>75%</div><div>16%</div><div>7%</div></div>
1	G	130	<div><div></div><div>2%</div><div>81%</div><div>12%</div><div>7%</div></div>
1	H	130	<div><div></div><div>7%</div><div>75%</div><div>14%</div><div>10%</div></div>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 8232 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sensor protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	120	Total	C	N	O	Se	0	2	0
			949	581	178	188	2			
1	B	124	Total	C	N	O	Se	0	2	0
			981	598	184	197	2			
1	C	122	Total	C	N	O	Se	0	3	0
			967	592	181	192	2			
1	D	117	Total	C	N	O	Se	0	1	0
			920	564	172	182	2			
1	E	123	Total	C	N	O	Se	0	1	0
			963	587	179	195	2			
1	F	121	Total	C	N	O	Se	0	3	0
			963	589	181	191	2			
1	G	121	Total	C	N	O	Se	0	3	0
			960	586	180	192	2			
1	H	117	Total	C	N	O	Se	0	1	0
			926	567	174	183	2			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	SER	-	expression tag	UNP Q9HT87
A	40	ASN	-	expression tag	UNP Q9HT87
A	41	ALA	-	expression tag	UNP Q9HT87
B	39	SER	-	expression tag	UNP Q9HT87
B	40	ASN	-	expression tag	UNP Q9HT87
B	41	ALA	-	expression tag	UNP Q9HT87
C	39	SER	-	expression tag	UNP Q9HT87
C	40	ASN	-	expression tag	UNP Q9HT87
C	41	ALA	-	expression tag	UNP Q9HT87
D	39	SER	-	expression tag	UNP Q9HT87
D	40	ASN	-	expression tag	UNP Q9HT87
D	41	ALA	-	expression tag	UNP Q9HT87
E	39	SER	-	expression tag	UNP Q9HT87

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	40	ASN	-	expression tag	UNP Q9HT87
E	41	ALA	-	expression tag	UNP Q9HT87
F	39	SER	-	expression tag	UNP Q9HT87
F	40	ASN	-	expression tag	UNP Q9HT87
F	41	ALA	-	expression tag	UNP Q9HT87
G	39	SER	-	expression tag	UNP Q9HT87
G	40	ASN	-	expression tag	UNP Q9HT87
G	41	ALA	-	expression tag	UNP Q9HT87
H	39	SER	-	expression tag	UNP Q9HT87
H	40	ASN	-	expression tag	UNP Q9HT87
H	41	ALA	-	expression tag	UNP Q9HT87


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	83	Total O 83 83	0	0
2	B	81	Total O 81 81	0	0
2	C	90	Total O 90 90	0	0
2	D	69	Total O 69 69	0	0
2	E	78	Total O 78 78	0	0
2	F	74	Total O 74 74	0	0
2	G	80	Total O 80 80	0	0
2	H	48	Total O 48 48	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Sensor protein




- Molecule 1: Sensor protein




- Molecule 1: Sensor protein




- Molecule 1: Sensor protein




- Molecule 1: Sensor protein




- Molecule 1: Sensor protein



• Molecule 1: Sensor protein



• Molecule 1: Sensor protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.51Å 103.08Å 69.47Å 90.00° 90.01° 90.00°	Depositor
Resolution (Å)	41.39 – 1.70 42.04 – 1.70	Depositor EDS
% Data completeness (in resolution range)	92.7 (41.39-1.70) 98.8 (42.04-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 1.70Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
R, R_{free}	0.182 , 0.225 0.185 , 0.224	Depositor DCC
R_{free} test set	4709 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	24.5	Xtriage
Anisotropy	0.436	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.5	EDS
Estimated twinning fraction	0.117 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 93950 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8232	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/963	0.50	0/1299
1	B	0.33	0/995	0.44	0/1341
1	C	0.33	0/984	0.48	0/1326
1	D	0.34	0/931	0.48	0/1255
1	E	0.32	0/974	0.44	0/1313
1	F	0.33	0/980	0.47	0/1322
1	G	0.34	0/977	0.48	0/1316
1	H	0.31	0/937	0.48	0/1262
All	All	0.33	0/7741	0.47	0/10434

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	949	0	934	20	0
1	B	981	0	959	11	0
1	C	967	0	954	9	0
1	D	920	0	903	14	0
1	E	963	0	936	9	0
1	F	963	0	948	18	0
1	G	960	0	943	10	0
1	H	926	0	909	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	83	0	0	4	0
2	B	81	0	0	1	0
2	C	90	0	0	1	0
2	D	69	0	0	2	0
2	E	78	0	0	2	0
2	F	74	0	0	4	0
2	G	80	0	0	2	0
2	H	48	0	0	2	0
All	All	8232	0	7486	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:46:MSE:HA	1:B:46:MSE:HE2	1.55	0.87
1:A:139:PHE:CE1	1:B:67:LEU:HD11	2.14	0.83
1:E:39:SER:HB3	1:E:42:GLN:HB3	1.63	0.79
1:E:130:ALA:HA	1:F:129:MSE:HE1	1.63	0.78
1:A:84:GLN:HE22	1:A:121[B]:THR:HG21	1.48	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/130 (92%)	117 (98%)	2 (2%)	1 (1%)	24	7
1	B	124/130 (95%)	124 (100%)	0	0	100	100
1	C	123/130 (95%)	122 (99%)	1 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	116/130 (89%)	114 (98%)	2 (2%)	0	100	100
1	E	122/130 (94%)	122 (100%)	0	0	100	100
1	F	122/130 (94%)	120 (98%)	2 (2%)	0	100	100
1	G	122/130 (94%)	121 (99%)	1 (1%)	0	100	100
1	H	116/130 (89%)	114 (98%)	2 (2%)	0	100	100
All	All	965/1040 (93%)	954 (99%)	10 (1%)	1 (0%)	56	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	128	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	99/103 (96%)	94 (95%)	5 (5%)	29	10
1	B	103/103 (100%)	101 (98%)	2 (2%)	65	46
1	C	102/103 (99%)	96 (94%)	6 (6%)	24	7
1	D	96/103 (93%)	92 (96%)	4 (4%)	36	14
1	E	101/103 (98%)	98 (97%)	3 (3%)	48	26
1	F	101/103 (98%)	94 (93%)	7 (7%)	19	5
1	G	102/103 (99%)	99 (97%)	3 (3%)	50	27
1	H	97/103 (94%)	96 (99%)	1 (1%)	82	72
All	All	801/824 (97%)	770 (96%)	31 (4%)	40	16

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	104	GLN
1	E	42	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	129	MSE
1	D	111	ARG
1	E	61	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 38 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	78	GLN
1	E	150	GLN
1	H	116	GLN
1	E	97	ASN
1	E	151	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	118/130 (90%)	0.34	9 (7%) 17 19	16, 28, 80, 92	0
1	B	122/130 (93%)	-0.05	3 (2%) 61 65	16, 27, 53, 92	2 (1%)
1	C	120/130 (92%)	0.06	3 (2%) 61 65	15, 25, 51, 71	0
1	D	115/130 (88%)	0.21	3 (2%) 59 64	15, 29, 59, 85	1 (0%)
1	E	121/130 (93%)	-0.16	3 (2%) 61 65	17, 29, 50, 86	1 (0%)
1	F	119/130 (91%)	0.25	10 (8%) 14 15	16, 28, 69, 90	1 (0%)
1	G	119/130 (91%)	0.06	3 (2%) 61 65	14, 26, 70, 80	0
1	H	115/130 (88%)	0.40	9 (7%) 16 18	15, 38, 82, 118	0
All	All	949/1040 (91%)	0.14	43 (4%) 37 41	14, 29, 69, 118	5 (0%)

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	99	VAL	8.7
1	G	39	SER	6.8
1	H	99	VAL	6.7
1	A	42	GLN	5.7
1	A	41	ALA	5.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.