



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 30, 2016 – 11:30 AM EST

PDB ID : 5L3B
Title : Human LSD1/CoREST: LSD1 D556G mutation
Authors : Pilotto, S.; Speranzini, V.; Marabelli, C.
Deposited on : 2016-04-06
Resolution : 3.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028320
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028320


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2060 (3.40-3.20)
Clashscore	102246	1058 (3.38-3.22)
Ramachandran outliers	100387	1038 (3.38-3.22)
Sidechain outliers	100360	1037 (3.38-3.22)
RSRZ outliers	91569	2070 (3.40-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	852	 68% 10% 22%
2	B	482	 23% 72%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6342 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysine-specific histone demethylase 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	666	Total	C	N	O	S	0	0	0
			5213	3322	906	965	20			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	556	GLY	ASP	engineered mutation	UNP O60341

- Molecule 2 is a protein called REST corepressor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	133	Total	C	N	O	S	0	0	0
			1076	676	194	203	3			

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lysine-specific histone demethylase 1A



- Molecule 2: REST corepressor 1





4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	119.09Å 179.12Å 234.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.30 45.79 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (50.00-3.30) 99.5 (45.79-3.30)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 3.32Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.195 , 0.208 0.202 , 0.216	Depositor DCC
R_{free} test set	740 reflections (1.99%)	DCC
Wilson B-factor (Å ²)	81.5	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 53.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6342	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/5327	0.66	2/7226 (0.0%)
2	B	0.45	0/1091	0.69	0/1471
All	All	0.44	0/6418	0.67	2/8697 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	820	ARG	NE-CZ-NH1	5.44	123.02	120.30
1	A	815	LEU	CA-CB-CG	5.41	127.74	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5213	0	5251	39	0
2	B	1076	0	1091	9	0
3	A	53	0	31	3	0
All	All	6342	0	6373	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:ILE:HD11	1:A:248:LEU:HD11	1.71	0.71
1:A:670:PHE:CZ	1:A:740:VAL:HG12	2.38	0.59
1:A:801:GLU:HG3	1:A:809:ALA:HA	1.83	0.59
1:A:566:THR:HG21	1:A:697:LEU:HD13	1.86	0.57
1:A:631:LYS:HE2	1:A:651:VAL:O	2.06	0.56
1:A:353:LEU:HB3	1:A:565:LEU:HD23	1.89	0.55
1:A:392:LEU:HD23	1:A:398:PHE:CD2	2.41	0.55
1:A:761:TYR:CD1	1:A:809:ALA:HB1	2.40	0.55
1:A:801:GLU:O	1:A:802:HIS:ND1	2.40	0.54
1:A:670:PHE:CE1	1:A:740:VAL:HG12	2.44	0.52
1:A:331:ALA:HA	3:A:901:FAD:N5	2.25	0.52
1:A:564:HIS:C	1:A:565:LEU:HD12	2.32	0.50
1:A:442:LYS:HE3	2:B:355:THR:HG21	1.94	0.49
1:A:538:PHE:CE1	1:A:706:LEU:HD23	2.48	0.49
1:A:694:PHE:HA	1:A:704:LEU:O	2.12	0.48
1:A:435:VAL:HG13	2:B:349:ILE:HD12	1.95	0.48
1:A:760:SER:HB2	3:A:901:FAD:HM83	1.96	0.47
1:A:366:ASN:OD1	1:A:367:GLY:N	2.47	0.47
1:A:756:TRP:O	1:A:757:ALA:HB2	2.15	0.46
1:A:340:ASN:OD1	1:A:342:MET:N	2.43	0.45
1:A:366:ASN:OD1	1:A:368:GLN:N	2.49	0.45
2:B:368:GLU:OE2	2:B:371:ARG:NH1	2.48	0.45
2:B:327:ASN:ND2	2:B:330:ALA:HB2	2.32	0.45
1:A:801:GLU:O	1:A:802:HIS:CG	2.71	0.44
2:B:395:ILE:HG22	2:B:433:VAL:HG12	1.99	0.43
1:A:659:LEU:HD11	3:A:901:FAD:HM73	2.00	0.43
1:A:418:LEU:CD1	2:B:324:VAL:HG21	2.48	0.43
1:A:214:ARG:O	1:A:218:LEU:HD12	2.19	0.42
1:A:671:TRP:O	1:A:673:PRO:HD3	2.20	0.42
1:A:700:ALA:HB1	1:A:701:PRO:HD2	2.02	0.42
1:A:804:ILE:O	1:A:804:ILE:HG23	2.20	0.41
1:A:384:ARG:NH1	2:B:312:LYS:O	2.48	0.41
1:A:534:ALA:HB2	1:A:688:ARG:HG3	2.02	0.41
1:A:632:GLN:CD	1:A:758:ARG:HE	2.24	0.41
1:A:666:PHE:O	1:A:701:PRO:HG2	2.21	0.41
1:A:695:TRP:CE3	1:A:697:LEU:HD11	2.56	0.41
1:A:668:ARG:HD3	1:A:741:PRO:HG3	2.03	0.41
1:A:374:LYS:NZ	1:A:525:ASP:OD1	2.54	0.41
1:A:473:ASP:OD1	1:A:475:THR:N	2.53	0.40
1:A:418:LEU:HD12	2:B:324:VAL:HG21	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:456:LYS:HA	2:B:370:TYR:CD1	2.57	0.40
1:A:594:ARG:HD2	1:A:603:ILE:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	664/852 (78%)	634 (96%)	28 (4%)	2 (0%)	46	81
2	B	131/482 (27%)	119 (91%)	10 (8%)	2 (2%)	13	49
All	All	795/1334 (60%)	753 (95%)	38 (5%)	4 (0%)	34	71

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	757	ALA
2	B	326	ALA
2	B	439	ALA
1	A	468	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	565/698 (81%)	533 (94%)	32 (6%)	25	65
2	B	117/395 (30%)	105 (90%)	12 (10%)	9	34
All	All	682/1093 (62%)	638 (94%)	44 (6%)	21	59

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	191	GLN
1	A	243	ASN
1	A	246	THR
1	A	269	ARG
1	A	273	LEU
1	A	328	ASP
1	A	368	GLN
1	A	437	THR
1	A	448	MET
1	A	458	LEU
1	A	469	LYS
1	A	473	ASP
1	A	479	LEU
1	A	485	ARG
1	A	492	LYS
1	A	508	LEU
1	A	511	LEU
1	A	523	SER
1	A	525	ASP
1	A	526	ARG
1	A	571	TYR
1	A	591	ARG
1	A	594	ARG
1	A	610	THR
1	A	611	SER
1	A	668	ARG
1	A	677	LEU
1	A	684	THR
1	A	696	ASN
1	A	719	SER
1	A	750	ARG
1	A	776	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	314	MET
2	B	316	LEU
2	B	317	SER
2	B	332	THR
2	B	337	GLN
2	B	344	SER
2	B	349	ILE
2	B	371	ARG
2	B	384	THR
2	B	385	THR
2	B	425	ARG
2	B	426	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	243	ASN
1	A	828	GLN
2	B	327	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FAD	A	901	-	52,58,58	1.32	8 (15%)	52,89,89	2.27	14 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	901	-	-	0/30/50/50	0/6/6/6

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	901	FAD	C6-C5X	-2.01	1.38	1.41
3	A	901	FAD	C2A-N3A	2.02	1.35	1.32
3	A	901	FAD	C10-N10	2.64	1.42	1.39
3	A	901	FAD	C5A-C4A	2.95	1.47	1.40
3	A	901	FAD	C8-C7	3.03	1.49	1.41
3	A	901	FAD	C9A-N10	3.24	1.43	1.38
3	A	901	FAD	C4X-C10	3.58	1.47	1.40
3	A	901	FAD	C9A-C5X	3.71	1.50	1.42

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	901	FAD	N3A-C2A-N1A	-6.47	123.79	128.87
3	A	901	FAD	N3-C2-N1	-4.10	120.79	127.69
3	A	901	FAD	C4-C4X-C10	-4.04	117.36	119.94
3	A	901	FAD	C4X-C10-N10	-3.38	118.06	120.52
3	A	901	FAD	C1B-N9A-C4A	-3.12	123.33	126.81
3	A	901	FAD	C4X-C4-N3	-2.92	119.71	123.52
3	A	901	FAD	C4B-O4B-C1B	-2.32	107.18	109.64
3	A	901	FAD	C6-C5X-N5	-2.22	116.15	118.92
3	A	901	FAD	O4'-C4'-C3'	2.39	115.12	108.96
3	A	901	FAD	C4X-N5-C5X	2.58	119.77	116.72
3	A	901	FAD	N6A-C6A-N1A	2.91	123.39	118.52
3	A	901	FAD	C6-C5X-C9A	3.10	122.53	119.11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	901	FAD	C1'-N10-C9A	4.49	124.04	118.83
3	A	901	FAD	C4-N3-C2	8.57	122.31	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	901	FAD	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	666/852 (78%)	0.05	4 (0%) 90 88	47, 79, 113, 143	0
2	B	133/482 (27%)	0.35	3 (2%) 64 57	71, 107, 130, 148	0
All	All	799/1334 (59%)	0.10	7 (0%) 85 82	47, 83, 120, 148	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	242	TYR	2.5
1	A	171	PRO	2.5
2	B	374	GLU	2.4
1	A	509	GLN	2.2
2	B	312	LYS	2.2
1	A	270	ILE	2.2
2	B	365	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	FAD	A	901	53/53	0.97	0.22	-0.22	49,58,68,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.