



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:15 AM GMT

PDB ID : 3L75  
Title : Cytochrome BC1 complex from chicken with fenamidone bound  
Authors : Huang, L.; Berry, E.A.  
Deposited on : 2009-12-28  
Resolution : 2.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	446	<div> <div>2%</div> <div>65%</div> <div>31%</div> <div>..</div> </div>
1	N	446	<div> <div>5%</div> <div>67%</div> <div>29%</div> <div>..</div> </div>
2	B	441	<div> <div>5%</div> <div>55%</div> <div>38%</div> <div>• 5%</div> </div>
2	O	441	<div> <div>4%</div> <div>55%</div> <div>39%</div> <div>• •</div> </div>
3	C	380	<div> <div>80%</div> <div>20%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	P	380	
4	D	241	
4	Q	241	
5	E	196	
5	R	196	
6	F	110	
6	S	110	
7	G	81	
7	T	81	
8	H	77	
8	U	77	
9	I	47	
9	V	47	
10	J	61	
10	W	61	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
11	PEE	C	2005	-	-	-	X
11	PEE	C	2007	-	-	-	X
11	PEE	N	3008	-	X	-	-
11	PEE	P	3005	-	-	-	X
11	PEE	P	3007	-	-	-	X
12	UNL	C	2046	-	-	-	X
12	UNL	C	2048	-	-	-	X
12	UNL	P	3046	-	-	-	X
15	UQ	C	2002	-	-	-	X
15	UQ	P	3002	-	-	-	X
16	AZI	C	2011	-	-	-	X
16	AZI	P	3011	-	-	-	X

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
17	BOG	D	2091	-	-	-	X
17	BOG	P	2010	-	-	-	X
19	CDL	G	2004	-	-	-	X
20	FES	R	501	-	-	X	-

## 2 Entry composition

There are 21 unique types of molecules in this entry. The entry contains 32691 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MITOCHONDRIAL UBIQUINOL-CYTOCHROME-C REDUCTASE COMPLEX CORE PROTEIN I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	443	Total	C	N	O	S	0	0	1
			3440	2155	606	658	21			
1	N	442	Total	C	N	O	S	0	0	0
			3437	2154	605	657	21			

- Molecule 2 is a protein called MITOCHONDRIAL UBIQUINOL-CYTOCHROME-C REDUCTASE COMPLEX CORE PROTEIN 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	421	Total	C	N	O	S	0	0	0
			3141	1974	545	613	9			
2	O	422	Total	C	N	O	S	0	0	0
			3147	1977	546	614	10			

- Molecule 3 is a protein called CYTOCHROME B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	380	Total	C	N	O	S	0	0	0
			3017	2022	478	505	12			
3	P	379	Total	C	N	O	S	0	0	0
			3012	2019	477	504	12			

- Molecule 4 is a protein called MITOCHONDRIAL CYTOCHROME C1, HEME PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	241	Total	C	N	O	S	0	0	0
			1898	1212	327	347	12			
4	Q	241	Total	C	N	O	S	0	0	0
			1898	1212	327	347	12			

- Molecule 5 is a protein called CYTOCHROME B-C1 COMPLEX SUBUNIT 5, RIESKE IRONSULFUR PROTEIN, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	196	Total	C	N	O	S	0	0	0
			1513	952	263	292	6			
5	R	196	Total	C	N	O	S	0	0	0
			1512	952	262	292	6			

- Molecule 6 is a protein called MITOCHONDRIAL UBIQUINOL-CYTOCHROME C REDUCTASE 14 KDA PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	101	Total	C	N	O	S	0	0	0
			891	570	159	159	3			
6	S	101	Total	C	N	O	S	0	0	0
			891	570	159	159	3			

- Molecule 7 is a protein called MITOCHONDRIAL UBIQUINOL-CYTOCHROME C REDUCTASE UBIQUINONE-BINDING PROTEIN QP-C.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	G	81	Total	C	N	O	0	0	0
			676	439	120	117			
7	T	78	Total	C	N	O	0	0	0
			654	428	116	110			

- Molecule 8 is a protein called MITOCHONDRIAL UBIQUINOL-CYTOCHROME C REDUCTASE 11 KDA PROTEIN, COMPLEX III SUBUNIT VIII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	70	Total	C	N	O	S	0	0	0
			574	350	105	114	5			
8	U	67	Total	C	N	O	S	0	0	0
			553	338	103	107	5			


- Molecule 9 is a protein called CYTOCHROME B-C1 COMPLEX SUBUNIT RIESKE, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	46	Total	C	N	O	S	0	0	0
			288	172	58	56	2			
9	V	44	Total	C	N	O	S	0	0	1
			278	167	56	53	2			

- Molecule 10 is a protein called MITOCHONDRIAL UBIQUINOL-CYTOCHROME C REDUCTASE 7.2 KDA PROTEIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
10	J	61	Total	C	N	O	0	0	0
			497	321	87	89			
10	W	60	Total	C	N	O	0	0	1
			479	311	86	82			

- Molecule 11 is 1,2-DIOLEOYL-SN-GLYCERO-3-PHOSPHOETHANOLAMINE (three-letter code: PEE) (formula: C<sub>41</sub>H<sub>83</sub>NO<sub>8</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
11	A	1	Total	C	O	P		0	0
			21	12	8	1			
11	C	1	Total	C	N	O	P	0	0
			50	40	1	8	1		
11	C	1	Total	C	N	O	P	0	0
			48	38	1	8	1		
11	N	1	Total	O	P			0	0
			5	4	1				
11	P	1	Total	C	N	O	P	0	0
			50	40	1	8	1		
11	P	1	Total	C	N	O	P	0	0
			48	38	1	8	1		

- Molecule 12 is UNKNOWN LIGAND (three-letter code: UNL) (formula: ).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	P	5	Total	O	0	0
			7	7		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	R	1	Total O 1 1	0	0
12	A	1	Total O 1 1	0	0
12	D	1	Total O 2 2	0	0
12	C	3	Total O 5 5	0	0

- Molecule 13 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
13	C	1	Total C Fe N O 43 34 1 4 4	0	0
13	C	1	Total C Fe N O 43 34 1 4 4	0	0
13	P	1	Total C Fe N O 43 34 1 4 4	0	0
13	P	1	Total C Fe N O 43 34 1 4 4	0	0

- Molecule 14 is (5S)-5-METHYL-2-(METHYLSULFANYL)-5-PHENYL-3-(PHENYLAMINO)-3,5-DIHYDRO-4H-IMIDAZOL-4-ONE (three-letter code: FNM) (formula:  $C_{17}H_{17}N_3OS$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
14	C	1	Total	C	N	O	S	0	0
			22	17	3	1	1		
14	P	1	Total	C	N	O	S	0	0
			22	17	3	1	1		

- Molecule 15 is COENZYME Q10, (2Z,6E,10Z,14E,18E,22E,26Z)-ISOMER (three-letter code: UQ) (formula: C<sub>59</sub>H<sub>90</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
15	C	1	Total	C	O	0	0
			19	15	4		

Continued on next page...

*Continued from previous page...*


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
15	P	1	Total	C	O	0	0
			19	15	4		

- Molecule 16 is AZIDE ION (three-letter code: AZI) (formula: N<sub>3</sub>).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	C	1	Total	N	0	0
			3	3		
16	P	1	Total	N	0	0
			3	3		

- Molecule 17 is SUGAR (B-OCTYLGLUCOSIDE) (three-letter code: BOG) (formula: C<sub>14</sub>H<sub>28</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
17	C	1	Total	C	O	0	0
			12	10	2		
17	D	1	Total	C	O	0	0
			20	14	6		
17	D	1	Total	C	O	0	0
			20	14	6		
17	P	1	Total	C	O	0	0
			19	13	6		
17	Q	1	Total	C	O	0	0
			20	14	6		
17	Q	1	Total	C	O	0	0
			20	14	6		

- Molecule 18 is HEME C (three-letter code: HEC) (formula: C<sub>34</sub>H<sub>34</sub>FeN<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
18	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
18	Q	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 19 is CARDIOLIPIN (three-letter code: CDL) (formula:  $C_{81}H_{156}O_{17}P_2$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
19	D	1	Total	C	O	P	0	0
			42	23	17	2		
19	G	1	Total	C	O	P	0	0
			40	21	17	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
19	Q	1	Total	C	O	P	0	0
			42	23	17	2		
19	T	1	Total	C	O	P	0	0
			40	21	17	2		

- Molecule 20 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe<sub>2</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
20	E	1	Total	Fe	S	0	0
			4	2	2		
20	R	1	Total	Fe	S	0	0
			4	2	2		


- Molecule 21 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
21	A	2	Total	O	0	0
			2	2		
21	C	9	Total	O	0	0
			9	9		
21	E	3	Total	O	0	0
			3	3		
21	P	10	Total	O	0	0
			10	10		
21	R	4	Total	O	0	0
			4	4		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: MITOCHONDRIAL UBIQUINOL-CYTOCHROME-C REDUCTASE COMPLEX CORE PROTEIN I




#### • Molecule 1: MITOCHONDRIAL UBIQUINOL-CYTOCHROME-C REDUCTASE COMPLEX CORE PROTEIN I




• Molecule 2: MITOCHONDRIAL UBIQUINOL-CYTOCHROME-C REDUCTASE COMPLEX CORE PROTEIN 2




• Molecule 2: MITOCHONDRIAL UBIQUINOL-CYTOCHROME-C REDUCTASE COMPLEX CORE PROTEIN 2





• Molecule 3: CYTOCHROME B









• Molecule 3: CYTOCHROME B



• Molecule 4: MITOCHONDRIAL CYTOCHROME C1, HEME PROTEIN




• Molecule 4: MITOCHONDRIAL CYTOCHROME C1, HEME PROTEIN




• Molecule 5: CYTOCHROME B-C1 COMPLEX SUBUNIT 5, RIESKE IRONSULFUR PROTEIN, MITOCHONDRIAL







- Molecule 5: CYTOCHROME B-C1 COMPLEX SUBUNIT 5, RIESKE IRONSULFUR PROTEIN, MITOCHONDRIAL



- Molecule 6: MITOCHONDRIAL UBIQUINOL-CYTOCHROME C REDUCTASE 14 KDA PROTEIN



- Molecule 6: MITOCHONDRIAL UBIQUINOL-CYTOCHROME C REDUCTASE 14 KDA PROTEIN



- Molecule 7: MITOCHONDRIAL UBIQUINOL-CYTOCHROME C REDUCTASE UBIQUINONE-BINDING PROTEIN QP-C




• Molecule 7: MITOCHONDRIAL UBIQUINOL-CYTOCHROME C REDUCTASE UBIQUINONE-BINDING PROTEIN QP-C

Chain T: 




• Molecule 8: MITOCHONDRIAL UBIQUINOL-CYTOCHROME C REDUCTASE 11 KDA PROTEIN, COMPLEX III SUBUNIT VIII

Chain H: 





• Molecule 8: MITOCHONDRIAL UBIQUINOL-CYTOCHROME C REDUCTASE 11 KDA PROTEIN, COMPLEX III SUBUNIT VIII

Chain U: 



• Molecule 9: CYTOCHROME B-C1 COMPLEX SUBUNIT RIESKE, MITOCHONDRIAL

Chain I: 




• Molecule 9: CYTOCHROME B-C1 COMPLEX SUBUNIT RIESKE, MITOCHONDRIAL

Chain V: 




• Molecule 10: MITOCHONDRIAL UBIQUINOL-CYTOCHROME C REDUCTASE 7.2 KDA PROTEIN

Chain J: 



● Molecule 10: MITOCHONDRIAL UBIQUINOL-CYTOCHROME C REDUCTASE 7.2 KDA PROTEIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	172.28Å 182.14Å 241.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.78 – 2.79 58.78 – 2.79	Depositor EDS
% Data completeness (in resolution range)	97.6 (58.78-2.79) 97.3 (58.78-2.79)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.03 (at 2.77Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.248 , 0.275 0.243 , 0.269	Depositor DCC
$R_{free}$ test set	3610 reflections (2.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	64.6	Xtriage
Anisotropy	0.413	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 53.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 183994 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	32691	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.81% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: AZI, CDL, UQ, FES, HEC, PEE, FNM, UNL, HEM, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/3511	0.63	0/4757
1	N	0.42	0/3508	0.63	0/4753
2	B	0.37	0/3196	0.60	0/4334
2	O	0.38	0/3202	0.62	1/4343 (0.0%)
3	C	0.52	0/3119	0.65	0/4270
3	P	0.46	0/3114	0.63	0/4263
4	D	0.47	0/1956	0.63	0/2658
4	Q	0.39	0/1956	0.60	0/2658
5	E	0.37	0/1547	0.59	0/2103
5	R	0.35	0/1545	0.57	0/2098
6	F	0.51	0/911	0.65	0/1219
6	S	0.42	0/911	0.61	0/1219
7	G	0.49	0/698	0.66	0/946
7	T	0.44	0/676	0.65	0/918
8	H	0.43	0/582	0.59	0/779
8	U	0.31	0/561	0.55	0/751
9	I	0.36	0/218	0.61	0/293
9	V	0.36	0/218	0.58	0/293
10	J	0.43	0/508	0.59	0/682
10	W	0.42	0/490	0.57	0/660
All	All	0.43	0/32427	0.62	1/43997 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	O	227	ARG	N-CA-C	5.48	125.81	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3440	0	3353	135	0
1	N	3437	0	3349	137	0
2	B	3141	0	3142	187	0
2	O	3147	0	3146	178	0
3	C	3017	0	3063	69	0
3	P	3012	0	3058	60	0
4	D	1898	0	1846	48	0
4	Q	1898	0	1846	47	0
5	E	1513	0	1478	67	0
5	R	1512	0	1476	95	0
6	F	891	0	893	15	0
6	S	891	0	893	22	0
7	G	676	0	659	26	0
7	T	654	0	641	24	0
8	H	574	0	548	17	0
8	U	553	0	535	23	0
9	I	288	0	254	42	0
9	V	278	0	252	31	0
10	J	497	0	490	11	0
10	W	479	0	478	14	0
11	A	21	0	13	1	0
11	C	98	0	147	0	0
11	N	5	0	0	0	0
11	P	98	0	147	1	0
12	A	1	0	0	0	0
12	C	5	0	0	0	0
12	D	2	0	0	0	0
12	P	7	0	0	0	0
12	R	1	0	0	0	0
13	C	86	0	60	5	0
13	P	86	0	60	5	0
14	C	22	0	17	2	0
14	P	22	0	17	1	0
15	C	19	0	17	4	0
15	P	19	0	17	3	0
16	C	3	0	0	0	0
16	P	3	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
17	C	12	0	18	0	0
17	D	40	0	56	2	0
17	P	19	0	24	1	0
17	Q	40	0	56	1	0
18	D	43	0	30	3	0
18	Q	43	0	30	1	0
19	D	42	0	28	2	0
19	G	40	0	24	1	0
19	Q	42	0	28	3	0
19	T	40	0	24	1	0
20	E	4	0	0	1	0
20	R	4	0	0	2	0
21	A	2	0	0	0	0
21	C	9	0	0	1	0
21	E	3	0	0	0	0
21	P	10	0	0	1	0
21	R	4	0	0	0	0
All	All	32691	0	32213	1132	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 1132 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:O:353:THR:HG22	2:O:355:GLU:H	1.16	1.11
5:R:83:GLU:HG2	5:R:102:THR:HG22	1.33	1.07
2:B:353:THR:HG22	2:B:355:GLU:H	1.20	1.06
2:B:157:VAL:HG23	9:I:64:LEU:HD21	1.38	1.03
1:N:178:THR:HG22	1:N:180:ALA:H	1.22	1.02

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/446 (99%)	412 (93%)	23 (5%)	6 (1%)	14	42
1	N	440/446 (99%)	411 (93%)	22 (5%)	7 (2%)	12	38
2	B	419/441 (95%)	374 (89%)	35 (8%)	10 (2%)	7	25
2	O	420/441 (95%)	372 (89%)	41 (10%)	7 (2%)	11	36
3	C	378/380 (100%)	364 (96%)	14 (4%)	0	100	100
3	P	377/380 (99%)	360 (96%)	17 (4%)	0	100	100
4	D	239/241 (99%)	228 (95%)	11 (5%)	0	100	100
4	Q	239/241 (99%)	224 (94%)	15 (6%)	0	100	100
5	E	194/196 (99%)	170 (88%)	18 (9%)	6 (3%)	5	17
5	R	192/196 (98%)	156 (81%)	28 (15%)	8 (4%)	3	11
6	F	99/110 (90%)	93 (94%)	5 (5%)	1 (1%)	19	52
6	S	99/110 (90%)	91 (92%)	6 (6%)	2 (2%)	9	30
7	G	79/81 (98%)	69 (87%)	10 (13%)	0	100	100
7	T	76/81 (94%)	68 (90%)	8 (10%)	0	100	100
8	H	68/77 (88%)	62 (91%)	6 (9%)	0	100	100
8	U	65/77 (84%)	60 (92%)	4 (6%)	1 (2%)	13	40
9	I	29/47 (62%)	26 (90%)	2 (7%)	1 (3%)	5	16
9	V	29/47 (62%)	24 (83%)	5 (17%)	0	100	100
10	J	59/61 (97%)	55 (93%)	3 (5%)	1 (2%)	11	36
10	W	58/61 (95%)	54 (93%)	2 (3%)	2 (3%)	5	16
All	All	4000/4160 (96%)	3673 (92%)	275 (7%)	52 (1%)	15	44

5 of 52 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	282	ARG
2	B	26	ILE
2	B	226	ILE
2	B	227	ARG
2	B	389	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/368 (99%)	355 (97%)	10 (3%)	52	85
1	N	365/368 (99%)	351 (96%)	14 (4%)	40	74
2	B	332/347 (96%)	327 (98%)	5 (2%)	72	93
2	O	333/347 (96%)	327 (98%)	6 (2%)	66	91
3	C	328/329 (100%)	323 (98%)	5 (2%)	72	93
3	P	328/329 (100%)	323 (98%)	5 (2%)	72	93
4	D	200/200 (100%)	197 (98%)	3 (2%)	72	93
4	Q	200/200 (100%)	197 (98%)	3 (2%)	72	93
5	E	166/166 (100%)	165 (99%)	1 (1%)	90	98
5	R	165/166 (99%)	159 (96%)	6 (4%)	42	76
6	F	93/96 (97%)	91 (98%)	2 (2%)	60	89
6	S	93/96 (97%)	90 (97%)	3 (3%)	46	80
7	G	71/71 (100%)	71 (100%)	0	100	100
7	T	69/71 (97%)	68 (99%)	1 (1%)	74	94
8	H	65/71 (92%)	64 (98%)	1 (2%)	72	93
8	U	63/71 (89%)	62 (98%)	1 (2%)	70	93
9	I	23/26 (88%)	22 (96%)	1 (4%)	35	70
9	V	23/26 (88%)	23 (100%)	0	100	100
10	J	49/49 (100%)	47 (96%)	2 (4%)	37	72
10	W	47/49 (96%)	46 (98%)	1 (2%)	61	90
All	All	3378/3446 (98%)	3308 (98%)	70 (2%)	61	90

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	N	18	THR
1	N	281	ASP
6	S	14	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	N	49	ASN
1	N	90	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 85 such sidechains are listed below:

Mol	Chain	Res	Type
7	G	44	GLN
1	N	126	GLN
5	R	164	HIS
7	G	79	ASN
1	N	10	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 41 ligands modelled in this entry, 11 are unknown - leaving 30 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
11	PEE	A	2008	-	20,20,50	1.67	5 (25%)	21,25,55	0.64	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
14	FNM	C	2001	-	21,24,24	1.78	2 (9%)	21,34,34	1.93	2 (9%)
15	UQ	C	2002	-	19,19,63	2.46	11 (57%)	23,26,79	1.08	3 (13%)
11	PEE	C	2005	-	49,49,50	1.37	9 (18%)	50,54,55	0.90	5 (10%)
11	PEE	C	2007	-	47,47,50	1.18	6 (12%)	48,52,55	0.90	5 (10%)
16	AZI	C	2011	-	0,2,2	0.00	-	0,1,1	0.00	-
17	BOG	C	3010	-	10,11,20	0.84	1 (10%)	9,11,25	0.91	0
13	HEM	C	501	3	30,50,50	2.84	9 (30%)	24,82,82	2.27	7 (29%)
13	HEM	C	502	3	30,50,50	2.51	7 (23%)	24,82,82	2.25	6 (25%)
19	CDL	D	2003	-	41,41,99	1.12	1 (2%)	43,53,111	1.08	5 (11%)
17	BOG	D	2009	-	20,20,20	0.96	1 (5%)	25,25,25	0.95	2 (8%)
17	BOG	D	2091	-	20,20,20	1.10	2 (10%)	25,25,25	0.99	1 (4%)
18	HEC	D	501	4	24,50,50	1.63	3 (12%)	19,82,82	3.16	7 (36%)
20	FES	E	501	5	0,4,4	0.00	-	0,4,4	0.00	-
19	CDL	G	2004	-	39,39,99	1.20	3 (7%)	41,51,111	1.12	4 (9%)
11	PEE	N	3008	-	4,4,50	3.66	4 (100%)	6,6,55	0.53	0
17	BOG	P	2010	-	18,18,20	1.16	3 (16%)	22,22,25	0.57	0
14	FNM	P	3001	-	21,24,24	1.79	3 (14%)	21,34,34	1.79	2 (9%)
15	UQ	P	3002	-	19,19,63	2.31	10 (52%)	23,26,79	1.06	3 (13%)
11	PEE	P	3005	-	49,49,50	1.36	9 (18%)	50,54,55	0.90	5 (10%)
11	PEE	P	3007	-	47,47,50	1.24	6 (12%)	48,52,55	0.86	4 (8%)
16	AZI	P	3011	-	0,2,2	0.00	-	0,1,1	0.00	-
13	HEM	P	501	3	30,50,50	2.82	9 (30%)	24,82,82	2.32	8 (33%)
13	HEM	P	502	3	30,50,50	2.61	9 (30%)	24,82,82	2.26	7 (29%)
19	CDL	Q	3003	-	41,41,99	1.17	2 (4%)	43,53,111	1.08	4 (9%)
17	BOG	Q	3009	-	20,20,20	0.93	1 (5%)	25,25,25	0.87	2 (8%)
17	BOG	Q	3091	-	20,20,20	1.12	2 (10%)	25,25,25	0.91	2 (8%)
18	HEC	Q	501	4	24,50,50	2.14	2 (8%)	19,82,82	3.06	6 (31%)
20	FES	R	501	5	0,4,4	0.00	-	0,4,4	0.00	-
19	CDL	T	3004	-	39,39,99	1.20	3 (7%)	41,51,111	1.11	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	PEE	A	2008	-	-	0/24/24/54	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
14	FNM	C	2001	-	-	0/12/31/31	0/3/3/3
15	UQ	C	2002	-	-	0/11/35/87	0/1/1/1
11	PEE	C	2005	-	-	0/53/53/54	0/0/0/0
11	PEE	C	2007	-	-	0/51/51/54	0/0/0/0
16	AZI	C	2011	-	-	0/0/0/0	0/0/0/0
17	BOG	C	3010	-	-	0/8/9/31	0/0/0/1
13	HEM	C	501	3	-	0/10/54/54	0/0/8/8
13	HEM	C	502	3	-	0/10/54/54	0/0/8/8
19	CDL	D	2003	-	-	0/51/51/110	0/0/0/0
17	BOG	D	2009	-	-	0/11/31/31	0/1/1/1
17	BOG	D	2091	-	-	0/11/31/31	0/1/1/1
18	HEC	D	501	4	-	0/6/54/54	0/0/8/8
20	FES	E	501	5	-	0/0/4/4	0/1/1/1
19	CDL	G	2004	-	-	0/49/49/110	0/0/0/0
11	PEE	N	3008	-	-	0/0/0/54	0/0/0/0
17	BOG	P	2010	-	-	0/6/26/31	0/1/1/1
14	FNM	P	3001	-	-	0/12/31/31	0/3/3/3
15	UQ	P	3002	-	-	0/11/35/87	0/1/1/1
11	PEE	P	3005	-	-	0/53/53/54	0/0/0/0
11	PEE	P	3007	-	-	0/51/51/54	0/0/0/0
16	AZI	P	3011	-	-	0/0/0/0	0/0/0/0
13	HEM	P	501	3	-	0/10/54/54	0/0/8/8
13	HEM	P	502	3	-	0/10/54/54	0/0/8/8
19	CDL	Q	3003	-	-	0/51/51/110	0/0/0/0
17	BOG	Q	3009	-	-	0/11/31/31	0/1/1/1
17	BOG	Q	3091	-	-	0/11/31/31	0/1/1/1
18	HEC	Q	501	4	-	0/6/54/54	0/0/8/8
20	FES	R	501	5	-	0/0/4/4	0/1/1/1
19	CDL	T	3004	-	-	0/49/49/110	0/0/0/0

The worst 5 of 123 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
13	C	501	HEM	C3B-CAB	-7.52	1.37	1.51
13	P	501	HEM	C3B-C4B	-7.25	1.45	1.51
13	C	501	HEM	C3B-C4B	-7.12	1.45	1.51
13	P	501	HEM	C3B-CAB	-6.98	1.38	1.51
18	Q	501	HEC	C3B-C2B	-6.87	1.33	1.40

The worst 5 of 93 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
18	Q	501	HEC	CBB-CAB-C3B	-7.55	110.58	127.35
18	D	501	HEC	CBB-CAB-C3B	-7.29	111.15	127.35
18	D	501	HEC	CBC-CAC-C3C	-6.07	113.87	127.35
18	Q	501	HEC	CBC-CAC-C3C	-5.64	114.81	127.35
13	C	501	HEM	CAA-C2A-C1A	-3.39	123.33	127.01

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

22 monomers are involved in 40 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	A	2008	PEE	1	0
14	C	2001	FNM	2	0
15	C	2002	UQ	4	0
13	C	501	HEM	1	0
13	C	502	HEM	4	0
19	D	2003	CDL	2	0
17	D	2009	BOG	1	0
17	D	2091	BOG	1	0
18	D	501	HEC	3	0
20	E	501	FES	1	0
19	G	2004	CDL	1	0
17	P	2010	BOG	1	0
14	P	3001	FNM	1	0
15	P	3002	UQ	3	0
11	P	3007	PEE	1	0
13	P	501	HEM	1	0
13	P	502	HEM	4	0
19	Q	3003	CDL	3	0
17	Q	3091	BOG	1	0
18	Q	501	HEC	1	0
20	R	501	FES	2	0
19	T	3004	CDL	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	443/446 (99%)	0.17	9 (2%) 68 58	29, 61, 87, 104	0
1	N	442/446 (99%)	0.34	21 (4%) 34 23	39, 69, 95, 104	0
2	B	421/441 (95%)	0.43	24 (5%) 27 17	50, 78, 116, 141	0
2	O	422/441 (95%)	0.30	17 (4%) 42 30	37, 74, 106, 120	0
3	C	380/380 (100%)	0.20	1 (0%) 94 92	21, 39, 73, 109	0
3	P	379/380 (99%)	0.23	5 (1%) 79 71	30, 62, 84, 96	0
4	D	241/241 (100%)	0.04	3 (1%) 81 73	30, 42, 78, 94	0
4	Q	241/241 (100%)	0.28	10 (4%) 41 29	50, 69, 99, 119	0
5	E	196/196 (100%)	1.18	56 (28%) 1 0	38, 88, 119, 125	0
5	R	196/196 (100%)	2.54	92 (46%) 0 0	42, 112, 159, 163	127 (64%)
6	F	101/110 (91%)	-0.11	0 100 100	29, 42, 60, 96	0
6	S	101/110 (91%)	0.29	4 (3%) 42 30	55, 68, 111, 135	0
7	G	81/81 (100%)	0.33	2 (2%) 61 48	32, 52, 93, 108	0
7	T	78/81 (96%)	0.82	10 (12%) 5 2	48, 82, 144, 161	0
8	H	70/77 (90%)	0.20	2 (2%) 55 43	39, 59, 84, 123	0
8	U	67/77 (87%)	1.14	13 (19%) 1 1	96, 115, 133, 140	0
9	I	31/47 (65%)	2.15	15 (48%) 0 0	81, 105, 125, 126	0
9	V	31/47 (65%)	2.10	13 (41%) 0 0	69, 109, 142, 145	0
10	J	61/61 (100%)	0.44	5 (8%) 14 7	44, 60, 92, 130	0
10	W	60/61 (98%)	1.06	5 (8%) 14 7	56, 77, 99, 111	0
All	All	4042/4160 (97%)	0.48	307 (7%) 17 9	21, 65, 116, 163	127 (3%)

The worst 5 of 307 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	R	193	VAL	14.2
5	R	195	VAL	13.9
5	R	133	VAL	13.8
10	W	62	SER	11.7
10	W	63	GLU	11.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
12	UNL	P	3046	2/-	0.83	0.72	32.14	85,85,85,85	0
16	AZI	P	3011	3/3	0.91	0.50	19.13	74,74,76,77	0
12	UNL	C	2048	2/-	0.89	0.41	18.33	62,62,62,63	0
12	UNL	C	2046	2/-	0.76	0.51	14.42	87,87,87,91	0
17	BOG	D	2091	20/20	0.24	0.60	13.76	177,189,190,190	0
16	AZI	C	2011	3/3	0.83	0.50	11.18	68,68,69,71	0
17	BOG	P	2010	19/20	0.59	0.47	9.94	102,183,184,184	0
11	PEE	P	3007	48/51	0.90	0.36	7.50	76,90,105,106	0
15	UQ	C	2002	19/63	0.83	0.33	4.74	85,87,89,89	0
15	UQ	P	3002	19/63	0.84	0.39	4.70	138,141,143,143	0
11	PEE	C	2005	50/51	0.83	0.32	2.77	83,93,101,104	0
19	CDL	G	2004	40/100	0.91	0.22	2.59	60,70,85,86	0
11	PEE	C	2007	48/51	0.95	0.23	2.38	41,52,81,82	0
11	PEE	P	3005	50/51	0.83	0.31	2.17	74,100,105,105	0
12	UNL	A	3015	1/-	0.92	0.21	1.92	48,48,48,48	0
12	UNL	P	3048	2/-	0.61	0.25	1.83	92,92,92,95	0
19	CDL	Q	3003	42/100	0.84	0.26	1.62	119,124,137,138	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
11	PEE	A	2008	21/51	0.88	0.24	1.23	95,113,116,118	0
19	CDL	D	2003	42/100	0.87	0.21	0.97	81,93,97,98	0
14	FNM	P	3001	22/22	0.96	0.22	0.83	49,53,59,62	0
17	BOG	D	2009	20/20	0.90	0.20	0.77	56,63,65,67	0
17	BOG	C	3010	12/20	0.86	0.35	0.65	94,96,97,98	0
19	CDL	T	3004	40/100	0.87	0.21	0.63	94,97,106,107	0
18	HEC	D	501	43/43	0.98	0.18	0.17	23,30,35,37	0
17	BOG	Q	3009	20/20	0.90	0.19	0.14	71,79,81,82	0
13	HEM	P	501	43/43	0.98	0.21	0.02	40,43,54,58	0
13	HEM	C	502	43/43	0.99	0.19	-0.00	23,26,33,39	0
13	HEM	P	502	43/43	0.98	0.17	-0.11	34,40,55,62	0
13	HEM	C	501	43/43	0.99	0.22	-0.11	23,31,38,43	0
14	FNM	C	2001	22/22	0.98	0.18	-0.22	38,40,42,44	0
18	HEC	Q	501	43/43	0.96	0.18	-0.26	47,55,64,66	0
20	FES	R	501	4/4	0.96	0.17	-1.30	106,106,106,107	4
20	FES	E	501	4/4	0.98	0.11	-1.49	83,84,85,86	0
12	UNL	P	3013	1/-	0.92	0.30	-	56,56,56,56	0
12	UNL	D	2012	2/-	0.77	0.32	-	59,59,59,60	0
12	UNL	P	3014	1/-	0.85	0.31	-	55,55,55,55	0
17	BOG	Q	3091	20/20	0.28	0.64	-	178,186,187,187	0
12	UNL	C	2047	1/-	0.96	0.35	-	34,34,34,34	0
12	UNL	P	3047	1/-	0.76	0.28	-	47,47,47,47	0
12	UNL	R	3012	1/-	0.94	0.19	-	38,38,38,38	0
11	PEE	N	3008	5/51	0.95	0.15	-	88,88,89,89	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.