



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 1, 2016 – 10:19 AM EST

PDB ID : 5LAL
Title : Structure of Arabidopsis dirigent protein AtDIR6
Authors : Gasper, R.; Kolesinski, P.; Terlecka, B.; Effenberger, I.; Schaller, A.; Hofmann, E.
Deposited on : 2016-06-14
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028320
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028320


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1199 (1.40-1.40)
Clashscore	102246	1295 (1.40-1.40)
Ramachandran outliers	100387	1259 (1.40-1.40)
Sidechain outliers	100360	1258 (1.40-1.40)
RSRZ outliers	91569	1198 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	158	<div> <div>8%</div> <div>86%</div> <div>9%</div> <div>..</div> </div>
1	B	158	<div> <div>8%</div> <div>87%</div> <div>10%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	208	-	-	-	X
4	MAN	B	204	-	-	-	X
4	MAN	B	211	-	-	-	X
7	EDO	A	210	-	-	-	X
7	EDO	A	213	-	-	-	X
7	EDO	A	214	-	-	-	X
7	EDO	B	215	-	-	-	X
7	EDO	B	217	-	-	-	X
8	GLY	A	215	-	-	-	X
8	GLY	A	216	-	-	-	X
8	GLY	B	218	-	-	-	X
8	GLY	B	220	-	-	-	X

2 Entry composition [i](#)


There are 9 unique types of molecules in this entry. The entry contains 5863 atoms, of which 2745 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dirigent protein 6.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	153	Total	C	H	N	O	S	0	13	0
			2495	851	1200	202	225	17			
1	B	155	Total	C	H	N	O	S	0	12	0
			2538	859	1229	204	230	16			

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			26	8	12	1	5		
2	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
2	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
2	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	H	N	O	
			26	8	12	1	5	
2	B	1	Total	C	H	N	O	
			27	8	13	1	5	
2	B	1	Total	C	H	N	O	
			25	8	11	1	5	
2	B	1	Total	C	H	N	O	
			26	8	12	1	5	

- Molecule 3 is BETA-D-MANNOSE (three-letter code: BMA) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O		
			19	6	8	5		
3	B	1	Total	C	H	O		
			20	6	9	5		
3	B	1	Total	C	H	O		
			18	6	7	5		

- Molecule 4 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			22	6	11	5		
4	A	1	Total	C	H	O	0	0
			22	6	11	5		
4	B	1	Total	C	H	O	0	0
			22	6	11	5		
4	B	1	Total	C	H	O	0	0
			22	6	11	5		
4	B	1	Total	C	H	O	0	0
			22	6	11	5		
4	B	1	Total	C	H	O	0	0
			22	6	11	5		

- Molecule 5 is BETA-D-XYLOPYRANOSE (three-letter code: XYP) (formula: C₅H₁₀O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			18	5	9	4		
5	B	1	Total	C	H	O	0	0
			18	5	9	4		
5	B	1	Total	C	H	O	0	0
			18	5	9	4		

- Molecule 6 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula: C₆H₁₂O₅).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			21	6	11	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	H	O	0	0
			21	6	11	4		
6	B	1	Total	C	H	O	0	0
			21	6	11	4		

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			10	2	6	2		
7	A	1	Total	C	H	O	0	0
			10	2	6	2		
7	A	1	Total	C	H	O	0	0
			10	2	6	2		
7	A	1	Total	C	H	O	0	0
			10	2	6	2		
7	B	1	Total	C	H	O	0	0
			10	2	6	2		
7	B	1	Total	C	H	O	0	0
			10	2	6	2		
7	B	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 8 is GLYCINE (three-letter code: GLY) (formula: $C_2H_5NO_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	A	1	Total	C	H	N	O	0	0
			7	2	2	1	2		
8	A	1	Total	C	H	N	O	0	0
			10	2	5	1	2		
8	B	1	Total	C	H	N	O	0	0
			7	2	2	1	2		
8	B	1	Total	C	H	N	O	0	0
			10	2	5	1	2		
8	B	1	Total	C	H	N	O	0	0
			10	2	5	1	2		



- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	96	Total	O	0	0
			96	96		
9	B	93	Total	O	0	0
			93	93		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dirigent protein 6



• Molecule 1: Dirigent protein 6



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	101.09 Å 101.09 Å 90.24 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.11 – 1.40 40.11 – 1.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (40.11-1.40) 99.9 (40.11-1.40)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 1.40 Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.169 , 0.186 0.169 , 0.187	Depositor DCC
R_{free} test set	3377 reflections (4.99%)	DCC
Wilson B-factor (Å ²)	20.5	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.056 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5863	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: XYP, BMA, NAG, EDO, FUC, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/1370	0.76	0/1837
1	B	0.50	0/1375	0.75	1/1845 (0.1%)
All	All	0.50	0/2745	0.76	1/3682 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	43	PHE	CB-CA-C	-6.30	97.80	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1295	1200	1293	17	1
1	B	1309	1229	1301	12	0
2	A	56	51	48	2	0
2	B	56	48	46	0	0
3	A	11	8	7	0	0
3	B	22	16	14	5	0
4	A	22	22	20	0	0
4	B	44	44	40	5	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	9	9	8	0	0
5	B	18	18	16	0	0
6	A	10	11	10	0	0
6	B	20	22	20	0	0
7	A	20	30	30	0	0
7	B	12	18	18	0	0
8	A	10	7	4	0	0
8	B	15	12	6	2	0
9	A	96	0	0	2	0
9	B	93	0	0	2	0
All	All	3118	2745	2881	32	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:203:BMA:O3	4:B:207:MAN:C1	1.66	1.43
3:B:203:BMA:C3	4:B:207:MAN:C1	2.38	1.01
1:A:78[B]:LYS:NZ	9:A:301:HOH:O	2.06	0.89
1:A:42:HIS:NE2	1:A:180[B]:ASP:OD2	2.12	0.82
1:A:159[B]:ARG:NH1	1:A:185:GLU:OE2	2.12	0.82
1:B:42:HIS:NE2	1:B:180[B]:ASP:OD2	2.16	0.79
3:B:203:BMA:H3	4:B:207:MAN:C1	2.16	0.75
1:B:180[A]:ASP:OD2	9:B:301:HOH:O	2.07	0.73
1:B:41:LYS:HE3	1:B:43:PHE:HE1	1.56	0.70
1:A:36:GLN:HG2	1:A:37:LYS:H	1.57	0.69
1:A:42:HIS:HE2	1:A:180[B]:ASP:CG	1.97	0.68
3:B:203:BMA:O3	4:B:207:MAN:C2	2.42	0.67
1:B:41:LYS:HE3	1:B:43:PHE:CE1	2.36	0.59
2:A:208:NAG:H81	9:A:391:HOH:O	2.07	0.54
1:A:36:GLN:HB3	1:B:33:THR:OG1	2.10	0.52
1:A:36:GLN:CG	1:A:37:LYS:H	2.22	0.50
1:A:35:ASP:HB2	1:B:187:TYR:CE1	2.49	0.48
1:A:123:ASN:HD22	2:A:208:NAG:H83	1.80	0.47
1:B:137:ASP:OD2	8:B:218:GLY:N	2.48	0.47
8:B:219:GLY:N	9:B:303:HOH:O	2.47	0.46
1:A:35:ASP:O	1:A:36:GLN:HB2	2.16	0.46
1:A:36:GLN:HG2	1:A:37:LYS:N	2.29	0.45
3:B:203:BMA:O3	4:B:207:MAN:C5	2.63	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:35:ASP:N	1:A:184:TYR:HB3	2.32	0.45
1:A:35:ASP:OD1	1:A:35:ASP:N	2.50	0.45
1:A:38:LYS:HD2	1:A:39:PRO:HD2	2.00	0.42
1:A:40[B]:CYS:HB3	1:B:35:ASP:HB2	2.02	0.41
1:B:113:ASN:OD1	1:B:138:LEU:HA	2.20	0.41
1:B:113:ASN:HA	1:B:139:MET:HG2	2.01	0.41
1:A:35:ASP:HB2	1:B:187:TYR:HE1	1.85	0.41
1:B:69:PRO:HD2	1:B:76:PHE:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:154:ASP:OD2	4:B:211:MAN:O4[8_444]	1.91	0.29

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	164/158 (104%)	159 (97%)	5 (3%)	0	100	100
1	B	165/158 (104%)	161 (98%)	4 (2%)	0	100	100
All	All	329/316 (104%)	320 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	143/135 (106%)	139 (97%)	4 (3%)	51	14
1	B	144/135 (107%)	139 (96%)	5 (4%)	43	9
All	All	287/270 (106%)	278 (97%)	9 (3%)	47	12

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	ASP
1	A	37	LYS
1	A	45	PHE
1	A	144	ARG
1	B	45	PHE
1	B	144	ARG
1	B	157[A]	MET
1	B	157[B]	MET
1	B	169	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

36 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	201	1,2	14,14,15	0.39	0	15,19,21	0.39	0
2	NAG	A	202	3,2	14,14,15	0.30	0	15,19,21	0.36	0
3	BMA	A	203	2,5,4	11,11,12	1.04	1 (9%)	15,15,17	1.01	2 (13%)
4	MAN	A	204	3	11,11,12	1.12	1 (9%)	15,15,17	0.90	0
5	XYP	A	205	3	9,9,10	1.92	3 (33%)	12,12,14	1.06	1 (8%)
4	MAN	A	206	-	11,11,12	0.73	0	15,15,17	1.33	3 (20%)
6	FUC	A	207	-	10,10,11	1.18	1 (10%)	13,14,16	0.88	0
2	NAG	A	208	1,2	14,14,15	0.45	0	15,19,21	0.58	0
2	NAG	A	209	2	14,14,15	0.32	0	15,19,21	0.28	0
7	EDO	A	210	-	3,3,3	0.44	0	2,2,2	0.25	0
7	EDO	A	211	-	3,3,3	0.46	0	2,2,2	0.64	0
7	EDO	A	212	-	3,3,3	0.38	0	2,2,2	0.23	0
7	EDO	A	213	-	3,3,3	0.45	0	2,2,2	0.27	0
7	EDO	A	214	-	3,3,3	0.37	0	2,2,2	0.42	0
8	GLY	A	215	-	1,4,4	0.52	0	0,4,4	0.00	-
8	GLY	A	216	-	1,4,4	0.51	0	0,4,4	0.00	-
2	NAG	B	201	1,2,6	14,14,15	0.77	1 (7%)	15,19,21	0.45	0
2	NAG	B	202	3,2	14,14,15	0.39	0	15,19,21	0.41	0
3	BMA	B	203	2,5,4	11,11,12	1.64	1 (9%)	15,15,17	1.24	1 (6%)
4	MAN	B	204	3	11,11,12	0.85	0	15,15,17	1.12	2 (13%)
5	XYP	B	205	3	9,9,10	2.12	4 (44%)	12,12,14	1.27	1 (8%)
6	FUC	B	206	2	10,10,11	1.09	1 (10%)	13,14,16	0.85	0
4	MAN	B	207	-	11,11,12	0.77	0	15,15,17	1.27	3 (20%)
2	NAG	B	208	1,2,6	14,14,15	0.32	0	15,19,21	0.31	0
2	NAG	B	209	3,2	14,14,15	0.38	0	15,19,21	0.31	0
3	BMA	B	210	2,5,4	11,11,12	0.65	0	15,15,17	0.84	0
4	MAN	B	211	3	11,11,12	0.90	0	15,15,17	1.01	1 (6%)
4	MAN	B	212	3	11,11,12	0.63	0	15,15,17	1.16	1 (6%)
5	XYP	B	213	3	9,9,10	1.93	3 (33%)	12,12,14	0.76	0
6	FUC	B	214	2	10,10,11	0.82	0	13,14,16	0.74	0
7	EDO	B	215	-	3,3,3	0.46	0	2,2,2	0.44	0
7	EDO	B	216	-	3,3,3	0.41	0	2,2,2	0.29	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	EDO	B	217	-	3,3,3	0.41	0	2,2,2	0.46	0
8	GLY	B	218	-	1,4,4	0.48	0	0,4,4	0.00	-
8	GLY	B	219	-	1,4,4	0.51	0	0,4,4	0.00	-
8	GLY	B	220	-	1,4,4	0.53	0	0,4,4	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	201	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	202	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	203	2,5,4	-	0/2/19/22	0/1/1/1
4	MAN	A	204	3	-	0/2/19/22	0/1/1/1
5	XYP	A	205	3	-	0/0/14/17	0/1/1/1
4	MAN	A	206	-	-	0/2/19/22	0/1/1/1
6	FUC	A	207	-	-	0/0/17/20	0/1/1/1
2	NAG	A	208	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	209	2	-	0/6/23/26	0/1/1/1
7	EDO	A	210	-	-	0/1/1/1	0/0/0/0
7	EDO	A	211	-	-	0/1/1/1	0/0/0/0
7	EDO	A	212	-	-	0/1/1/1	0/0/0/0
7	EDO	A	213	-	-	0/1/1/1	0/0/0/0
7	EDO	A	214	-	-	0/1/1/1	0/0/0/0
8	GLY	A	215	-	-	0/0/2/2	0/0/0/0
8	GLY	A	216	-	-	0/0/2/2	0/0/0/0
2	NAG	B	201	1,2,6	-	0/6/23/26	0/1/1/1
2	NAG	B	202	3,2	-	0/6/23/26	0/1/1/1
3	BMA	B	203	2,5,4	-	0/2/19/22	0/1/1/1
4	MAN	B	204	3	-	0/2/19/22	0/1/1/1
5	XYP	B	205	3	-	0/0/14/17	0/1/1/1
6	FUC	B	206	2	-	0/0/17/20	0/1/1/1
4	MAN	B	207	-	-	0/2/19/22	0/1/1/1
2	NAG	B	208	1,2,6	-	0/6/23/26	0/1/1/1
2	NAG	B	209	3,2	-	0/6/23/26	0/1/1/1
3	BMA	B	210	2,5,4	-	0/2/19/22	0/1/1/1
4	MAN	B	211	3	-	0/2/19/22	0/1/1/1
4	MAN	B	212	3	-	0/2/19/22	0/1/1/1
5	XYP	B	213	3	-	0/0/14/17	0/1/1/1
6	FUC	B	214	2	-	0/0/17/20	0/1/1/1
7	EDO	B	215	-	-	0/1/1/1	0/0/0/0
7	EDO	B	216	-	-	0/1/1/1	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	B	217	-	-	0/1/1/1	0/0/0/0
8	GLY	B	218	-	-	0/0/2/2	0/0/0/0
8	GLY	B	219	-	-	0/0/2/2	0/0/0/0
8	GLY	B	220	-	-	0/0/2/2	0/0/0/0

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	203	BMA	O5-C1	-4.42	1.36	1.43
6	A	207	FUC	O5-C1	-3.08	1.38	1.43
4	A	204	MAN	O5-C1	-2.85	1.39	1.43
5	B	205	XYP	C2B-C3B	-2.78	1.48	1.52
6	B	206	FUC	O5-C1	-2.66	1.39	1.43
3	A	203	BMA	O5-C1	-2.61	1.39	1.43
5	B	205	XYP	C4B-C3B	-2.51	1.49	1.52
5	B	213	XYP	C2B-C3B	-2.36	1.49	1.52
2	B	201	NAG	O5-C1	-2.25	1.40	1.43
5	A	205	XYP	C2B-C3B	-2.04	1.49	1.52
5	A	205	XYP	O5B-C5B	2.67	1.48	1.42
5	B	213	XYP	O5B-C5B	2.83	1.48	1.42
5	B	205	XYP	O5B-C5B	2.86	1.48	1.42
5	B	213	XYP	O5B-C1B	3.81	1.50	1.42
5	B	205	XYP	O5B-C1B	3.81	1.50	1.42
5	A	205	XYP	O5B-C1B	4.14	1.51	1.42

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	203	BMA	C1-C2-C3	-3.03	105.88	109.55
3	A	203	BMA	O5-C5-C4	-2.66	105.74	110.13
4	B	207	MAN	O2-C2-C3	-2.58	104.99	110.19
4	A	206	MAN	O2-C2-C3	-2.50	105.15	110.19
4	B	212	MAN	O2-C2-C3	-2.38	105.38	110.19
4	B	204	MAN	O2-C2-C3	-2.27	105.62	110.19
3	A	203	BMA	O2-C2-C3	-2.23	105.69	110.19
4	B	207	MAN	C1-C2-C3	2.02	112.00	109.55
4	B	204	MAN	O5-C1-C2	2.10	114.25	110.89
5	A	205	XYP	C5B-C4B-C3B	2.12	112.11	109.55
4	B	207	MAN	C1-O5-C5	2.34	115.59	112.14
4	A	206	MAN	C1-O5-C5	2.35	115.60	112.14
4	A	206	MAN	O5-C1-C2	2.39	114.71	110.89
4	B	211	MAN	C1-O5-C5	2.40	115.67	112.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	205	XYP	C1B-C2B-C3B	3.75	114.09	109.55

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	208	NAG	2	0
3	B	203	BMA	5	0
4	B	207	MAN	5	0
4	B	211	MAN	0	1
8	B	218	GLY	1	0
8	B	219	GLY	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	153/158 (96%)	0.38	12 (7%) 16 14	14, 21, 37, 66	0
1	B	155/158 (98%)	0.33	12 (7%) 16 14	14, 22, 43, 63	0
All	All	308/316 (97%)	0.36	24 (7%) 16 14	14, 22, 41, 66	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	36	GLN	4.0
1	B	37	LYS	3.3
1	A	79	PHE	3.2
1	A	65	ILE	3.1
1	B	125	THR	3.1
1	A	125	THR	3.0
1	A	35	ASP	2.8
1	A	81	ILE	2.8
1	A	80	VAL	2.7
1	B	171	GLY	2.6
1	A	103	PHE	2.5
1	B	169	PHE	2.5
1	B	79	PHE	2.5
1	B	65	ILE	2.3
1	A	66	VAL	2.3
1	B	157[A]	MET	2.3
1	A	105	PHE	2.2
1	B	80	VAL	2.2
1	B	156	PHE	2.1
1	B	140	MET	2.1
1	B	103	PHE	2.1
1	A	115	TRP	2.1
1	B	142	PRO	2.0
1	A	64	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	EDO	B	217	4/4	0.83	0.39	10.86	23,36,45,47	10
7	EDO	B	215	4/4	0.92	0.24	9.23	32,40,55,66	10
8	GLY	B	218	5/5	0.63	0.27	7.20	38,41,46,49	7
7	EDO	A	213	4/4	0.79	0.30	6.23	29,35,44,53	10
8	GLY	A	216	5/5	0.81	0.34	4.51	47,60,73,73	10
7	EDO	A	210	4/4	0.91	0.16	3.69	26,36,53,63	10
8	GLY	A	215	5/5	0.81	0.17	3.49	38,44,46,47	7
7	EDO	A	214	4/4	0.91	0.24	3.19	30,42,50,54	0
2	NAG	A	208	14/15	0.78	0.25	2.89	40,55,76,78	0
4	MAN	B	211	11/12	0.64	0.38	2.86	59,75,89,101	0
8	GLY	B	220	5/5	0.81	0.21	2.72	26,37,57,57	10
4	MAN	B	204	11/12	0.94	0.13	2.38	27,32,42,57	0
7	EDO	A	212	4/4	0.96	0.13	1.88	22,35,44,44	10
8	GLY	B	219	5/5	0.94	0.20	1.60	41,48,56,56	10
7	EDO	B	216	4/4	0.96	0.14	1.53	23,40,48,48	10
7	EDO	A	211	4/4	0.85	0.17	1.17	30,44,46,55	0
3	BMA	B	203	11/12	0.94	0.09	0.48	23,27,36,36	0
2	NAG	B	201	14/15	0.97	0.07	0.21	18,23,44,44	0
4	MAN	A	204	11/12	0.96	0.09	-0.07	23,31,40,63	0
2	NAG	A	201	14/15	0.96	0.07	-0.44	19,25,37,37	0
2	NAG	B	202	14/15	0.96	0.07	-0.77	19,25,34,41	0
2	NAG	A	202	14/15	0.97	0.07	-0.88	20,25,31,45	0
4	MAN	A	206	11/12	0.88	0.16	-	37,50,63,76	0
6	FUC	A	207	10/11	0.90	0.17	-	33,45,58,70	0
5	XYP	B	205	9/10	0.94	0.18	-	30,46,62,71	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAG	B	209	14/15	0.86	0.27	-	47,65,87,87	0
2	NAG	B	208	14/15	0.74	0.26	-	50,58,70,87	0
5	XYP	A	205	9/10	0.91	0.13	-	29,39,48,52	0
3	BMA	A	203	11/12	0.95	0.07	-	23,29,37,39	0
4	MAN	B	207	11/12	0.85	0.15	-	25,44,57,62	0
5	XYP	B	213	9/10	0.87	0.39	-	64,79,95,95	0
4	MAN	B	212	11/12	0.74	0.43	-	75,83,97,100	0
6	FUC	B	214	10/11	0.81	0.28	-	52,67,86,92	0
6	FUC	B	206	10/11	0.94	0.12	-	28,35,46,48	0
3	BMA	B	210	11/12	0.74	0.28	-	65,71,85,85	0
2	NAG	A	209	14/15	0.65	0.40	-	68,90,113,119	0

6.5 Other polymers [i](#)

There are no such residues in this entry.