



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2016 – 10:42 AM EDT

PDB ID : 5LBV  
Title : Structural basis of zika and dengue virus potent antibody cross-neutralization  
Authors : Barba-Spaeth, G.  
Deposited on : 2016-06-17  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : unknown  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027939  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	447	<div> <div>5%</div> <div> <div></div> <div>78%</div> <div>9%</div> <div>13%</div> </div> </div>
1	B	447	<div> <div>3%</div> <div> <div></div> <div>76%</div> <div>13%</div> <div>12%</div> </div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6074 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called envelope protein E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	390	Total	C	N	O	S	0	0	0
			2961	1850	514	573	24			
1	B	394	Total	C	N	O	S	0	0	0
			3011	1882	525	578	26			

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	409	GLY	-	expression tag	UNP A0A120IIH9
A	410	PRO	-	expression tag	UNP A0A120IIH9
A	411	PHE	-	expression tag	UNP A0A120IIH9
A	412	GLU	-	expression tag	UNP A0A120IIH9
A	413	ASP	-	expression tag	UNP A0A120IIH9
A	414	ASP	-	expression tag	UNP A0A120IIH9
A	415	ASP	-	expression tag	UNP A0A120IIH9
A	416	ASP	-	expression tag	UNP A0A120IIH9
A	417	LYS	-	expression tag	UNP A0A120IIH9
A	418	ALA	-	expression tag	UNP A0A120IIH9
A	419	GLY	-	expression tag	UNP A0A120IIH9
A	420	TRP	-	expression tag	UNP A0A120IIH9
A	421	SER	-	expression tag	UNP A0A120IIH9
A	422	HIS	-	expression tag	UNP A0A120IIH9
A	423	PRO	-	expression tag	UNP A0A120IIH9
A	424	GLN	-	expression tag	UNP A0A120IIH9
A	425	PHE	-	expression tag	UNP A0A120IIH9
A	426	GLU	-	expression tag	UNP A0A120IIH9
A	427	LYS	-	expression tag	UNP A0A120IIH9
A	428	GLY	-	expression tag	UNP A0A120IIH9
A	429	GLY	-	expression tag	UNP A0A120IIH9
A	430	GLY	-	expression tag	UNP A0A120IIH9
A	431	SER	-	expression tag	UNP A0A120IIH9
A	432	GLY	-	expression tag	UNP A0A120IIH9
A	433	GLY	-	expression tag	UNP A0A120IIH9

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	434	GLY	-	expression tag	UNP A0A120IIH9
A	435	SER	-	expression tag	UNP A0A120IIH9
A	436	GLY	-	expression tag	UNP A0A120IIH9
A	437	GLY	-	expression tag	UNP A0A120IIH9
A	438	GLY	-	expression tag	UNP A0A120IIH9
A	439	SER	-	expression tag	UNP A0A120IIH9
A	440	TRP	-	expression tag	UNP A0A120IIH9
A	441	SER	-	expression tag	UNP A0A120IIH9
A	442	HIS	-	expression tag	UNP A0A120IIH9
A	443	PRO	-	expression tag	UNP A0A120IIH9
A	444	GLN	-	expression tag	UNP A0A120IIH9
A	445	PHE	-	expression tag	UNP A0A120IIH9
A	446	GLU	-	expression tag	UNP A0A120IIH9
A	447	LYS	-	expression tag	UNP A0A120IIH9
B	409	GLY	-	expression tag	UNP A0A120IIH9
B	410	PRO	-	expression tag	UNP A0A120IIH9
B	411	PHE	-	expression tag	UNP A0A120IIH9
B	412	GLU	-	expression tag	UNP A0A120IIH9
B	413	ASP	-	expression tag	UNP A0A120IIH9
B	414	ASP	-	expression tag	UNP A0A120IIH9
B	415	ASP	-	expression tag	UNP A0A120IIH9
B	416	ASP	-	expression tag	UNP A0A120IIH9
B	417	LYS	-	expression tag	UNP A0A120IIH9
B	418	ALA	-	expression tag	UNP A0A120IIH9
B	419	GLY	-	expression tag	UNP A0A120IIH9
B	420	TRP	-	expression tag	UNP A0A120IIH9
B	421	SER	-	expression tag	UNP A0A120IIH9
B	422	HIS	-	expression tag	UNP A0A120IIH9
B	423	PRO	-	expression tag	UNP A0A120IIH9
B	424	GLN	-	expression tag	UNP A0A120IIH9
B	425	PHE	-	expression tag	UNP A0A120IIH9
B	426	GLU	-	expression tag	UNP A0A120IIH9
B	427	LYS	-	expression tag	UNP A0A120IIH9
B	428	GLY	-	expression tag	UNP A0A120IIH9
B	429	GLY	-	expression tag	UNP A0A120IIH9
B	430	GLY	-	expression tag	UNP A0A120IIH9
B	431	SER	-	expression tag	UNP A0A120IIH9
B	432	GLY	-	expression tag	UNP A0A120IIH9
B	433	GLY	-	expression tag	UNP A0A120IIH9
B	434	GLY	-	expression tag	UNP A0A120IIH9
B	435	SER	-	expression tag	UNP A0A120IIH9
B	436	GLY	-	expression tag	UNP A0A120IIH9

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	437	GLY	-	expression tag	UNP A0A120IIH9
B	438	GLY	-	expression tag	UNP A0A120IIH9
B	439	SER	-	expression tag	UNP A0A120IIH9
B	440	TRP	-	expression tag	UNP A0A120IIH9
B	441	SER	-	expression tag	UNP A0A120IIH9
B	442	HIS	-	expression tag	UNP A0A120IIH9
B	443	PRO	-	expression tag	UNP A0A120IIH9
B	444	GLN	-	expression tag	UNP A0A120IIH9
B	445	PHE	-	expression tag	UNP A0A120IIH9
B	446	GLU	-	expression tag	UNP A0A120IIH9
B	447	LYS	-	expression tag	UNP A0A120IIH9

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is BETA-D-MANNOSE (three-letter code: BMA) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula:  $C_6H_{12}O_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Na	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	17	Total	O	0	0
			17	17		
7	B	24	Total	O	0	0
			24	24		





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.63Å 213.57Å 124.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.55 – 2.20 38.55 – 2.20	Depositor EDS
% Data completeness (in resolution range)	66.4 (38.55-2.20) 66.1 (38.55-2.20)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.65 (at 2.20Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.207 , 0.239 0.227 , 0.262	Depositor DCC
$R_{free}$ test set	1466 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	45.9	Xtriage
Anisotropy	0.317	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 40.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6074	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, NA, BMA, NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/3019	0.74	0/4088
1	B	0.52	0/3074	0.73	0/4162
All	All	0.51	0/6093	0.73	0/8250

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2961	0	2876	20	0
1	B	3011	0	2936	25	0
2	B	28	0	23	0	0
3	B	11	0	9	0	0
4	B	10	0	10	0	0
5	B	11	0	10	0	0
6	B	1	0	0	0	0
7	A	17	0	0	0	0
7	B	24	0	0	0	0
All	All	6074	0	5864	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:97:VAL:HG11	1:B:250:ALA:HA	1.50	0.94
1:A:97:VAL:HG11	1:A:250:ALA:HA	1.55	0.86
1:B:65:ILE:HG12	1:B:257:VAL:HG13	1.62	0.81
1:B:99:ARG:HA	1:B:103:ASN:HD22	1.54	0.73
1:A:65:ILE:HG12	1:A:257:VAL:HG23	1.81	0.62

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	380/447 (85%)	374 (98%)	5 (1%)	1 (0%)	46	50
1	B	388/447 (87%)	378 (97%)	8 (2%)	2 (0%)	34	35
All	All	768/894 (86%)	752 (98%)	13 (2%)	3 (0%)	39	42

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	193	ARG
1	B	16	SER
1	B	15	MET

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/367 (88%)	310 (97%)	11 (3%)	44	54
1	B	329/367 (90%)	319 (97%)	10 (3%)	48	60
All	All	650/734 (89%)	629 (97%)	21 (3%)	46	57

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	368	SER
1	B	19	THR
1	B	301	LYS
1	A	357	ARG
1	B	357	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	163	ASN
1	A	208	ASN
1	B	103	ASN
1	B	163	ASN
1	B	208	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	B	501	1,2,4	14,14,15	0.26	0	15,19,21	0.51	0
2	NAG	B	502	3,2	14,14,15	0.33	0	15,19,21	0.58	0
3	BMA	B	503	2,5	11,11,12	0.37	0	15,15,17	1.37	1 (6%)
4	FUC	B	504	2	10,10,11	0.37	0	13,14,16	1.05	1 (7%)
5	MAN	B	505	3	11,11,12	0.33	0	15,15,17	1.23	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	501	1,2,4	-	0/6/23/26	0/1/1/1
2	NAG	B	502	3,2	-	0/6/23/26	0/1/1/1
3	BMA	B	503	2,5	-	0/2/19/22	0/1/1/1
4	FUC	B	504	2	-	0/0/17/20	0/1/1/1
5	MAN	B	505	3	-	0/2/19/22	1/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	504	FUC	C3-C4-C5	2.24	113.00	109.66
3	B	503	BMA	C1-O5-C5	3.15	116.77	112.14
5	B	505	MAN	C1-O5-C5	4.54	118.82	112.14

There are no chirality outliers.

There are no torsion outliers.

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	505	MAN	C1-C2-C3-C4-C5-O5

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	390/447 (87%)	0.30	23 (5%)	26 25	34, 68, 103, 125	0
1	B	394/447 (88%)	0.04	12 (3%)	54 53	28, 54, 93, 120	0
All	All	784/894 (87%)	0.17	35 (4%)	37 36	28, 61, 100, 125	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	271	GLY	4.8
1	A	52	ASN	4.8
1	B	271	GLY	4.2
1	B	251	LYS	3.8
1	A	156	THR	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	BMA	B	503	11/12	0.89	0.15	1.51	67,72,77,80	0
5	MAN	B	505	11/12	0.85	0.27	0.85	82,88,89,89	0
2	NAG	B	501	14/15	0.89	0.14	-0.30	75,78,80,81	0
6	NA	B	506	1/1	0.93	0.09	-1.20	51,51,51,51	0
4	FUC	B	504	10/11	0.88	0.15	-	64,71,74,75	0
2	NAG	B	502	14/15	0.83	0.19	-	70,74,82,83	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.