



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:22 PM GMT

PDB ID : 4LEO
Title : Crystal structure of anti-HER3 Fab RG7116 in complex with the extracellular domains of human Her3 (ERBB3)
Authors : Schiller C.B.; Hopfner K.P.
Deposited on : 2013-06-26
Resolution : 2.64 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3377 (2.68-2.60)
Clashscore	102246	3781 (2.68-2.60)
Ramachandran outliers	100387	3722 (2.68-2.60)
Sidechain outliers	100360	3722 (2.68-2.60)
RSRZ outliers	91569	3388 (2.68-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	<div> <div>2%</div> <div>83% 10% 6%</div> </div>
2	B	220	<div> <div>%</div> <div>88% 11%</div> </div>
3	C	621	<div> <div>3%</div> <div>83% 13% . .</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	C	705	-	-	-	X
4	NAG	C	706	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8185 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RG7116 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	60	0	0
			1618	1020	273	318	7			

- Molecule 2 is a protein called RG7116 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	219	Total	C	N	O	S	61	0	0
			1697	1062	282	348	5			

- Molecule 3 is a protein called Receptor tyrosine-protein kinase erbB-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	601	Total	C	N	O	S	257	0	0
			4622	2861	834	868	59			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	613	ALA	-	EXPRESSION TAG	UNP P21860
C	614	ALA	-	EXPRESSION TAG	UNP P21860
C	615	ALA	-	EXPRESSION TAG	UNP P21860
C	616	LEU	-	EXPRESSION TAG	UNP P21860
C	617	GLU	-	EXPRESSION TAG	UNP P21860
C	618	VAL	-	EXPRESSION TAG	UNP P21860
C	619	LEU	-	EXPRESSION TAG	UNP P21860
C	620	PHE	-	EXPRESSION TAG	UNP P21860
C	621	GLN	-	EXPRESSION TAG	UNP P21860

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	5	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		



- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	29	Total	O	0	0
			29	29		
6	B	38	Total	O	0	0
			38	38		
6	C	83	Total	O	0	0
			83	83		

- Molecule 1: RG7116 Fab heavy chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.46 Å 59.36 Å 103.66 Å 97.64° 106.85° 96.83°	Depositor
Resolution (Å)	48.83 – 2.64 48.83 – 2.64	Depositor EDS
% Data completeness (in resolution range)	95.8 (48.83-2.64) 90.4 (48.83-2.64)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 2.65 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.228 , 0.258 0.226 , 0.258	Depositor DCC
R_{free} test set	1710 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	43.6	Xtriage
Anisotropy	0.464	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 34226 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	8185	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/1658	0.41	0/2261
2	B	0.32	0/1734	0.41	0/2356
3	C	0.28	0/4733	0.48	2/6422 (0.0%)
All	All	0.28	0/8125	0.45	2/11039 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	116	ARG	NE-CZ-NH2	9.43	125.01	120.30
3	C	307	LEU	CA-CB-CG	6.75	130.83	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1618	0	1583	11	0
2	B	1697	0	1638	11	0
3	C	4622	0	4386	45	0
4	C	70	0	65	0	0
5	C	28	0	25	0	0
6	A	29	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	38	0	0	0	0
6	C	83	0	0	2	0
All	All	8185	0	7697	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 63 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:34:ASN:HB3	3:C:154:GLU:HG2	1.60	0.84
3:C:247:ASN:HB3	3:C:250:THR:HG22	1.67	0.76
1:A:18:VAL:HG23	1:A:86:LEU:HD11	1.67	0.76
3:C:522:ARG:NH2	3:C:540:MET:O	2.20	0.75
2:B:34:ASN:CB	3:C:154:GLU:HG2	2.18	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/227 (92%)	204 (97%)	6 (3%)	0	100	100
2	B	217/220 (99%)	212 (98%)	5 (2%)	0	100	100
3	C	597/621 (96%)	556 (93%)	41 (7%)	0	100	100
All	All	1024/1068 (96%)	972 (95%)	52 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/194 (94%)	169 (93%)	13 (7%)	18	34
2	B	194/195 (100%)	186 (96%)	8 (4%)	37	64
3	C	521/535 (97%)	494 (95%)	27 (5%)	29	52
All	All	897/924 (97%)	849 (95%)	48 (5%)	27	50

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	13	THR
3	C	106	THR
3	C	448	SER
3	C	14	LEU
3	C	35	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	28	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	C	703	3,5	14,14,15	1.21	1 (7%)	15,19,21	1.46	2 (13%)
5	NAG	C	704	5	14,14,15	1.20	1 (7%)	15,19,21	1.58	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	703	3,5	-	0/6/23/26	0/1/1/1
5	NAG	C	704	5	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	704	NAG	C1-C2	2.41	1.55	1.52
5	C	703	NAG	C1-C2	2.72	1.56	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	703	NAG	C4-C3-C2	2.01	114.35	111.23
5	C	704	NAG	C3-C4-C5	2.19	114.01	110.20
5	C	704	NAG	C4-C3-C2	2.42	114.99	111.23
5	C	704	NAG	O3-C3-C2	3.10	115.25	109.11
5	C	703	NAG	O3-C3-C2	3.14	115.33	109.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	C	701	3	14,14,15	1.15	1 (7%)	15,19,21	1.56	3 (20%)
4	NAG	C	702	3	14,14,15	1.19	1 (7%)	15,19,21	1.62	3 (20%)
4	NAG	C	705	3	14,14,15	1.21	1 (7%)	15,19,21	1.60	4 (26%)
4	NAG	C	706	3	14,14,15	1.18	1 (7%)	15,19,21	1.66	5 (33%)
4	NAG	C	707	3	14,14,15	1.20	1 (7%)	15,19,21	1.59	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	701	3	-	0/6/23/26	0/1/1/1
4	NAG	C	702	3	-	0/6/23/26	0/1/1/1
4	NAG	C	705	3	-	0/6/23/26	0/1/1/1
4	NAG	C	706	3	-	0/6/23/26	0/1/1/1
4	NAG	C	707	3	-	0/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	702	NAG	C1-C2	2.01	1.55	1.52
4	C	706	NAG	C1-C2	2.12	1.55	1.52
4	C	701	NAG	C1-C2	2.22	1.55	1.52
4	C	707	NAG	C1-C2	2.30	1.55	1.52
4	C	705	NAG	C1-C2	2.43	1.55	1.52

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	C	706	NAG	C2-N2-C7	-2.55	119.77	123.04
4	C	705	NAG	O4-C4-C3	-2.01	105.82	110.34
4	C	706	NAG	O4-C4-C3	-2.00	105.83	110.34
4	C	701	NAG	C3-C4-C5	2.09	113.84	110.20
4	C	707	NAG	C4-C3-C2	2.21	114.66	111.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/227 (94%)	0.21	5 (2%) 64 57	29, 52, 77, 97	21 (9%)
2	B	219/220 (99%)	0.10	2 (0%) 85 83	29, 49, 80, 106	25 (11%)
3	C	601/621 (96%)	0.29	16 (2%) 58 52	28, 54, 88, 124	81 (13%)
All	All	1034/1068 (96%)	0.23	23 (2%) 65 59	28, 52, 86, 124	127 (12%)

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	587	VAL	4.8
3	C	37	CYS	3.9
3	C	574	VAL	3.8
3	C	296	ASN	3.7
3	C	300	MET	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	NAG	C	703	14/15	0.87	0.15	-0.86	57,60,60,61	0
5	NAG	C	704	14/15	0.75	0.23	-	96,97,99,99	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	NAG	C	706	14/15	0.71	0.35	6.33	106,109,111,111	0
4	NAG	C	705	14/15	0.87	0.23	2.21	64,66,67,67	0
4	NAG	C	701	14/15	0.95	0.17	0.04	45,47,49,49	0
4	NAG	C	707	14/15	0.83	0.19	-0.07	80,82,83,83	0
4	NAG	C	702	14/15	0.86	0.14	-	75,76,78,79	5

6.5 Other polymers [i](#)

There are no such residues in this entry.