



# wwPDB NMR Structure Validation Summary Report ⓘ

Apr 27, 2016 – 01:12 AM BST

PDB ID : 2LHU  
Title : Structural Insight into the Unique Cardiac Myosin Binding Protein-C Motif:  
A Partially Folded Domain  
Authors : Howarth, J.W.; Rosevear, P.R.; Ramisetty, S.; Nolan, K.; Sadayappan, S.  
Deposited on : 2011-08-18

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 41%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	124	 27% . 72%

## 2 Ensemble composition and analysis ⓘ

This entry contains 20 models. Model 15 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:317-A:349 (33)	0.32	15

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20
Single-model clusters	8

### 3 Entry composition


There is only 1 type of molecule in this entry. The entry contains 589 atoms, of which 295 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Mybpc3 protein.

Mol	Chain	Residues	Atoms						Trace
1	A	35	Total	C	H	N	O	S	0
			589	187	295	54	52	1	

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	234	MET	-	EXPRESSION TAG	UNP A9JR55
A	235	GLY	-	EXPRESSION TAG	UNP A9JR55
A	236	SER	-	EXPRESSION TAG	UNP A9JR55
A	237	SER	-	EXPRESSION TAG	UNP A9JR55
A	238	HIS	-	EXPRESSION TAG	UNP A9JR55
A	239	HIS	-	EXPRESSION TAG	UNP A9JR55
A	240	HIS	-	EXPRESSION TAG	UNP A9JR55
A	241	HIS	-	EXPRESSION TAG	UNP A9JR55
A	242	HIS	-	EXPRESSION TAG	UNP A9JR55
A	243	HIS	-	EXPRESSION TAG	UNP A9JR55
A	244	SER	-	EXPRESSION TAG	UNP A9JR55
A	245	SER	-	EXPRESSION TAG	UNP A9JR55
A	246	GLY	-	EXPRESSION TAG	UNP A9JR55
A	247	LEU	-	EXPRESSION TAG	UNP A9JR55
A	249	PRO	-	EXPRESSION TAG	UNP A9JR55
A	250	ARG	-	EXPRESSION TAG	UNP A9JR55
A	251	GLY	-	EXPRESSION TAG	UNP A9JR55
A	252	SER	-	EXPRESSION TAG	UNP A9JR55
A	253	HIS	-	EXPRESSION TAG	UNP A9JR55
A	254	MET	-	EXPRESSION TAG	UNP A9JR55



## 5 Refinement protocol and experimental data overview

Of the 5000 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CS-Rosetta	structure solution	
CS-Rosetta	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2lhu_cs.str
Number of chemical shift lists	1
Total number of shifts	608
Number of shifts mapped to atoms	608
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	41%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	277	285	286	1±1
All	All	5540	5700	5720	16

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

5 of 7 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:348:LEU:C	1:A:348:LEU:HD23	0.61	2.16	14	5
1:A:348:LEU:HD23	1:A:348:LEU:C	0.58	2.18	9	3
1:A:348:LEU:O	1:A:348:LEU:HD23	0.56	2.00	14	3
1:A:348:LEU:CD2	1:A:348:LEU:C	0.49	2.80	17	2
1:A:348:LEU:C	1:A:348:LEU:CD2	0.49	2.80	14	1

### 6.3 Torsion angles [i](#)

#### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	32/124 (26%)	32±0 (100±1%)	0±0 (0±1%)	0±0 (0±0%)	100	100
All	All	640/2480 (26%)	639 (100%)	1 (0%)	0 (0%)	100	100

There are no Ramachandran outliers.

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	29/104 (28%)	29±0 (99±1%)	0±0 (1±1%)	89	98
All	All	580/2080 (28%)	576 (99%)	4 (1%)	89	98

All 1 unique residues with a non-rotameric sidechain are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	321	LEU	4

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.6 Ligand geometry ⓘ

There are no ligands in this entry.



## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 41% for the well-defined parts and 42% for the entire structure.

### 7.1 Chemical shift list 1

File name: 2lhu\_cs.str

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	608
Number of shifts mapped to atoms	608
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

#### 7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	102	$0.11 \pm 0.15$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	94	$0.35 \pm 0.13$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	102	$-0.15 \pm 0.15$	None needed ( $< 0.5$ ppm)
$^{15}\text{N}$	99	$0.26 \pm 0.43$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 41%, i.e. 188 atoms were assigned a chemical shift out of a possible 454. 0 out of 6 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	158/161 (98%)	63/64 (98%)	64/66 (97%)	31/31 (100%)
Sidechain	30/258 (12%)	0/153 (0%)	30/89 (34%)	0/16 (0%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	0/35 (0%)	0/19 (0%)	0/15 (0%)	0/1 (0%)
Overall	188/454 (41%)	63/236 (27%)	94/170 (55%)	31/48 (65%)

#### 7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	298	LYS	NZ	120.41	49.86 – 18.16	27.3
1	A	331	ARG	CB	41.06	39.81 – 21.51	5.7

#### 7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

