



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 21, 2017 – 10:05 PM EST

PDB ID : 5LHZ  
Title : PB3 Domain of Human PLK4 in Complex with Coiled-Coil Domain of STIL  
Authors : Cottee, M.A.; Lea, S.M.  
Deposited on : 2016-07-13  
Resolution : 2.51 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	91	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>•</div> <div>14%</div> </div> </div>
1	B	91	<div> <div>3%</div> <div> <div></div> <div>78%</div> <div>• •</div> <div>16%</div> </div> </div>
1	C	91	<div> <div>2%</div> <div> <div></div> <div>80%</div> <div>•</div> <div>15%</div> </div> </div>
2	D	28	<div> <div>7%</div> <div> <div></div> <div>71%</div> <div>7%</div> <div>21%</div> </div> </div>
2	E	28	<div> <div>7%</div> <div> <div></div> <div>68%</div> <div>7%</div> <div>25%</div> </div> </div>
2	F	28	<div> <div>7%</div> <div> <div></div> <div>75%</div> <div>•</div> <div>21%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4731 atoms, of which 2389 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase PLK4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	78	Total	C	H	N	O	S	0	0	0
			1214	387	609	100	116	2			
1	B	76	Total	C	H	N	O	S	0	0	0
			1186	379	596	97	112	2			
1	C	77	Total	C	H	N	O	S	0	0	0
			1197	382	601	98	114	2			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	880	GLY	-	expression tag	UNP O00444
A	881	PRO	-	expression tag	UNP O00444
A	882	MET	-	expression tag	UNP O00444
A	883	GLY	-	expression tag	UNP O00444
B	880	GLY	-	expression tag	UNP O00444
B	881	PRO	-	expression tag	UNP O00444
B	882	MET	-	expression tag	UNP O00444
B	883	GLY	-	expression tag	UNP O00444
C	880	GLY	-	expression tag	UNP O00444
C	881	PRO	-	expression tag	UNP O00444
C	882	MET	-	expression tag	UNP O00444
C	883	GLY	-	expression tag	UNP O00444

- Molecule 2 is a protein called SCL-interrupting locus protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	D	22	Total	C	H	N	O	S	0	0	0
			382	113	197	37	34	1			
2	E	21	Total	C	H	N	O	S	0	0	0
			365	108	189	35	32	1			
2	F	22	Total	C	H	N	O	S	0	0	0
			382	113	197	37	34	1			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	723	GLY	-	expression tag	UNP Q15468
D	724	GLY	-	expression tag	UNP Q15468
D	725	SER	-	expression tag	UNP Q15468
E	723	GLY	-	expression tag	UNP Q15468
E	724	GLY	-	expression tag	UNP Q15468
E	725	SER	-	expression tag	UNP Q15468
F	723	GLY	-	expression tag	UNP Q15468
F	724	GLY	-	expression tag	UNP Q15468
F	725	SER	-	expression tag	UNP Q15468



- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	3	Total O 3 3	0	0
3	E	1	Total O 1 1	0	0
3	F	1	Total O 1 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Serine/threonine-protein kinase PLK4




- Molecule 1: Serine/threonine-protein kinase PLK4




- Molecule 1: Serine/threonine-protein kinase PLK4





- Molecule 2: SCL-interrupting locus protein



- Molecule 2: SCL-interrupting locus protein



- Molecule 2: SCL-interrupting locus protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.36 Å 68.36 Å 137.39 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.22 – 2.51 36.22 – 2.51	Depositor EDS
% Data completeness (in resolution range)	99.7 (36.22-2.51) 99.7 (36.22-2.51)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.63 (at 2.51 Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.250 , 0.282 0.248 , 0.280	Depositor DCC
$R_{free}$ test set	669 reflections (5.06%)	DCC
Wilson B-factor (Å <sup>2</sup> )	63.0	Xtriage
Anisotropy	0.331	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 47.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.098 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	4731	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/616	0.44	0/835
1	B	0.25	0/601	0.43	0/815
1	C	0.25	0/607	0.43	0/823
2	D	0.23	0/184	0.36	0/243
2	E	0.23	0/175	0.39	0/231
2	F	0.23	0/184	0.39	0/243
All	All	0.25	0/2367	0.42	0/3190

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	605	609	609	2	0
1	B	590	596	596	3	0
1	C	596	601	601	3	0
2	D	185	197	197	1	0
2	E	176	189	189	2	0
2	F	185	197	197	1	0
3	A	3	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
All	All	2342	2389	2389	8	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 8 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:921:ALA:O	2:F:739:GLN:NE2	2.18	0.76
1:A:930:SER:OG	1:A:932:ASN:OD1	2.13	0.65
1:B:921:ALA:O	2:E:739:GLN:NE2	2.31	0.63
1:C:930:SER:OG	1:C:932:ASN:OD1	2.16	0.63
1:A:923:VAL:O	2:D:739:GLN:NE2	2.44	0.48

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	76/91 (84%)	74 (97%)	2 (3%)	0	100	100
1	B	74/91 (81%)	72 (97%)	2 (3%)	0	100	100
1	C	75/91 (82%)	72 (96%)	3 (4%)	0	100	100
2	D	20/28 (71%)	20 (100%)	0	0	100	100
2	E	19/28 (68%)	19 (100%)	0	0	100	100
2	F	20/28 (71%)	20 (100%)	0	0	100	100
All	All	284/357 (80%)	277 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	69/79 (87%)	69 (100%)	0	100	100
1	B	67/79 (85%)	66 (98%)	1 (2%)	72	91
1	C	68/79 (86%)	68 (100%)	0	100	100
2	D	20/24 (83%)	19 (95%)	1 (5%)	30	53
2	E	19/24 (79%)	19 (100%)	0	100	100
2	F	20/24 (83%)	20 (100%)	0	100	100
All	All	263/309 (85%)	261 (99%)	2 (1%)	86	96

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	893	VAL
2	D	733	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	78/91 (85%)	0.45	2 (2%) 59 63	42, 58, 90, 102	0
1	B	76/91 (83%)	0.41	3 (3%) 43 48	48, 68, 97, 101	0
1	C	77/91 (84%)	0.52	2 (2%) 59 63	51, 69, 102, 107	0
2	D	22/28 (78%)	0.51	2 (9%) 11 12	51, 64, 99, 120	0
2	E	21/28 (75%)	0.84	2 (9%) 10 11	62, 80, 110, 124	0
2	F	22/28 (78%)	0.75	2 (9%) 11 12	65, 79, 98, 109	0
All	All	296/357 (82%)	0.51	13 (4%) 38 43	42, 68, 101, 124	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	962	PHE	4.6
1	A	887	LEU	3.8
1	B	961	MET	3.7
2	F	750	MET	3.2
2	E	749	LEU	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.