



Full wwPDB NMR Structure Validation Report ⓘ

Apr 27, 2016 – 01:01 AM BST

PDB ID : 2LIF
Title : Solution Structure of KKGF
Authors : Montserret, R.; Penin, F.
Deposited on : 2011-08-29

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 63%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	27	 89% 11%

2 Ensemble composition and analysis ⓘ

This entry contains 27 models.

Cyrange was unable to find well-defined residues.

Error message: Only domains with < 8 residues could be identified.

NmrClust was unable to cluster the ensemble.

Error message: Wrapper check: not enough residues in core to run NmrClust

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 428 atoms, of which 225 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Core protein p21.

Mol	Chain	Residues	Atoms						Trace
1	A	27	Total	C	H	N	O	S	0
			428	138	225	31	33	1	

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	LYS	-	SEE REMARK 999	UNP Q99IB8
A	2	LYS	-	SEE REMARK 999	UNP Q99IB8

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Core protein p21



4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Core protein p21




4.2.2 Score per residue for model 2

- Molecule 1: Core protein p21




4.2.3 Score per residue for model 3

- Molecule 1: Core protein p21




4.2.4 Score per residue for model 4

- Molecule 1: Core protein p21




4.2.5 Score per residue for model 5

- Molecule 1: Core protein p21




4.2.6 Score per residue for model 6

- Molecule 1: Core protein p21




4.2.7 Score per residue for model 7

- Molecule 1: Core protein p21




4.2.8 Score per residue for model 8

- Molecule 1: Core protein p21



4.2.9 Score per residue for model 9

- Molecule 1: Core protein p21




4.2.10 Score per residue for model 10

- Molecule 1: Core protein p21




4.2.11 Score per residue for model 11

- Molecule 1: Core protein p21




4.2.12 Score per residue for model 12

- Molecule 1: Core protein p21




4.2.13 Score per residue for model 13

- Molecule 1: Core protein p21




4.2.14 Score per residue for model 14

- Molecule 1: Core protein p21




4.2.15 Score per residue for model 15

- Molecule 1: Core protein p21



4.2.16 Score per residue for model 16

- Molecule 1: Core protein p21




4.2.17 Score per residue for model 17

- Molecule 1: Core protein p21




4.2.18 Score per residue for model 18

- Molecule 1: Core protein p21




4.2.19 Score per residue for model 19

- Molecule 1: Core protein p21



4.2.20 Score per residue for model 20

- Molecule 1: Core protein p21




4.2.21 Score per residue for model 21

- Molecule 1: Core protein p21




4.2.22 Score per residue for model 22


- Molecule 1: Core protein p21



4.2.23 Score per residue for model 23

- Molecule 1: Core protein p21

Chain A:  78% 22%



4.2.24 Score per residue for model 24

- Molecule 1: Core protein p21


Chain A:  93% 7%



4.2.25 Score per residue for model 25


- Molecule 1: Core protein p21

Chain A:  67% 33%



4.2.26 Score per residue for model 26

- Molecule 1: Core protein p21

Chain A:  85% 15%



4.2.27 Score per residue for model 27

- Molecule 1: Core protein p21

Chain A:  67% 26% 7%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *DGSA-distance geometry simulated annealing*.

Of the 50 calculated structures, 27 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	
Xplor	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2lif_cs.str
Number of chemical shift lists	1
Total number of shifts	239
Number of shifts mapped to atoms	239
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	63%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	203	225	228	2±2
All	All	5481	6075	6156	65

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:182:LEU:HA	1:A:185:ILE:HD12	0.74	1.57	21	3
1:A:185:ILE:HD13	1:A:186:THR:N	0.68	2.04	21	1
1:A:181:LEU:O	1:A:185:ILE:HG23	0.65	1.92	21	1
1:A:186:THR:O	1:A:189:VAL:HG22	0.62	1.94	1	1
1:A:194:VAL:HG13	1:A:195:LYS:HE2	0.57	1.75	27	1
1:A:187:VAL:CG2	1:A:188:PRO:HD3	0.55	2.32	14	3
1:A:182:LEU:HD13	1:A:185:ILE:HD11	0.55	1.79	13	2
1:A:187:VAL:HG23	1:A:188:PRO:HD3	0.53	1.79	14	2
1:A:174:PHE:O	1:A:178:LEU:HD13	0.53	2.04	8	1
1:A:190:SER:O	1:A:194:VAL:HG23	0.52	2.04	22	3
1:A:185:ILE:O	1:A:189:VAL:HG23	0.52	2.05	6	3
1:A:174:PHE:O	1:A:178:LEU:HD23	0.51	2.05	17	1
1:A:194:VAL:HG13	1:A:195:LYS:CE	0.50	2.36	27	1
1:A:187:VAL:N	1:A:188:PRO:CD	0.50	2.75	17	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:182:LEU:HD12	1:A:185:ILE:HD11	0.49	1.84	1	1
1:A:186:THR:O	1:A:189:VAL:HG12	0.48	2.07	12	2
1:A:191:ALA:HA	1:A:194:VAL:HG22	0.48	1.85	8	6
1:A:187:VAL:HB	1:A:188:PRO:CD	0.47	2.39	2	3
1:A:182:LEU:HA	1:A:185:ILE:CG1	0.47	2.40	2	8
1:A:182:LEU:HA	1:A:185:ILE:HG12	0.47	1.85	19	3
1:A:191:ALA:HA	1:A:194:VAL:CG2	0.47	2.40	17	4
1:A:183:SER:O	1:A:187:VAL:HG23	0.46	2.10	1	1
1:A:187:VAL:HB	1:A:188:PRO:HD3	0.44	1.89	21	4
1:A:182:LEU:CA	1:A:185:ILE:HD12	0.43	2.36	21	1
1:A:187:VAL:N	1:A:188:PRO:HD2	0.42	2.30	14	5
1:A:175:SER:HA	1:A:178:LEU:HB2	0.42	1.92	15	1
1:A:185:ILE:HA	1:A:188:PRO:HG2	0.40	1.93	27	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	25/27 (93%)	24±1 (96±3%)	1±1 (3±3%)	0±0 (0±1%)	50	83
All	All	675/729 (93%)	650 (96%)	23 (3%)	2 (0%)	50	83

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	171	GLY	1
1	A	173	PRO	1

6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	23/23 (100%)	19±2 (81±7%)	4±2 (19±7%)	5	39
All	All	621/621 (100%)	506 (81%)	115 (19%)	5	39

All 17 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	182	LEU	15
1	A	195	LYS	14
1	A	1	LYS	13
1	A	193	GLN	11
1	A	2	LYS	11
1	A	179	LEU	10
1	A	189	VAL	7
1	A	175	SER	6
1	A	181	LEU	6
1	A	178	LEU	5
1	A	172	PHE	4
1	A	174	PHE	4
1	A	177	PHE	3
1	A	184	CYS	2
1	A	183	SER	2
1	A	185	ILE	1
1	A	186	THR	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 63% for the well-defined parts and 63% for the entire structure.

7.1 Chemical shift list 1

File name: 2lif_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	239
Number of shifts mapped to atoms	239
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	27	-1.83 ± 0.57	Should be applied
$^{13}\text{C}_\beta$	26	-0.28 ± 0.24	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	—
^{15}N	0	—	—

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 63%, i.e. 206 atoms were assigned a chemical shift out of a possible 328. 1 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	75/131 (57%)	48/52 (92%)	27/54 (50%)	0/25 (0%)
Sidechain	131/170 (77%)	84/100 (84%)	47/66 (71%)	0/4 (0%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	0/27 (0%)	0/15 (0%)	0/12 (0%)	0/0 (—%)
Overall	206/328 (63%)	132/167 (79%)	74/132 (56%)	0/29 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 63%, i.e. 206 atoms were assigned a chemical shift out of a possible 328. 1 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	75/131 (57%)	48/52 (92%)	27/54 (50%)	0/25 (0%)
Sidechain	131/170 (77%)	84/100 (84%)	47/66 (71%)	0/4 (0%)
Aromatic	0/27 (0%)	0/15 (0%)	0/12 (0%)	0/0 (—%)
Overall	206/328 (63%)	132/167 (79%)	74/132 (56%)	0/29 (0%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

