



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 2, 2017 – 05:45 PM EST

PDB ID : 5LIV
Title : Crystal structure of myxobacterial CYP260A1
Authors : Carius, Y.; Khatri, Y.; Bernhardt, R.; Lancaster, C.R.D.
Deposited on : 2016-07-15
Resolution : 2.67 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028442
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2780 (2.70-2.66)
Clashscore	102246	3138 (2.70-2.66)
Ramachandran outliers	100387	3089 (2.70-2.66)
Sidechain outliers	100360	3089 (2.70-2.66)
RSRZ outliers	91569	2789 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	450	<div> <div>5%</div> <div> <div></div> <div>76%</div> <div>10%</div> <div>•</div> <div>12%</div> </div> </div>
1	B	450	<div> <div>9%</div> <div> <div></div> <div>72%</div> <div>13%</div> <div>•</div> <div>14%</div> </div> </div>
1	C	450	<div> <div>%</div> <div> <div></div> <div>73%</div> <div>13%</div> <div>•</div> <div>12%</div> </div> </div>
1	D	450	<div> <div>3%</div> <div> <div></div> <div>72%</div> <div>13%</div> <div>•</div> <div>14%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	502	-	-	-	X
3	GOL	B	504	-	-	-	X
3	GOL	C	504	-	-	-	X
3	GOL	C	505	-	-	X	-
3	GOL	C	506	-	-	X	-
3	GOL	D	507	-	-	-	X
5	SO4	C	503	-	-	-	X
5	SO4	D	510	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 12879 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 CYP260A1,Cytochrome P450 CYP260A1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	397	Total	C	N	O	S	0	2	0
			3092	1955	565	555	17			
1	B	389	Total	C	N	O	S	0	1	0
			3003	1908	536	543	16			
1	C	395	Total	C	N	O	S	0	2	0
			3091	1952	562	561	16			
1	D	389	Total	C	N	O	S	0	0	0
			2997	1896	542	543	16			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	445	HIS	-	expression tag	UNP A9FDB7
A	446	HIS	-	expression tag	UNP A9FDB7
A	447	HIS	-	expression tag	UNP A9FDB7
A	448	HIS	-	expression tag	UNP A9FDB7
A	449	HIS	-	expression tag	UNP A9FDB7
A	450	HIS	-	expression tag	UNP A9FDB7
B	445	HIS	-	expression tag	UNP A9FDB7
B	446	HIS	-	expression tag	UNP A9FDB7
B	447	HIS	-	expression tag	UNP A9FDB7
B	448	HIS	-	expression tag	UNP A9FDB7
B	449	HIS	-	expression tag	UNP A9FDB7
B	450	HIS	-	expression tag	UNP A9FDB7
C	445	HIS	-	expression tag	UNP A9FDB7
C	446	HIS	-	expression tag	UNP A9FDB7
C	447	HIS	-	expression tag	UNP A9FDB7
C	448	HIS	-	expression tag	UNP A9FDB7
C	449	HIS	-	expression tag	UNP A9FDB7
C	450	HIS	-	expression tag	UNP A9FDB7
D	445	HIS	-	expression tag	UNP A9FDB7
D	446	HIS	-	expression tag	UNP A9FDB7
D	447	HIS	-	expression tag	UNP A9FDB7

Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	448	HIS	-	expression tag	UNP A9FDB7
D	449	HIS	-	expression tag	UNP A9FDB7
D	450	HIS	-	expression tag	UNP A9FDB7

- # HEM

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	B	1	Total	C	O	S	0	0
			4	2	1	1		
4	C	1	Total	C	O	S	0	0
			4	2	1	1		
4	D	1	Total	C	O	S	0	0
			4	2	1	1		
4	D	1	Total	C	O	S	0	0
			4	2	1	1		
4	D	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	78	Total	O	0	0
			78	78		
7	B	66	Total	O	0	0
			66	66		
7	C	117	Total	O	0	0
			117	117		
7	D	76	Total	O	0	0
			76	76		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cytochrome P450 CYP260A1,Cytochrome P450 CYP260A1




- Molecule 1: Cytochrome P450 CYP260A1,Cytochrome P450 CYP260A1




- Molecule 1: Cytochrome P450 CYP260A1,Cytochrome P450 CYP260A1





● Molecule 1: Cytochrome P450 CYP260A1,Cytochrome P450 CYP260A1



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	234.56 Å 234.56 Å 96.43 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	69.93 – 2.67 69.93 – 2.67	Depositor EDS
% Data completeness (in resolution range)	99.7 (69.93-2.67) 99.7 (69.93-2.67)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 2.65 Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.187 , 0.223 0.193 , 0.226	Depositor DCC
R_{free} test set	4262 reflections (5.21%)	DCC
Wilson B-factor (Å ²)	40.5	Xtriage
Anisotropy	0.157	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12879	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, DMS, SO4, MES, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.83	1/3161 (0.0%)	0.95	6/4284 (0.1%)
1	B	0.79	1/3068 (0.0%)	0.97	9/4164 (0.2%)
1	C	0.94	2/3157 (0.1%)	0.98	10/4281 (0.2%)
1	D	0.85	0/3058	0.99	8/4146 (0.2%)
All	All	0.85	4/12444 (0.0%)	0.97	33/16875 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
1	D	0	2
All	All	0	4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	328	GLY	N-CA	-7.92	1.34	1.46
1	C	361	GLU	CG-CD	6.98	1.62	1.51
1	A	113	MET	CG-SD	-5.87	1.65	1.81
1	C	345	SER	CB-OG	-5.80	1.34	1.42

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	104	ARG	NE-CZ-NH1	-9.16	115.72	120.30
1	D	208	MET	CG-SD-CE	-8.16	87.14	100.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	104	ARG	NE-CZ-NH2	7.84	124.22	120.30
1	C	72	ARG	CG-CD-NE	7.17	126.86	111.80
1	A	113	MET	CB-CG-SD	6.90	133.09	112.40

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	418	PRO	Peptide
1	C	168	GLY	Peptide
1	D	168	GLY	Peptide
1	D	242	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3092	0	3071	41	0
1	B	3003	0	2977	41	1
1	C	3091	0	3085	48	1
1	D	2997	0	2971	50	0
2	A	43	0	30	4	0
2	B	43	0	30	5	0
2	C	43	0	30	5	0
2	D	43	0	30	6	0
3	A	24	0	32	0	0
3	B	6	0	8	0	0
3	C	18	0	24	9	0
3	D	18	0	24	6	0
4	A	4	0	6	0	0
4	B	4	0	6	1	0
4	C	4	0	6	0	0
4	D	12	0	18	0	0
5	A	20	0	0	0	0
5	B	20	0	0	1	0
5	C	20	0	0	1	0
5	D	25	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	12	0	13	0	0
7	A	78	0	0	1	0
7	B	66	0	0	2	0
7	C	117	0	0	2	0
7	D	76	0	0	4	0
All	All	12879	0	12361	199	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 199 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:65:ASP:OD2	1:C:67:THR:HB	1.43	1.17
1:D:60:PHE:HA	1:D:425:MET:HE2	1.45	0.95
1:A:104:ARG:HD2	3:D:507:GOL:H2	1.49	0.92
1:D:145:ARG:HD2	1:D:145:ARG:H	1.35	0.92
1:A:445:HIS:O	1:A:446:HIS:CG	2.28	0.87

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:225:GLN:NE2	1:C:423:SER:OG[3_455]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	395/450 (88%)	375 (95%)	18 (5%)	2 (0%)	34 61
1	B	384/450 (85%)	365 (95%)	15 (4%)	4 (1%)	19 42

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	C	395/450 (88%)	379 (96%)	15 (4%)	1 (0%)	46 73
1	D	383/450 (85%)	370 (97%)	12 (3%)	1 (0%)	46 73
All	All	1557/1800 (86%)	1489 (96%)	60 (4%)	8 (0%)	34 61

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	245	ASP
1	B	218	PRO
1	C	52	ASP
1	D	244	ARG
1	A	445	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	316/361 (88%)	308 (98%)	8 (2%)	55 83
1	B	307/361 (85%)	299 (97%)	8 (3%)	54 81
1	C	320/361 (89%)	307 (96%)	13 (4%)	37 66
1	D	305/361 (84%)	295 (97%)	10 (3%)	45 74
All	All	1248/1444 (86%)	1209 (97%)	39 (3%)	47 76

5 of 39 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	67	THR
1	C	200	GLU
1	D	245	ASP
1	C	100	THR
1	C	113	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	149	GLN
1	B	307	GLN
1	C	149	GLN
1	A	358	ASN
1	C	78	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

39 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	501	1	24,50,50	1.33	4 (16%)	16,82,82	2.19	5 (31%)
3	GOL	A	502	-	5,5,5	0.68	0	5,5,5	1.31	0
3	GOL	A	503	-	5,5,5	0.72	0	5,5,5	0.45	0
3	GOL	A	504	-	5,5,5	0.96	0	5,5,5	0.76	0
4	DMS	A	505	-	3,3,3	0.22	0	3,3,3	1.14	0
3	GOL	A	506	-	5,5,5	1.05	0	5,5,5	1.21	0
5	SO4	A	507	-	4,4,4	0.98	0	6,6,6	0.54	0
5	SO4	A	508	-	4,4,4	0.41	0	6,6,6	0.88	0
6	MES	A	509	-	12,12,12	2.60	2 (16%)	15,16,16	1.71	4 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	A	510	-	4,4,4	0.32	0	6,6,6	0.49	0
5	SO4	A	511	-	4,4,4	0.76	0	6,6,6	0.86	0
2	HEM	B	501	1	24,50,50	1.53	4 (16%)	16,82,82	1.82	5 (31%)
5	SO4	B	502	-	4,4,4	0.84	0	6,6,6	0.46	0
5	SO4	B	503	-	4,4,4	0.68	0	6,6,6	0.45	0
3	GOL	B	504	-	5,5,5	0.41	0	5,5,5	1.40	1 (20%)
4	DMS	B	505	-	3,3,3	0.47	0	3,3,3	1.59	1 (33%)
5	SO4	B	506	-	4,4,4	0.71	0	6,6,6	0.60	0
5	SO4	B	507	-	4,4,4	0.59	0	6,6,6	0.14	0
2	HEM	C	501	1	24,50,50	2.02	8 (33%)	16,82,82	2.19	8 (50%)
5	SO4	C	502	-	4,4,4	0.55	0	6,6,6	0.19	0
5	SO4	C	503	-	4,4,4	0.84	0	6,6,6	0.44	0
3	GOL	C	504	-	5,5,5	0.65	0	5,5,5	1.57	0
3	GOL	C	505	-	5,5,5	0.51	0	5,5,5	0.92	0
3	GOL	C	506	-	5,5,5	0.64	0	5,5,5	1.76	2 (40%)
4	DMS	C	507	-	3,3,3	0.41	0	3,3,3	1.36	1 (33%)
5	SO4	C	508	-	4,4,4	0.94	0	6,6,6	0.92	0
5	SO4	C	509	-	4,4,4	0.56	0	6,6,6	0.23	0
2	HEM	D	501	1	24,50,50	1.64	5 (20%)	16,82,82	1.93	6 (37%)
3	GOL	D	502	-	5,5,5	0.69	0	5,5,5	1.36	0
4	DMS	D	503	-	3,3,3	0.34	0	3,3,3	0.57	0
4	DMS	D	504	-	3,3,3	0.25	0	3,3,3	1.36	1 (33%)
3	GOL	D	505	-	5,5,5	0.59	0	5,5,5	0.65	0
4	DMS	D	506	-	3,3,3	0.44	0	3,3,3	0.64	0
3	GOL	D	507	-	5,5,5	1.13	0	5,5,5	1.48	1 (20%)
5	SO4	D	508	-	4,4,4	0.77	0	6,6,6	0.42	0
5	SO4	D	509	-	4,4,4	0.69	0	6,6,6	0.38	0
5	SO4	D	510	-	4,4,4	1.13	0	6,6,6	0.60	0
5	SO4	D	511	-	4,4,4	0.56	0	6,6,6	0.42	0
5	SO4	D	512	-	4,4,4	0.50	0	6,6,6	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	501	1	-	0/6/54/54	0/0/8/8
3	GOL	A	502	-	-	0/4/4/4	0/0/0/0
3	GOL	A	503	-	-	0/4/4/4	0/0/0/0
3	GOL	A	504	-	-	0/4/4/4	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DMS	A	505	-	-	0/0/0/0	0/0/0/0
3	GOL	A	506	-	-	0/4/4/4	0/0/0/0
5	SO4	A	507	-	-	0/0/0/0	0/0/0/0
5	SO4	A	508	-	-	0/0/0/0	0/0/0/0
6	MES	A	509	-	-	0/6/14/14	0/1/1/1
5	SO4	A	510	-	-	0/0/0/0	0/0/0/0
5	SO4	A	511	-	-	0/0/0/0	0/0/0/0
2	HEM	B	501	1	-	0/6/54/54	0/0/8/8
5	SO4	B	502	-	-	0/0/0/0	0/0/0/0
5	SO4	B	503	-	-	0/0/0/0	0/0/0/0
3	GOL	B	504	-	-	0/4/4/4	0/0/0/0
4	DMS	B	505	-	-	0/0/0/0	0/0/0/0
5	SO4	B	506	-	-	0/0/0/0	0/0/0/0
5	SO4	B	507	-	-	0/0/0/0	0/0/0/0
2	HEM	C	501	1	-	0/6/54/54	0/0/8/8
5	SO4	C	502	-	-	0/0/0/0	0/0/0/0
5	SO4	C	503	-	-	0/0/0/0	0/0/0/0
3	GOL	C	504	-	-	0/4/4/4	0/0/0/0
3	GOL	C	505	-	-	0/4/4/4	0/0/0/0
3	GOL	C	506	-	-	0/4/4/4	0/0/0/0
4	DMS	C	507	-	-	0/0/0/0	0/0/0/0
5	SO4	C	508	-	-	0/0/0/0	0/0/0/0
5	SO4	C	509	-	-	0/0/0/0	0/0/0/0
2	HEM	D	501	1	-	0/6/54/54	0/0/8/8
3	GOL	D	502	-	-	0/4/4/4	0/0/0/0
4	DMS	D	503	-	-	0/0/0/0	0/0/0/0
4	DMS	D	504	-	-	0/0/0/0	0/0/0/0
3	GOL	D	505	-	-	0/4/4/4	0/0/0/0
4	DMS	D	506	-	-	0/0/0/0	0/0/0/0
3	GOL	D	507	-	-	0/4/4/4	0/0/0/0
5	SO4	D	508	-	-	0/0/0/0	0/0/0/0
5	SO4	D	509	-	-	0/0/0/0	0/0/0/0
5	SO4	D	510	-	-	0/0/0/0	0/0/0/0
5	SO4	D	511	-	-	0/0/0/0	0/0/0/0
5	SO4	D	512	-	-	0/0/0/0	0/0/0/0

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	509	MES	C8-S	-8.34	1.65	1.77
2	C	501	HEM	C3B-C2B	-4.26	1.35	1.40
2	C	501	HEM	C3C-C2C	-4.01	1.35	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	HEM	C1B-NB	-3.90	1.31	1.36
2	C	501	HEM	C3B-CAB	-3.62	1.40	1.47

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	HEM	C3B-C4B-NB	-4.31	103.64	109.21
2	C	501	HEM	C3B-CAB-CBB	-4.06	118.24	126.40
2	A	501	HEM	CBA-CAA-C2A	-3.82	105.77	112.49
2	C	501	HEM	CBA-CAA-C2A	-3.77	105.86	112.49
2	B	501	HEM	C3B-C4B-NB	-3.74	104.37	109.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

13 monomers are involved in 38 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	HEM	4	0
2	B	501	HEM	5	0
4	B	505	DMS	1	0
5	B	506	SO4	1	0
2	C	501	HEM	5	0
5	C	502	SO4	1	0
3	C	504	GOL	2	0
3	C	505	GOL	7	0
3	C	506	GOL	6	0
2	D	501	HEM	6	0
3	D	502	GOL	3	0
3	D	505	GOL	3	0
3	D	507	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	397/450 (88%)	0.06	22 (5%) 29 26	17, 42, 87, 111	0
1	B	389/450 (86%)	0.25	41 (10%) 8 6	23, 46, 93, 126	0
1	C	395/450 (87%)	-0.22	5 (1%) 79 79	17, 30, 59, 97	0
1	D	389/450 (86%)	-0.04	13 (3%) 50 49	18, 38, 83, 107	0
All	All	1570/1800 (87%)	0.01	81 (5%) 31 29	17, 38, 86, 126	0

The worst 5 of 81 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	239	VAL	5.3
1	B	194	PRO	5.2
1	B	264	ASP	4.6
1	A	244	ARG	4.5
1	B	238	VAL	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	D	507	6/6	0.83	0.28	5.77	19,32,38,43	0
5	SO4	D	510	5/5	0.87	0.23	5.33	75,75,91,100	0
5	SO4	C	503	5/5	0.80	0.27	4.85	65,93,107,111	0
3	GOL	B	504	6/6	0.93	0.20	3.96	40,42,45,50	0
3	GOL	C	504	6/6	0.89	0.22	2.80	34,42,46,47	0
3	GOL	A	502	6/6	0.86	0.24	2.52	53,60,64,66	0
3	GOL	D	502	6/6	0.90	0.21	1.95	29,49,56,66	0
5	SO4	A	508	5/5	0.95	0.34	1.38	69,82,86,87	0
3	GOL	A	506	6/6	0.86	0.19	1.33	43,48,56,60	0
3	GOL	A	503	6/6	0.89	0.17	0.44	33,45,53,56	0
2	HEM	D	501	43/43	0.98	0.16	0.21	18,23,30,40	0
2	HEM	A	501	43/43	0.98	0.14	-0.34	22,25,35,38	0
2	HEM	B	501	43/43	0.98	0.14	-0.38	19,22,29,34	0
4	DMS	D	504	4/4	0.87	0.16	-0.56	51,54,56,61	0
2	HEM	C	501	43/43	0.98	0.14	-0.64	15,19,21,25	0
5	SO4	B	503	5/5	0.98	0.08	-1.60	50,51,58,63	0
4	DMS	B	505	4/4	0.95	0.17	-	50,53,58,60	0
4	DMS	A	505	4/4	0.95	0.13	-	48,54,55,60	0
5	SO4	A	507	5/5	0.97	0.13	-	47,48,58,61	0
3	GOL	A	504	6/6	0.90	0.20	-	35,51,59,60	0
3	GOL	D	505	6/6	0.91	0.27	-	57,62,67,74	0
5	SO4	D	512	5/5	0.95	0.23	-	83,84,85,92	0
5	SO4	B	507	5/5	0.94	0.15	-	68,74,82,87	0
4	DMS	D	506	4/4	0.94	0.14	-	90,92,102,107	0
5	SO4	C	508	5/5	0.93	0.20	-	67,69,76,76	0
4	DMS	C	507	4/4	0.94	0.18	-	48,57,58,62	0
5	SO4	D	511	5/5	0.90	0.32	-	71,97,101,105	0
5	SO4	A	510	5/5	0.96	0.16	-	59,61,64,65	5
5	SO4	B	506	5/5	0.95	0.23	-	60,65,75,75	0
5	SO4	D	508	5/5	0.92	0.28	-	71,76,81,85	0
5	SO4	B	502	5/5	0.94	0.19	-	56,73,81,82	0
3	GOL	C	506	6/6	0.95	0.19	-	36,37,41,48	0
5	SO4	D	509	5/5	0.91	0.17	-	62,70,84,87	0
5	SO4	C	502	5/5	0.93	0.23	-	61,79,86,94	0
5	SO4	A	511	5/5	0.89	0.28	-	66,68,85,95	0
5	SO4	C	509	5/5	0.93	0.34	-	85,87,95,96	0
3	GOL	C	505	6/6	0.96	0.16	-	31,32,34,36	0
6	MES	A	509	12/12	0.97	0.17	-	39,49,54,55	0
4	DMS	D	503	4/4	0.96	0.14	-	52,66,69,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.