



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 19, 2016 – 01:25 PM EST

PDB ID : 5LKU
Title : Crystal structure of the p300 acetyltransferase catalytic core with coenzyme A.
Authors : Kaczmarska, Z.; Ortega, E.; Marquez, J.A.; Panne, D.
Deposited on : 2016-07-25
Resolution : 3.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028442
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1051 (3.60-3.40)
Clashscore	102246	1157 (3.60-3.40)
Ramachandran outliers	100387	1120 (3.60-3.40)
Sidechain outliers	100360	1121 (3.60-3.40)
RSRZ outliers	91569	1058 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	578	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; left: 0; top: -10px; width: 5px; height: 5px; background-color: red;"></div> <div style="position: absolute; left: 86%; top: -10px; width: 5px; height: 5px; background-color: yellow;"></div> <div style="position: absolute; left: 99%; top: -10px; width: 5px; height: 5px; background-color: grey;"></div> </div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 86% 8% 5% </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4580 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone acetyltransferase p300,Histone acetyltransferase p300.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	551	4512	2884	769	826	33	0	0	0


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1033	GLY	-	expression tag	UNP Q09472
A	1034	ALA	-	expression tag	UNP Q09472
A	1035	MET	-	expression tag	UNP Q09472
A	1036	ALA	-	expression tag	UNP Q09472
A	1037	GLY	-	expression tag	UNP Q09472
A	1038	LYS	-	expression tag	UNP Q09472
A	1039	ALA	-	expression tag	UNP Q09472
A	1040	VAL	-	expression tag	UNP Q09472
A	1041	PRO	-	expression tag	UNP Q09472
A	1042	MET	-	expression tag	UNP Q09472
A	1467	PHE	TYR	engineered mutation	UNP Q09472
A	1520	SER	-	linker	UNP Q09472
A	1521	GLY	-	linker	UNP Q09472
A	1522	GLY	-	linker	UNP Q09472
A	1523	SER	-	linker	UNP Q09472
A	1580	GLY	-	linker	UNP Q09472

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Zn	0	0
			4	4		

- Molecule 3 is COENZYME A (three-letter code: COA) (formula: C₂₁H₃₆N₇O₁₆P₃S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	48	21	7	16	3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	16	Total O 16 16	0	0

- Molecule 1: Histone acetyltransferase p300, Histone acetyltransferase p300



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	93.96 Å 156.40 Å 110.68 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.01 – 3.50 29.68 – 3.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.01-3.50) 99.9 (29.68-3.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 3.47 Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.200 , 0.242 0.203 , 0.234	Depositor DCC
R_{free} test set	539 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	63.2	Xtriage
Anisotropy	0.049	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 67.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.018 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.038 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4580	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: COA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/4633	0.71	3/6263 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	1297	ARG	NE-CZ-NH1	10.14	125.37	120.30
1	A	1297	ARG	NE-CZ-NH2	-5.38	117.61	120.30
1	A	1305	ARG	NE-CZ-NH2	-5.37	117.61	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4512	0	4409	30	0
2	A	4	0	0	0	1
3	A	48	0	32	0	0
4	A	16	0	0	0	0
All	All	4580	0	4441	30	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1366:ILE:HD12	1:A:1371:LEU:HD23	1.75	0.67
1:A:1060:PRO:HA	1:A:1063:GLU:HG2	1.79	0.63
1:A:1172:PRO:HG3	1:A:1187:ARG:HD3	1.81	0.60
1:A:1172:PRO:CG	1:A:1187:ARG:HD3	2.32	0.60
1:A:1105:LYS:HG2	1:A:1110:GLN:OE1	2.04	0.57
1:A:1235:LYS:N	1:A:1235:LYS:HD2	2.21	0.55
1:A:1198:TYR:CD2	1:A:1215:LEU:HD13	2.44	0.52
1:A:1359:ALA:HB2	1:A:1377:HIS:CD2	2.45	0.52
1:A:1402:HIS:HD2	1:A:1455:GLN:HB2	1.75	0.52
1:A:1297:ARG:HH11	1:A:1297:ARG:CG	2.23	0.52
1:A:1366:ILE:CD1	1:A:1371:LEU:HD23	2.44	0.48
1:A:1071:GLU:HA	1:A:1141:TYR:CD2	2.49	0.48
1:A:1052:GLU:O	1:A:1056:GLN:HG2	2.13	0.47
1:A:1353:PHE:CD2	1:A:1386:PRO:HG2	2.50	0.47
1:A:1355:TYR:HB3	1:A:1381:TYR:CE2	2.50	0.46
1:A:1436:TRP:CD1	1:A:1595:PHE:CE1	3.05	0.45
1:A:1415:HIS:HE1	1:A:1466:TRP:CE2	2.34	0.45
1:A:1086:ILE:HG23	1:A:1131:TYR:OH	2.17	0.45
1:A:1649:TRP:CZ3	1:A:1652:MET:CE	3.00	0.44
1:A:1177:CYS:SG	1:A:1204:CYS:SG	3.16	0.44
1:A:1121:ILE:HG22	1:A:1125:PHE:CE2	2.53	0.43
1:A:1402:HIS:CD2	1:A:1455:GLN:HB2	2.53	0.43
1:A:1459:LYS:HB3	1:A:1460:PRO:CD	2.50	0.42
1:A:1651:THR:HA	1:A:1654:MET:HE3	2.01	0.42
1:A:1523:SER:OG	1:A:1581:SER:N	2.52	0.42
1:A:1122:TRP:CD2	1:A:1167:LYS:HD2	2.54	0.41
1:A:1122:TRP:CG	1:A:1167:LYS:HD2	2.55	0.41
1:A:1172:PRO:HG2	1:A:1187:ARG:HD3	2.00	0.41
1:A:1436:TRP:CD1	1:A:1594:VAL:HG12	2.56	0.40
1:A:1329:SER:HA	1:A:1618:LEU:HD23	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1704:ZN:ZN	2:A:1704:ZN:ZN[3_554]	0.55	1.65

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	545/578 (94%)	530 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	501/522 (96%)	494 (99%)	7 (1%)	74	91

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1167	LYS
1	A	1263	GLU
1	A	1291	LYS
1	A	1297	ARG
1	A	1396	SER
1	A	1397	TYR
1	A	1415	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1256	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1402	HIS
1	A	1415	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	COA	A	1705	-	41,50,50	1.58	6 (14%)	49,75,75	2.46	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	COA	A	1705	-	-	0/44/64/64	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1705	COA	C5A-C4A	-3.10	1.33	1.40
3	A	1705	COA	C2B-C1B	-2.63	1.49	1.53
3	A	1705	COA	O4B-C1B	2.39	1.44	1.41
3	A	1705	COA	P3B-O9A	3.05	1.65	1.54
3	A	1705	COA	C2A-N1A	3.97	1.41	1.33
3	A	1705	COA	C2A-N3A	4.73	1.40	1.32

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1705	COA	N3A-C2A-N1A	-14.21	117.71	128.87
3	A	1705	COA	O4B-C1B-N9A	-5.34	98.01	108.11
3	A	1705	COA	C1B-N9A-C4A	-4.40	121.90	126.81
3	A	1705	COA	C2P-C3P-N4P	-2.55	107.39	112.44
3	A	1705	COA	CAP-C9P-N8P	-2.16	111.22	116.31
3	A	1705	COA	C6P-C5P-N4P	-2.09	112.82	116.46
3	A	1705	COA	O3B-P3B-O7A	2.87	114.33	107.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	551/578 (95%)	-0.30	7 (1%) 79 70	28, 60, 116, 161	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1209	GLN	3.0
1	A	1211	GLU	3.0
1	A	1084	LEU	2.9
1	A	1210	GLY	2.9
1	A	1107	ASP	2.6
1	A	1181	GLN	2.3
1	A	1182	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	COA	A	1705	48/48	0.94	0.17	-0.91	29,43,53,54	0
2	ZN	A	1702	1/1	0.99	0.05	-1.96	70,70,70,70	0
2	ZN	A	1701	1/1	0.95	0.09	-	142,142,142,142	0
2	ZN	A	1704	1/1	0.98	0.04	-	95,95,95,95	0
2	ZN	A	1703	1/1	0.97	0.12	-	59,59,59,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.