



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:18 AM GMT

PDB ID : 3LMB
Title : The crystal structure of the protein OLEI01261 with unknown function from Chlorobaculum tepidum TLS
Authors : Zhang, R.; Evdokimova, E.; Egorova, O.; Savchenko, A.; Edwards, A.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2010-01-29
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	165	 6% 70% 23% . .
1	B	165	 16% 77% 19% ...

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2709 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	164	Total	C	N	O	S	Se	0	0	0
			1289	820	213	246	4	6			
1	B	163	Total	C	N	O	S	Se	0	2	0
			1286	819	213	244	4	6			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	81	Total	O	0	0
			81	81		
2	B	53	Total	O	0	0
			53	53		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uncharacterized protein



• Molecule 1: Uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.09Å 72.96Å 79.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.83 – 2.10 32.88 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (53.83-2.10) 99.9 (32.88-2.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.166 , 0.222 0.173 , 0.220	Depositor DCC
R_{free} test set	1043 reflections (5.40%)	DCC
Wilson B-factor (Å ²)	30.8	Xtriage
Anisotropy	0.188	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 54.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 20380 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2709	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.20	4/1311 (0.3%)	1.03	9/1764 (0.5%)
1	B	0.98	0/1308	0.89	2/1761 (0.1%)
All	All	1.09	4/2619 (0.2%)	0.96	11/3525 (0.3%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	153	VAL	CB-CG1	-8.67	1.34	1.52
1	A	2	ASN	CB-CG	8.21	1.70	1.51
1	A	10	VAL	CB-CG1	-7.09	1.38	1.52
1	A	79	GLU	CB-CG	-5.17	1.42	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	88	VAL	CB-CA-C	-8.18	95.86	111.40
1	A	132	LEU	CA-CB-CG	8.13	134.01	115.30
1	A	88	VAL	CB-CA-C	-7.80	96.59	111.40
1	A	102	ARG	NE-CZ-NH1	6.83	123.71	120.30
1	A	43	THR	CB-CA-C	-6.37	94.40	111.60
1	A	89	THR	N-CA-CB	-6.31	98.32	110.30
1	B	43	THR	CB-CA-C	-6.27	94.67	111.60
1	A	139	ASP	CB-CG-OD1	6.08	123.78	118.30
1	A	88	VAL	CG1-CB-CG2	5.95	120.42	110.90
1	A	102	ARG	NE-CZ-NH2	-5.80	117.40	120.30
1	A	8	ASP	CB-CA-C	-5.21	99.98	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1289	0	1270	33	0
1	B	1286	0	1248	38	0
2	A	81	0	0	6	0
2	B	53	0	0	1	0
All	All	2709	0	2518	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:THR:CG2	1:A:158:GLN:HE21	1.46	1.28
1:B:18:PHE:CE1	1:B:67[B]:MSE:HE2	1.91	1.05
1:A:89:THR:CG2	1:A:158:GLN:NE2	2.21	1.01
1:B:18:PHE:HE1	1:B:67[B]:MSE:HE2	1.28	0.94
1:B:108:HIS:HE1	1:B:145:ILE:H	0.97	0.94
1:A:89:THR:HG22	1:A:158:GLN:HE21	1.30	0.94
1:B:13:LYS:HE2	1:B:78:GLN:NE2	1.86	0.89
1:B:108:HIS:CE1	1:B:145:ILE:H	1.88	0.88
1:B:6:THR:HG22	1:B:9:GLN:H	1.38	0.86
1:A:89:THR:HG21	1:A:158:GLN:NE2	1.90	0.84
1:B:18:PHE:HE1	1:B:67[B]:MSE:CE	1.89	0.83
1:B:18:PHE:CE1	1:B:67[B]:MSE:CE	2.64	0.75
1:A:146:GLU:HB2	1:A:148:GLU:OE2	1.87	0.74
1:B:108:HIS:HE1	1:B:145:ILE:N	1.80	0.72
1:A:59:GLY:HA3	1:B:63:ASN:HD21	1.54	0.72
1:B:5:LEU:HB3	2:B:190:HOH:O	1.92	0.68
1:A:28:MSE:HG2	1:A:49:ASN:HD21	1.58	0.67
1:B:55:THR:HG22	1:B:100[B]:TYR:CD1	2.30	0.67
1:B:30:LEU:HD22	1:B:67[A]:MSE:HE3	1.76	0.66
1:A:80:GLU:HB3	1:A:82:ILE:HD12	1.78	0.66
1:A:100:TYR:H	1:A:138:ASN:HD21	1.44	0.64
1:A:77:THR:CG2	2:A:230:HOH:O	2.50	0.60
1:B:13:LYS:HE2	1:B:78:GLN:HE22	1.62	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:43:THR:HG23	1:B:104:ARG:HG2	1.85	0.58
1:A:164:ASN:HB2	2:A:229:HOH:O	2.04	0.58
1:A:100:TYR:H	1:A:138:ASN:ND2	2.03	0.56
1:B:6:THR:CG2	1:B:9:GLN:H	2.14	0.56
1:B:98:PRO:HG2	1:B:100[B]:TYR:CZ	2.40	0.56
1:A:88:VAL:HG13	1:A:159:TYR:CE1	2.42	0.55
1:A:43:THR:HG21	1:A:102:ARG:HD3	1.89	0.54
1:A:99:VAL:HA	1:A:138:ASN:HD21	1.73	0.54
1:B:67[B]:MSE:HE1	1:B:70:TRP:CE3	2.42	0.54
1:A:43:THR:HG21	2:A:168:HOH:O	2.07	0.53
1:A:126:GLY:HA2	1:A:163:LYS:HE3	1.91	0.52
1:A:43:THR:HG23	1:A:104:ARG:HG2	1.91	0.51
1:A:59:GLY:HA3	1:B:63:ASN:ND2	2.23	0.51
1:A:77:THR:HG21	2:A:167:HOH:O	2.10	0.51
1:A:150:LYS:H	1:A:150:LYS:HD2	1.75	0.51
1:A:51:ASN:HA	1:B:22:LEU:HD23	1.92	0.51
1:B:6:THR:HG22	1:B:9:GLN:N	2.19	0.50
1:B:5:LEU:H	1:B:9:GLN:HE21	1.60	0.49
1:B:30:LEU:HD22	1:B:67[A]:MSE:CE	2.40	0.49
1:B:61:LEU:HD13	1:B:153:VAL:HG21	1.95	0.48
1:A:97:ALA:HB1	1:A:98:PRO:HD2	1.95	0.47
1:B:2:ASN:N	1:B:2:ASN:OD1	2.47	0.47
1:A:77:THR:HG23	2:A:230:HOH:O	2.10	0.47
1:B:82:ILE:HD12	1:B:119:PHE:CD1	2.49	0.47
1:A:89:THR:HG23	1:A:158:GLN:NE2	2.26	0.47
1:A:12:LYS:NZ	2:A:220:HOH:O	2.47	0.47
1:B:88:VAL:HG13	1:B:159:TYR:CE1	2.51	0.46
1:B:93:MSE:HE2	1:B:153:VAL:HG12	1.98	0.46
1:B:5:LEU:H	1:B:9:GLN:NE2	2.14	0.45
1:B:100[A]:TYR:CZ	1:B:138:ASN:ND2	2.85	0.44
1:B:93:MSE:HE2	1:B:153:VAL:CG1	2.48	0.44
1:A:121:HIS:HD2	1:A:127:LYS:O	2.00	0.44
1:B:30:LEU:CD2	1:B:67[A]:MSE:HE3	2.48	0.42
1:B:55:THR:HG22	1:B:100[A]:TYR:HB3	2.01	0.42
1:A:63:ASN:HB3	1:A:67:MSE:HE2	2.01	0.42
1:A:124:ARG:HH21	1:A:125:LYS:HE3	1.84	0.42
1:A:16:GLN:NE2	1:A:20:ASP:OD1	2.50	0.42
1:B:6:THR:HG22	1:B:9:GLN:HB2	2.01	0.41
1:A:24:ILE:HG21	1:B:57:PHE:CD2	2.55	0.41
1:A:53:LYS:O	1:A:54:GLN:HB2	2.19	0.41
1:A:107:CYS:SG	1:A:132:LEU:HD12	2.60	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:55:THR:HG22	1:B:100[B]:TYR:HD1	1.81	0.41
1:B:43:THR:HG21	1:B:102:ARG:HD3	2.03	0.40
1:B:142:VAL:CG1	1:B:144:LYS:O	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	162/165 (98%)	158 (98%)	4 (2%)	0	100	100
1	B	163/165 (99%)	159 (98%)	3 (2%)	1 (1%)	30	24
All	All	325/330 (98%)	317 (98%)	7 (2%)	1 (0%)	46	45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	148	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/134 (104%)	125 (90%)	14 (10%)	9	5
1	B	136/134 (102%)	127 (93%)	9 (7%)	21	17

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	275/268 (103%)	252 (92%)	23 (8%)	14 9

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	A	10	VAL
1	A	43	THR
1	A	77	THR
1	A	80	GLU
1	A	88	VAL
1	A	89	THR
1	A	123	GLU
1	A	132	LEU
1	A	144	LYS
1	A	148	GLU
1	A	149	THR
1	A	150	LYS
1	A	163	LYS
1	B	2	ASN
1	B	5	LEU
1	B	6	THR
1	B	43	THR
1	B	88	VAL
1	B	94	LYS
1	B	112	GLU
1	B	123	GLU
1	B	149	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	49	ASN
1	A	78	GLN
1	A	81	ASN
1	A	121	HIS
1	A	138	ASN
1	A	158	GLN
1	B	2	ASN
1	B	9	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	16	GLN
1	B	63	ASN
1	B	78	GLN
1	B	81	ASN
1	B	85	ASN
1	B	108	HIS
1	B	121	HIS
1	B	164	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	158/165 (95%)	0.30	10 (6%) 23 31	8, 18, 39, 51	0
1	B	158/165 (95%)	0.66	26 (16%) 2 3	12, 22, 43, 49	0
All	All	316/330 (95%)	0.48	36 (11%) 7 9	8, 20, 41, 51	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	123	GLU	5.2
1	B	3	ALA	4.9
1	A	148	GLU	4.8
1	B	124	ARG	4.8
1	B	149	THR	4.8
1	B	147	PRO	4.7
1	B	83	ALA	4.5
1	B	145	ILE	4.3
1	B	148	GLU	4.1
1	B	125	LYS	3.7
1	B	163	LYS	3.6
1	B	127	LYS	3.3
1	A	120	ASP	3.1
1	B	120	ASP	3.1
1	A	147	PRO	3.1
1	B	121	HIS	3.0
1	A	123	GLU	2.9
1	A	149	THR	2.9
1	B	62	TYR	2.8
1	B	162	LEU	2.8
1	B	111	ASP	2.8
1	A	113	GLU	2.7
1	B	66	VAL	2.7
1	B	164	ASN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	62	TYR	2.7
1	B	113	GLU	2.6
1	A	164	ASN	2.5
1	A	66	VAL	2.5
1	B	4	SER	2.4
1	B	122	PHE	2.4
1	A	61	LEU	2.3
1	B	2	ASN	2.2
1	B	142	VAL	2.2
1	B	117	ASN	2.1
1	B	146	GLU	2.0
1	B	150	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.