



Full wwPDB NMR Structure Validation Report ⓘ

Apr 27, 2016 – 02:20 AM BST

PDB ID : 2LQ6
Title : Solution structure of BRD1 PHD2 finger
Authors : Liu, L.; Wu, J.
Deposited on : 2012-02-25

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 85%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	87	<div> <div></div> <div>57%</div> <div>18%</div> <div>15%</div> <div>9%</div> </div>

2 Ensemble composition and analysis ⓘ

This entry contains 20 models. Model 3 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *minimized average structure*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:325-A:390 (66)	0.86	3

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

NmrClust was unable to cluster the ensemble.

Error message: Inconsistent models

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1201 atoms, of which 599 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Bromodomain-containing protein 1.

Mol	Chain	Residues	Atoms						Trace
1	A	79	Total	C	H	N	O	S	0
			1199	379	599	105	107	9	

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	308	MET	-	EXPRESSION TAG	UNP O95696
A	309	GLY	-	EXPRESSION TAG	UNP O95696
A	310	HIS	-	EXPRESSION TAG	UNP O95696
A	311	HIS	-	EXPRESSION TAG	UNP O95696
A	312	HIS	-	EXPRESSION TAG	UNP O95696
A	313	HIS	-	EXPRESSION TAG	UNP O95696
A	314	HIS	-	EXPRESSION TAG	UNP O95696
A	315	HIS	-	EXPRESSION TAG	UNP O95696
A	316	MET	-	EXPRESSION TAG	UNP O95696
A	338	SER	CYS	ENGINEERED MUTATION	UNP O95696
A	393	SER	CYS	ENGINEERED MUTATION	UNP O95696

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
2	A	2	Total	Zn
			2	2

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Bromodomain-containing protein 1




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Bromodomain-containing protein 1




4.2.2 Score per residue for model 2

- Molecule 1: Bromodomain-containing protein 1




4.2.3 Score per residue for model 3 (medoid)

- Molecule 1: Bromodomain-containing protein 1




4.2.4 Score per residue for model 4

- Molecule 1: Bromodomain-containing protein 1




4.2.5 Score per residue for model 5

- Molecule 1: Bromodomain-containing protein 1




4.2.6 Score per residue for model 6

- Molecule 1: Bromodomain-containing protein 1




4.2.7 Score per residue for model 7

- Molecule 1: Bromodomain-containing protein 1




4.2.8 Score per residue for model 8

- Molecule 1: Bromodomain-containing protein 1




4.2.9 Score per residue for model 9

- Molecule 1: Bromodomain-containing protein 1




4.2.10 Score per residue for model 10

- Molecule 1: Bromodomain-containing protein 1




4.2.11 Score per residue for model 11

- Molecule 1: Bromodomain-containing protein 1




4.2.12 Score per residue for model 12

- Molecule 1: Bromodomain-containing protein 1




4.2.13 Score per residue for model 13

- Molecule 1: Bromodomain-containing protein 1




4.2.14 Score per residue for model 14

- Molecule 1: Bromodomain-containing protein 1




4.2.15 Score per residue for model 15

- Molecule 1: Bromodomain-containing protein 1




4.2.16 Score per residue for model 16

- Molecule 1: Bromodomain-containing protein 1



4.2.17 Score per residue for model 17

- Molecule 1: Bromodomain-containing protein 1



5 Refinement protocol and experimental data overview

The models were refined using the following method: *DGSA-distance geometry simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	1.2
CNS	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2lq6_cs.str
Number of chemical shift lists	1
Total number of shifts	887
Number of shifts mapped to atoms	887
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	85%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	498	496	495	7±2
All	All	10000	9920	9891	134

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:385:CYS:SG	1:A:388:HIS:HB2	0.85	2.12	1	19
1:A:351:HIS:HB2	1:A:354:CYS:SG	0.80	2.16	3	18
1:A:342:HIS:HB3	1:A:384:TYR:CE1	0.74	2.18	18	6
1:A:342:HIS:HB3	1:A:384:TYR:CE2	0.71	2.21	20	1
1:A:389:THR:HB	1:A:390:PRO:HD3	0.63	1.69	8	3
1:A:369:GLU:HB2	1:A:376:THR:OG1	0.63	1.93	2	7
1:A:369:GLU:O	1:A:375:THR:HA	0.60	1.96	7	3
1:A:369:GLU:HB3	1:A:376:THR:OG1	0.60	1.96	19	4
1:A:389:THR:HB	1:A:390:PRO:CD	0.58	2.28	8	2
1:A:327:CYS:HB3	1:A:332:GLN:O	0.57	1.99	19	1
1:A:370:LEU:HD12	1:A:374:GLY:O	0.56	1.99	19	2
1:A:389:THR:CB	1:A:390:PRO:HD3	0.55	2.29	8	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:327:CYS:HA	1:A:349:ALA:O	0.55	2.01	8	5
1:A:330:CYS:SG	1:A:351:HIS:CE1	0.55	2.99	3	4
1:A:334:GLY:O	1:A:335:VAL:HB	0.55	2.02	17	1
1:A:346:CYS:SG	1:A:388:HIS:CE1	0.53	3.02	13	4
1:A:334:GLY:O	1:A:335:VAL:HG22	0.52	2.03	14	3
1:A:353:THR:O	1:A:357:LYS:HG2	0.51	2.05	4	2
1:A:340:GLN:NE2	1:A:347:TYR:HB3	0.50	2.22	7	3
1:A:389:THR:OG1	1:A:390:PRO:HD3	0.50	2.06	16	2
1:A:353:THR:O	1:A:357:LYS:HB2	0.50	2.07	6	5
1:A:329:LEU:O	1:A:330:CYS:SG	0.50	2.70	15	3
1:A:342:HIS:O	1:A:343:LYS:HB2	0.49	2.07	14	4
1:A:367:VAL:HG12	1:A:378:SER:O	0.48	2.08	1	1
1:A:337:ALA:O	1:A:351:HIS:HA	0.48	2.09	3	1
1:A:327:CYS:SG	1:A:350:PHE:HA	0.47	2.50	10	4
1:A:330:CYS:SG	1:A:332:GLN:HG2	0.47	2.49	4	1
1:A:342:HIS:O	1:A:343:LYS:HB3	0.47	2.09	4	1
1:A:389:THR:CG2	1:A:390:PRO:HD3	0.46	2.41	8	1
1:A:341:CYS:SG	1:A:384:TYR:HA	0.46	2.50	11	1
1:A:370:LEU:HA	1:A:375:THR:HA	0.46	1.87	9	2
1:A:334:GLY:O	1:A:335:VAL:HG12	0.46	2.11	12	1
1:A:340:GLN:HA	1:A:348:THR:O	0.46	2.11	3	1
1:A:357:LYS:N	1:A:357:LYS:HE2	0.44	2.27	13	1
1:A:362:MET:HG2	1:A:383:ALA:HB2	0.44	1.88	12	1
1:A:389:THR:CB	1:A:390:PRO:CD	0.43	2.94	8	1
1:A:333:LYS:HD3	1:A:333:LYS:H	0.43	1.73	6	1
1:A:365:GLU:HB2	1:A:380:ARG:CG	0.43	2.44	13	1
1:A:380:ARG:HG3	1:A:380:ARG:O	0.42	2.14	13	1
1:A:369:GLU:CD	1:A:370:LEU:H	0.42	2.17	19	1
1:A:387:VAL:O	1:A:387:VAL:HG22	0.42	2.14	17	1
1:A:328:TYR:CE2	1:A:329:LEU:HG	0.42	2.50	6	1
1:A:341:CYS:HA	1:A:383:ALA:O	0.41	2.14	10	1
1:A:341:CYS:O	1:A:347:TYR:HA	0.41	2.14	6	1
1:A:369:GLU:HB2	1:A:376:THR:HG23	0.41	1.92	14	1
1:A:389:THR:OG1	1:A:390:PRO:HD2	0.41	2.15	14	1
1:A:327:CYS:SG	1:A:330:CYS:HB2	0.41	2.56	12	1
1:A:331:LYS:O	1:A:331:LYS:HG2	0.41	2.15	5	1
1:A:342:HIS:NE2	1:A:382:THR:HG23	0.41	2.30	11	1
1:A:343:LYS:HE2	1:A:343:LYS:HA	0.40	1.93	12	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	66/87 (76%)	53±2 (81±3%)	9±3 (13±4%)	4±1 (6±2%)	4	22
All	All	1320/1740 (76%)	1066 (81%)	178 (13%)	76 (6%)	4	22

All 18 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	330	CYS	19
1	A	343	LYS	14
1	A	390	PRO	7
1	A	335	VAL	6
1	A	346	CYS	5
1	A	336	GLY	4
1	A	370	LEU	4
1	A	333	LYS	3
1	A	373	GLY	2
1	A	387	VAL	2
1	A	344	ALA	2
1	A	347	TYR	2
1	A	334	GLY	1
1	A	325	LEU	1
1	A	342	HIS	1
1	A	389	THR	1
1	A	372	GLY	1
1	A	366	PRO	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	54/72 (75%)	51±2 (94±3%)	3±2 (6±3%)	27 72
All	All	1080/1440 (75%)	1012 (94%)	68 (6%)	27 72

All 24 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	341	CYS	9
1	A	347	TYR	6
1	A	370	LEU	6
1	A	357	LYS	5
1	A	368	LYS	4
1	A	333	LYS	3
1	A	389	THR	3
1	A	376	THR	3
1	A	325	LEU	3
1	A	348	THR	3
1	A	369	GLU	3
1	A	377	PHE	2
1	A	342	HIS	2
1	A	380	ARG	2
1	A	335	VAL	2
1	A	350	PHE	2
1	A	339	ILE	2
1	A	329	LEU	2
1	A	331	LYS	1
1	A	364	MET	1
1	A	345	ASN	1
1	A	371	THR	1
1	A	381	LYS	1
1	A	332	GLN	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 85% for the well-defined parts and 84% for the entire structure.

7.1 Chemical shift list 1

File name: 2lq6_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	887
Number of shifts mapped to atoms	887
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	77	0.00 ± 0.24	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	69	-0.05 ± 0.17	None needed (< 0.5 ppm)
$^{13}\text{C}'$	75	-2.27 ± 0.37	Should be applied
^{15}N	72	-1.26 ± 0.54	Should be applied

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 85%, i.e. 649 atoms were assigned a chemical shift out of a possible 766. 0 out of 9 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	318/326 (98%)	128/130 (98%)	127/132 (96%)	63/64 (98%)
Sidechain	306/368 (83%)	198/217 (91%)	105/137 (77%)	3/14 (21%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	25/72 (35%)	25/38 (66%)	0/30 (0%)	0/4 (0%)
Overall	649/766 (85%)	351/385 (91%)	232/299 (78%)	66/82 (80%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 84%, i.e. 785 atoms were assigned a chemical shift out of a possible 931. 0 out of 9 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	373/385 (97%)	149/153 (97%)	152/158 (96%)	72/74 (97%)
Sidechain	382/462 (83%)	249/275 (91%)	129/168 (77%)	4/19 (21%)
Aromatic	30/84 (36%)	29/44 (66%)	0/35 (0%)	1/5 (20%)
Overall	785/931 (84%)	427/472 (90%)	281/361 (78%)	77/98 (79%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	318	ILE	CG2	37.01	24.63 – 10.43	13.7

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

