



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:27 AM GMT

PDB ID : 3LZX  
Title : Crystal structure of ferredoxin-NADP<sup>+</sup> oxidoreductase from *Bacillus subtilis* (FORM II)  
Authors : Komori, H.; Seo, D.; Sakurai, T.; Higuchi, Y.  
Deposited on : 2010-03-02  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	<div> <div>5%</div> <div>92%</div> <div>8%</div> </div>
1	B	332	<div> <div>8%</div> <div>92%</div> <div>7%</div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5860 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ferredoxin–NADP reductase 2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	329	Total	C	N	O	S	0	0	0
			2570	1641	430	491	8			
1	B	329	Total	C	N	O	S	0	0	0
			2570	1641	430	491	8			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		
4	A	1	Total	Na	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	286	Total	O	0	0
			286	286		
5	B	230	Total	O	0	0
			230	230		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Ferredoxin–NADP reductase 2



#### • Molecule 1: Ferredoxin–NADP reductase 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	207.47Å 64.85Å 61.12Å 90.00° 105.82° 90.00°	Depositor
Resolution (Å)	26.85 – 1.90 26.41 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (26.85-1.90) 96.8 (26.41-1.90)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.83 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.181 , 0.211 0.188 , 0.218	Depositor DCC
$R_{free}$ test set	3014 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	22.2	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 47.3	EDS
Estimated twinning fraction	0.025 for -h-2*k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 59742 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5860	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, NAP, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/2618	0.58	0/3537
1	B	0.41	0/2618	0.56	0/3537
All	All	0.42	0/5236	0.57	0/7074

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2570	0	2576	15	1
1	B	2570	0	2576	16	0
2	A	53	0	31	0	0
2	B	53	0	31	0	0
3	A	48	0	25	0	0
3	B	48	0	25	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	286	0	0	0	0
5	B	230	0	0	0	0
All	All	5860	0	5264	28	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35:ILE:HD11	1:B:77:MET:HE1	1.72	0.71
1:B:132:GLU:CD	1:B:132:GLU:H	1.97	0.66
1:B:35:ILE:CD1	1:B:77:MET:HE1	2.31	0.60
1:B:183:ILE:HG21	1:B:226:LEU:HD22	1.84	0.59
1:A:129:LEU:HG	1:A:131:LEU:HD13	1.86	0.57
1:B:149:LEU:HD11	1:B:169:TRP:CZ3	2.40	0.57
1:A:149:LEU:HD11	1:A:172:MET:CE	2.37	0.54
1:A:54:TYR:O	1:B:326:THR:HG22	2.08	0.53
1:B:168:ASP:HB3	1:B:172:MET:HE2	1.93	0.51
1:B:146:VAL:HG12	1:B:149:LEU:HD12	1.93	0.51
1:B:132:GLU:CD	1:B:132:GLU:N	2.64	0.49
1:A:322:PRO:HG2	1:B:295:LEU:HD21	1.94	0.49
1:B:146:VAL:HG12	1:B:149:LEU:CD1	2.46	0.46
1:A:142:LEU:HD13	1:A:242:LEU:HD23	1.98	0.45
1:A:237:LEU:HD12	1:A:237:LEU:N	2.31	0.45
1:A:155:ARG:O	1:A:177:ALA:HB1	2.17	0.45
1:A:225:VAL:HG22	1:A:236:ILE:HG12	1.99	0.44
1:A:130:GLU:O	1:A:131:LEU:HD12	2.18	0.44
1:B:213:ALA:HB2	1:B:227:GLU:HG2	2.00	0.44
1:B:29:ARG:O	1:B:311:LYS:HE2	2.18	0.43
1:A:183:ILE:HG21	1:A:226:LEU:HD22	2.00	0.43
1:A:183:ILE:HG12	1:A:226:LEU:HD11	2.01	0.43
1:B:142:LEU:HD13	1:B:242:LEU:HD23	2.01	0.43
1:B:29:ARG:O	1:B:311:LYS:CE	2.67	0.42
1:A:159:ILE:HD11	1:A:173:LEU:HD12	2.02	0.42
1:A:216:ILE:HB	1:A:223:GLN:HB2	2.02	0.41
1:A:149:LEU:HD11	1:A:172:MET:HE2	2.02	0.41
1:A:293:VAL:HG12	1:B:320:VAL:HG22	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:228:GLU:O	1:A:317:LYS:NZ[4_556]	2.16	0.04



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/332 (98%)	316 (97%)	11 (3%)	0	100	100
1	B	327/332 (98%)	315 (96%)	12 (4%)	0	100	100
All	All	654/664 (98%)	631 (96%)	23 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/278 (99%)	273 (99%)	2 (1%)	88	88
1	B	275/278 (99%)	269 (98%)	6 (2%)	60	53
All	All	550/556 (99%)	542 (98%)	8 (2%)	72	69

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	TYR
1	A	280	PHE
1	B	25	TYR
1	B	132	GLU
1	B	174	GLU
1	B	193	GLU
1	B	261	ASP
1	B	280	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	A	1000	-	48,58,58	1.22	6 (12%)	54,89,89	1.99	8 (14%)
3	NAP	A	2000	-	42,52,52	0.96	3 (7%)	54,80,80	1.84	6 (11%)
2	FAD	B	1001	-	48,58,58	1.30	5 (10%)	54,89,89	2.09	7 (12%)
3	NAP	B	2001	-	42,52,52	0.91	2 (4%)	54,80,80	1.80	7 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	1000	-	-	0/30/50/50	0/6/6/6
3	NAP	A	2000	-	-	0/27/67/67	0/5/5/5

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	1001	-	-	0/30/50/50	0/6/6/6
3	NAP	B	2001	-	-	0/27/67/67	0/5/5/5

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2001	NAP	C6N-N1N	2.01	1.40	1.35
3	A	2000	NAP	C6N-N1N	2.17	1.41	1.35
2	A	1000	FAD	O4B-C1B	2.21	1.44	1.41
2	B	1001	FAD	C10-N1	2.24	1.39	1.35
2	A	1000	FAD	C9A-N10	2.25	1.41	1.38
2	A	1000	FAD	C4-N3	2.43	1.37	1.33
3	A	2000	NAP	C3N-C7N	2.62	1.54	1.50
2	B	1001	FAD	C5X-N5	2.65	1.39	1.35
2	A	1000	FAD	C10-N1	2.76	1.40	1.35
3	B	2001	NAP	O4B-C1B	2.79	1.44	1.41
2	B	1001	FAD	C4-N3	2.98	1.38	1.33
3	A	2000	NAP	O4D-C1D	3.04	1.45	1.41
2	A	1000	FAD	C5X-N5	3.45	1.40	1.35
2	A	1000	FAD	C4X-N5	3.80	1.39	1.33
2	B	1001	FAD	O4B-C1B	3.87	1.46	1.41
2	B	1001	FAD	C4X-N5	4.66	1.40	1.33

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2001	NAP	N3A-C2A-N1A	-9.20	121.85	128.89
2	B	1001	FAD	N3A-C2A-N1A	-8.80	122.16	128.89
3	A	2000	NAP	N3A-C2A-N1A	-8.62	122.29	128.89
2	A	1000	FAD	N3A-C2A-N1A	-8.23	122.59	128.89
3	A	2000	NAP	O3-PN-O5D	-6.99	84.39	102.94
3	B	2001	NAP	O3-PN-O5D	-4.69	90.49	102.94
2	B	1001	FAD	C4X-C4-N3	-3.67	118.56	123.59
2	A	1000	FAD	C4B-O4B-C1B	-3.57	105.80	109.72
2	A	1000	FAD	C4X-C4-N3	-3.28	119.11	123.59
2	B	1001	FAD	C4B-O4B-C1B	-2.92	106.51	109.72
3	A	2000	NAP	O2N-PN-O3	-2.90	91.92	105.09
2	A	1000	FAD	P-O3P-PA	-2.80	124.87	132.73
2	B	1001	FAD	P-O3P-PA	-2.62	125.39	132.73
3	A	2000	NAP	C4A-C5A-N7A	-2.60	107.08	109.48
3	B	2001	NAP	O2N-PN-O3	-2.42	94.10	105.09
3	B	2001	NAP	C4B-O4B-C1B	-2.21	107.29	109.72

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2001	NAP	C4A-C5A-N7A	-2.05	107.59	109.48
3	A	2000	NAP	O2N-PN-O1N	2.11	123.95	112.53
3	B	2001	NAP	O2N-PN-O1N	2.18	124.33	112.53
2	A	1000	FAD	O3P-PA-O5B	2.34	109.13	102.94
3	A	2000	NAP	O5D-PN-O1N	2.40	118.92	109.62
2	B	1001	FAD	C4X-N5-C5X	2.60	119.75	116.76
2	B	1001	FAD	C5X-C9A-N10	2.82	119.76	117.62
3	B	2001	NAP	O5D-PN-O1N	3.12	121.71	109.62
2	A	1000	FAD	C1'-N10-C9A	3.14	122.39	118.86
2	A	1000	FAD	C4X-N5-C5X	3.54	120.84	116.76
2	A	1000	FAD	C4-N3-C2	7.63	121.84	115.25
2	B	1001	FAD	C4-N3-C2	8.60	122.68	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	329/332 (99%)	0.15	17 (5%) 31 34	11, 22, 43, 52	0
1	B	329/332 (99%)	0.40	26 (7%) 15 17	13, 26, 46, 61	0
All	All	658/664 (99%)	0.28	43 (6%) 22 25	11, 24, 44, 61	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	318	ALA	9.7
1	B	328	LEU	8.6
1	B	329	PHE	7.5
1	B	320	VAL	6.2
1	B	327	SER	5.6
1	A	328	LEU	4.9
1	B	317	LYS	4.3
1	B	319	ARG	4.2
1	B	322	PRO	4.1
1	A	329	PHE	3.9
1	B	230	LYS	3.7
1	B	323	LEU	3.5
1	A	218	GLU	3.3
1	B	1	MET	3.3
1	A	230	LYS	3.2
1	B	326	THR	3.2
1	B	264	LYS	2.9
1	A	188	LYS	2.8
1	B	147	ASP	2.8
1	B	316	PRO	2.6
1	B	99	GLY	2.6
1	A	160	LEU	2.6
1	B	315	ASP	2.5
1	A	327	SER	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	194	HIS	2.5
1	A	150	GLN	2.4
1	A	116	ILE	2.4
1	B	117	ILE	2.3
1	B	232	ASP	2.3
1	A	194	HIS	2.3
1	A	147	ASP	2.3
1	A	159	ILE	2.3
1	B	218	GLU	2.2
1	A	145	PHE	2.2
1	B	159	ILE	2.2
1	A	319	ARG	2.2
1	B	150	GLN	2.1
1	A	176	ILE	2.1
1	A	283	ALA	2.1
1	B	261	ASP	2.1
1	B	242	LEU	2.1
1	A	135	GLU	2.0
1	B	13	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAP	A	2000	48/48	0.95	0.10	-0.28	22,26,36,40	0
2	FAD	A	1000	53/53	0.98	0.09	-0.47	10,13,19,20	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAP	B	2001	48/48	0.97	0.08	-0.51	19,23,25,27	0
2	FAD	B	1001	53/53	0.97	0.09	-0.77	14,18,21,22	0
4	NA	B	3002	1/1	0.99	0.05	-0.94	24,24,24,24	0
4	NA	A	3001	1/1	0.99	0.04	-1.86	19,19,19,19	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.