



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:34 PM GMT

PDB ID : 4M1Z
Title : Crystal structure of MycP1 with the N-terminal propeptide removed
Authors : Sun, D.M.; He, Y.; Wang, C.L.; Zang, J.Y.; Tian, C.L.
Deposited on : 2013-08-04
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1640 (2.28-2.24)
Clashscore	102246	1095 (2.26-2.26)
Ramachandran outliers	100387	1063 (2.26-2.26)
Sidechain outliers	100360	1063 (2.26-2.26)
RSRZ outliers	91569	1647 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	<div> <div>4%</div> <div>78%</div> <div>7%</div> <div>15%</div> </div>
1	B	372	<div> <div>2%</div> <div>74%</div> <div>10%</div> <div>15%</div> </div>
2	C	20	<div> <div>5%</div> <div>40%</div> <div>10%</div> <div>50%</div> </div>
2	D	20	<div> <div>10%</div> <div>45%</div> <div>20%</div> <div>35%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5033 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Membrane-anchored mycosin mycp1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	317	Total	C	N	O	S	1	0	0
			2288	1424	406	451	7			
1	B	315	Total	C	N	O	S	2	0	0
			2271	1413	403	448	7			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	51	MET	-	EXPRESSION TAG	UNP A0QNL1
A	52	GLY	-	EXPRESSION TAG	UNP A0QNL1
A	53	HIS	-	EXPRESSION TAG	UNP A0QNL1
A	54	HIS	-	EXPRESSION TAG	UNP A0QNL1
A	55	HIS	-	EXPRESSION TAG	UNP A0QNL1
A	56	HIS	-	EXPRESSION TAG	UNP A0QNL1
A	57	HIS	-	EXPRESSION TAG	UNP A0QNL1
A	58	HIS	-	EXPRESSION TAG	UNP A0QNL1
A	59	GLY	-	EXPRESSION TAG	UNP A0QNL1
A	60	SER	-	EXPRESSION TAG	UNP A0QNL1
A	61	HIS	-	EXPRESSION TAG	UNP A0QNL1
A	62	MET	-	EXPRESSION TAG	UNP A0QNL1
B	51	MET	-	EXPRESSION TAG	UNP A0QNL1
B	52	GLY	-	EXPRESSION TAG	UNP A0QNL1
B	53	HIS	-	EXPRESSION TAG	UNP A0QNL1
B	54	HIS	-	EXPRESSION TAG	UNP A0QNL1
B	55	HIS	-	EXPRESSION TAG	UNP A0QNL1
B	56	HIS	-	EXPRESSION TAG	UNP A0QNL1
B	57	HIS	-	EXPRESSION TAG	UNP A0QNL1
B	58	HIS	-	EXPRESSION TAG	UNP A0QNL1
B	59	GLY	-	EXPRESSION TAG	UNP A0QNL1
B	60	SER	-	EXPRESSION TAG	UNP A0QNL1
B	61	HIS	-	EXPRESSION TAG	UNP A0QNL1
B	62	MET	-	EXPRESSION TAG	UNP A0QNL1

- Molecule 2 is a protein called Membrane-anchored mycosin mycp1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	10	Total	C	N	O	0	0	0
			78	55	10	13			
2	D	13	Total	C	N	O	0	0	0
			105	72	17	16			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	140	Total	O	0	0
			140	140		
3	B	142	Total	O	0	0
			142	142		
3	C	5	Total	O	0	0
			5	5		
3	D	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Membrane-anchored mycosin mycp1




- Molecule 1: Membrane-anchored mycosin mycp1



- Molecule 2: Membrane-anchored mycosin mycp1



- Molecule 2: Membrane-anchored mycosin mycp1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.31Å 83.61Å 98.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.28 – 2.25 37.28 – 2.25	Depositor EDS
% Data completeness (in resolution range)	91.5 (37.28-2.25) 91.5 (37.28-2.25)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.13 (at 2.24Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.166 , 0.222 0.167 , 0.218	Depositor DCC
R_{free} test set	1519 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	30.4	Xtriage
Anisotropy	0.114	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.1	EDS
Estimated twinning fraction	0.019 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 30404 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5033	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/2340	0.68	0/3208
1	B	0.58	2/2322 (0.1%)	0.66	0/3185
2	C	0.54	0/83	0.57	0/118
2	D	0.40	0/110	0.57	0/153
All	All	0.58	2/4855 (0.0%)	0.67	0/6664

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	298	TRP	CD2-CE2	5.39	1.47	1.41
1	B	261	TRP	CD2-CE2	5.11	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2288	0	2225	21	0
1	B	2271	0	2208	29	0
2	C	78	0	78	3	0
2	D	105	0	113	2	0
3	A	140	0	0	2	0
3	B	142	0	0	3	0
3	C	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	4	0	0	0	0
All	All	5033	0	4624	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:GLN:HE22	1:A:295:SER:H	1.05	1.02
1:B:82:THR:H	1:B:349:GLN:HE22	1.09	0.95
1:B:246:GLN:HE22	1:B:295:SER:H	1.13	0.90
1:A:239:ASN:HD21	1:A:333:THR:H	1.20	0.89
2:C:407:VAL:HG22	2:C:408:PRO:HD2	1.61	0.81
1:B:239:ASN:HD21	1:B:333:THR:H	1.34	0.74
1:A:82:THR:H	1:A:349:GLN:HE22	1.37	0.72
1:B:82:THR:H	1:B:349:GLN:NE2	1.87	0.72
1:A:246:GLN:HE21	1:A:247:ASN:H	1.38	0.71
1:B:353:ASP:HB2	3:B:564:HOH:O	1.89	0.70
2:C:407:VAL:CG2	2:C:408:PRO:HD2	2.23	0.67
1:A:314:ASP:HB2	1:A:316:GLN:OE1	1.98	0.63
1:A:246:GLN:NE2	1:A:295:SER:H	1.87	0.62
1:A:83:GLY:H	1:A:349:GLN:NE2	2.01	0.59
1:B:305:GLY:O	1:B:331:ASN:HA	2.03	0.58
1:A:82:THR:H	1:A:349:GLN:NE2	2.03	0.57
1:A:96:ASN:ND2	1:A:119:ASP:H	2.05	0.54
1:A:100:ARG:HD3	1:A:315:GLY:O	2.08	0.52
1:B:246:GLN:HE21	1:B:247:ASN:H	1.55	0.52
1:B:190:HIS:HD2	3:B:528:HOH:O	1.92	0.51
1:A:243:ASP:HA	3:A:534:HOH:O	2.11	0.50
1:B:65:ARG:NH1	1:B:140:GLY:O	2.44	0.50
1:B:246:GLN:HE22	1:B:295:SER:N	1.96	0.50
1:B:319:ASN:O	1:B:320:ALA:HB2	2.11	0.50
2:D:403:ARG:HG2	2:D:404:VAL:HG23	1.93	0.50
1:A:246:GLN:HE21	1:A:247:ASN:N	2.07	0.49
1:A:83:GLY:H	1:A:349:GLN:HE21	1.60	0.49
1:B:211:ARG:HG2	1:B:213:ILE:CD1	2.42	0.49
1:B:246:GLN:NE2	1:B:295:SER:H	1.96	0.49
1:B:88:VAL:HG11	1:B:131:ILE:HG21	1.95	0.48
1:B:83:GLY:H	1:B:349:GLN:NE2	2.10	0.48
1:A:215:GLU:HA	3:A:541:HOH:O	2.13	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:83:GLY:H	1:B:349:GLN:HE21	1.62	0.47
1:B:64:ASP:OD1	1:B:64:ASP:N	2.47	0.47
1:B:96:ASN:ND2	1:B:119:ASP:H	2.13	0.46
1:A:111:ASP:O	1:A:113:ALA:O	2.33	0.46
1:B:246:GLN:HE21	1:B:247:ASN:N	2.13	0.46
1:A:201:ILE:HB	1:A:235:VAL:HG12	1.97	0.45
1:A:96:ASN:HD21	1:A:119:ASP:H	1.64	0.45
1:A:239:ASN:ND2	1:A:333:THR:HG23	2.32	0.45
1:B:67:TRP:CE2	1:B:285:GLN:HG2	2.53	0.43
1:B:290:SER:HB3	1:B:292:PHE:CE2	2.54	0.43
1:A:67:TRP:CE2	1:A:285:GLN:HG2	2.53	0.43
1:B:129:ALA:CB	1:B:311:LEU:HB2	2.48	0.43
2:C:407:VAL:HG22	2:C:408:PRO:CD	2.39	0.43
1:B:126:MET:HG2	1:B:310:SER:HB3	2.00	0.43
1:B:350:ARG:NE	3:B:561:HOH:O	2.51	0.43
1:B:141:PHE:CE2	1:B:310:SER:HA	2.55	0.41
1:A:106:GLY:O	1:A:153:LEU:HG	2.21	0.41
1:A:247:ASN:HA	1:A:248:PRO:HD3	1.97	0.41
2:D:413:ILE:HA	2:D:414:PRO:HD3	1.88	0.41
1:B:123:HIS:CE1	1:B:334:SER:OG	2.74	0.41
1:B:239:ASN:ND2	1:B:333:THR:HG23	2.36	0.40
1:B:131:ILE:HG13	1:B:338:ALA:HB1	2.03	0.40
1:B:83:GLY:N	1:B:349:GLN:NE2	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/372 (84%)	308 (98%)	5 (2%)	0	100	100
1	B	311/372 (84%)	303 (97%)	8 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	8/20 (40%)	8 (100%)	0	0	100	100
2	D	11/20 (55%)	11 (100%)	0	0	100	100
All	All	643/784 (82%)	630 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/286 (84%)	237 (99%)	2 (1%)	86	92
1	B	237/286 (83%)	235 (99%)	2 (1%)	86	92
2	C	10/19 (53%)	10 (100%)	0	100	100
2	D	13/19 (68%)	13 (100%)	0	100	100
All	All	499/610 (82%)	495 (99%)	4 (1%)	86	92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	136	SER
1	A	272	TRP
1	B	64	ASP
1	B	149	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	ASN
1	A	200	ASN
1	A	239	ASN
1	A	246	GLN
1	A	265	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	349	GLN
1	B	69	ASN
1	B	96	ASN
1	B	190	HIS
1	B	200	ASN
1	B	223	ASN
1	B	239	ASN
1	B	246	GLN
1	B	265	GLN
1	B	349	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/372 (85%)	-0.11	16 (5%) 32 36	19, 28, 61, 94	4 (1%)
1	B	315/372 (84%)	-0.27	9 (2%) 55 60	20, 28, 52, 84	8 (2%)
2	C	10/20 (50%)	0.40	1 (10%) 9 10	28, 38, 59, 62	0
2	D	13/20 (65%)	0.52	2 (15%) 3 3	30, 37, 68, 76	0
All	All	655/784 (83%)	-0.17	28 (4%) 39 43	19, 29, 59, 94	12 (1%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	397	PRO	6.1
1	A	398	GLU	5.8
1	A	396	GLY	5.4
1	A	112	ALA	4.1
2	D	415	PRO	4.0
1	A	113	ALA	4.0
1	A	395	ASP	3.9
2	D	414	PRO	3.9
1	A	182	ARG	3.5
1	A	159	ALA	3.3
1	B	63	ALA	3.2
1	B	243	ASP	3.1
1	A	394	PRO	3.0
1	A	179	GLY	2.9
1	B	324	GLU	2.9
1	A	181	ILE	2.9
2	C	415	PRO	2.8
1	B	325	ASP	2.7
1	B	158	VAL	2.7
1	A	242	GLN	2.6
1	B	242	GLN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	64	ASP	2.6
1	A	180	SER	2.5
1	A	111	ASP	2.4
1	B	137	PRO	2.4
1	A	114	GLY	2.3
1	A	399	LYS	2.2
1	B	320	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.