



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 27, 2016 – 02:28 AM BST

PDB ID : 2M4F  
Title : Solution Structure of Outer surface protein E  
Authors : Bhattacharjee, A.; Oeemig, J.S.; Kolodziejczyk, R.; Meri, T.; Kajander, T.;  
Iwai, H.; Jokiranta, T.; Goldman, A.  
Deposited on : 2013-02-05

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 92%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	151	 61% 35%

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 9 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:43-A:66, A:71-A:77, A:83-A:114, A:123-A:134, A:140-A:149, A:156-A:168 (98)	0.15	9

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20
2	4, 18

### 3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 2382 atoms, of which 1190 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Outer surface protein E.


Mol	Chain	Residues	Atoms						Trace
1	A	151	Total	C	H	N	O	S	0
			2382	753	1190	194	244	1	

## 4 Residue-property plots [i](#)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Outer surface protein E




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1


- Molecule 1: Outer surface protein E



#### 4.2.2 Score per residue for model 2


- Molecule 1: Outer surface protein E






#### 4.2.3 Score per residue for model 3

- Molecule 1: Outer surface protein E




#### 4.2.4 Score per residue for model 4

- Molecule 1: Outer surface protein E




#### 4.2.5 Score per residue for model 5

- Molecule 1: Outer surface protein E




#### 4.2.6 Score per residue for model 6

- Molecule 1: Outer surface protein E




#### 4.2.7 Score per residue for model 7

- Molecule 1: Outer surface protein E




#### 4.2.8 Score per residue for model 8

- Molecule 1: Outer surface protein E




#### 4.2.9 Score per residue for model 9 (medoid)

- Molecule 1: Outer surface protein E




### 4.2.10 Score per residue for model 10

- Molecule 1: Outer surface protein E




### 4.2.11 Score per residue for model 11

- Molecule 1: Outer surface protein E




### 4.2.12 Score per residue for model 12

- Molecule 1: Outer surface protein E



### 4.2.13 Score per residue for model 13


- Molecule 1: Outer surface protein E






### 4.2.14 Score per residue for model 14

- Molecule 1: Outer surface protein E




### 4.2.15 Score per residue for model 15

- Molecule 1: Outer surface protein E




### 4.2.16 Score per residue for model 16

- Molecule 1: Outer surface protein E




### 4.2.17 Score per residue for model 17

- Molecule 1: Outer surface protein E




### 4.2.18 Score per residue for model 18

- Molecule 1: Outer surface protein E




### 4.2.19 Score per residue for model 19

- Molecule 1: Outer surface protein E



### 4.2.20 Score per residue for model 20

- Molecule 1: Outer surface protein E



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
AMBER	refinement	
CYANA	structure solution	3

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2m4f_cs.str
Number of chemical shift lists	1
Total number of shifts	1836
Number of shifts mapped to atoms	1836
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	92%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.73±0.01	0±0/794 (0.0±0.0%)	0.85±0.01	0±0/1068 (0.0±0.0%)
All	All	0.73	0/15880 (0.0%)	0.85	3/21360 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.1±0.3
All	All	0	2

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	66	ARG	NE-CZ-NH1	5.83	123.21	120.30	13	3

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	111	TYR	Sidechain	1
1	A	159	TYR	Sidechain	1

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	781	787	787	1±1
All	All	15620	15740	15740	13

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:100:MET:SD	1:A:166:LEU:HD13	0.53	2.44	20	1
1:A:100:MET:CG	1:A:166:LEU:HD13	0.49	2.37	1	4
1:A:100:MET:HG3	1:A:166:LEU:HD13	0.46	1.88	2	4
1:A:149:ILE:CG2	1:A:157:ALA:HB2	0.44	2.43	20	1
1:A:93:VAL:O	1:A:97:VAL:HG23	0.44	2.13	4	2
1:A:161:ILE:CG2	1:A:166:LEU:HD11	0.41	2.46	4	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	98/151 (65%)	92±1 (94±1%)	6±1 (6±1%)	0±0 (0±0%)	59	88
All	All	1960/3020 (65%)	1841 (94%)	118 (6%)	1 (0%)	59	88

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	114	TYR	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/133 (65%)	79±1 (92±2%)	7±1 (8±2%)	20	65
All	All	1720/2660 (65%)	1583 (92%)	137 (8%)	20	65

All 20 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	149	ILE	20
1	A	87	SER	19
1	A	88	LEU	17
1	A	100	MET	16
1	A	95	ASN	12
1	A	163	LEU	11
1	A	51	LYS	10
1	A	107	LYS	7
1	A	47	LYS	4
1	A	45	THR	4
1	A	73	ASP	4
1	A	132	LYS	3
1	A	43	LYS	2
1	A	98	LYS	2
1	A	127	LYS	1
1	A	53	LYS	1
1	A	147	ASP	1
1	A	50	ASN	1
1	A	52	ASP	1
1	A	77	ASN	1

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 92% for the well-defined parts and 90% for the entire structure.

### 7.1 Chemical shift list 1

File name: 2m4f\_cs.str

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1836
Number of shifts mapped to atoms	1836
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	13

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	151	$-0.05 \pm 0.10$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	137	$-0.21 \pm 0.15$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	151	$0.20 \pm 0.22$	None needed ( $< 0.5$ ppm)
$^{15}\text{N}$	150	$-0.17 \pm 0.31$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 92%, i.e. 1113 atoms were assigned a chemical shift out of a possible 1204. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	490/490 (100%)	196/196 (100%)	196/196 (100%)	98/98 (100%)
Sidechain	543/608 (89%)	331/351 (94%)	204/235 (87%)	8/22 (36%)

*Continued on next page...*



Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	80/106 (75%)	48/56 (86%)	31/49 (63%)	1/1 (100%)
Overall	1113/1204 (92%)	575/603 (95%)	431/480 (90%)	107/121 (88%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 90%, i.e. 1652 atoms were assigned a chemical shift out of a possible 1842. 0 out of 18 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	752/755 (100%)	300/302 (99%)	302/302 (100%)	150/151 (99%)
Sidechain	804/950 (85%)	486/553 (88%)	306/359 (85%)	12/38 (32%)
Aromatic	96/137 (70%)	60/73 (82%)	35/61 (57%)	1/3 (33%)
Overall	1652/1842 (90%)	846/928 (91%)	643/722 (89%)	163/192 (85%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	51	LYS	HE2	1.55	3.87 – 1.97	-7.2
1	A	51	LYS	HE3	1.55	3.86 – 1.96	-7.1
1	A	169	ASN	HB3	0.46	4.41 – 1.11	-7.0
1	A	51	LYS	HG2	-0.26	2.67 – 0.07	-6.3
1	A	136	ILE	HD12	-1.00	2.13 – -0.77	-5.8
1	A	136	ILE	HD13	-1.00	2.13 – -0.77	-5.8
1	A	136	ILE	HD11	-1.00	2.13 – -0.77	-5.8
1	A	65	VAL	HB	0.29	3.59 – 0.39	-5.3
1	A	160	ALA	HB2	0.05	2.61 – 0.11	-5.2
1	A	160	ALA	HB1	0.05	2.61 – 0.11	-5.2
1	A	160	ALA	HB3	0.05	2.61 – 0.11	-5.2
1	A	127	LYS	HE2	1.95	3.87 – 1.97	-5.1
1	A	129	ILE	HB	0.32	3.24 – 0.34	-5.1

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

