



Full wwPDB NMR Structure Validation Report ⓘ

Apr 27, 2016 – 02:56 AM BST

PDB ID : 2M6S
Title : Holo_YqcA
Authors : Jin, C.; Hu, Y.; Ye, Q.
Deposited on : 2013-04-09

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment is 90%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	149	 95%

2 Ensemble composition and analysis ⓘ

This entry contains 20 models. Model 17 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:2-A:59, A:63-A:149 (145)	0.29	17

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	3, 4, 6, 7, 8, 9, 11, 12, 14, 15, 17, 18, 20
2	2, 5, 13, 16, 19
Single-model clusters	1; 10


3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2297 atoms, of which 1118 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Flavodoxin.

Mol	Chain	Residues	Atoms						Trace
1	A	149	Total	C	H	N	O	S	0
			2247	730	1099	181	233	4	

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: $C_{17}H_{21}N_4O_9P$).




Mol	Chain	Residues	Atoms					
2	A	1	Total	C	H	N	O	P
			50	17	19	4	9	1

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Flavodoxin




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Flavodoxin



4.2.2 Score per residue for model 2

- Molecule 1: Flavodoxin



4.2.3 Score per residue for model 3

- Molecule 1: Flavodoxin

Chain A:  93% ..



4.2.4 Score per residue for model 4

- Molecule 1: Flavodoxin

Chain A:  92% 5% .



4.2.5 Score per residue for model 5

- Molecule 1: Flavodoxin

Chain A:  94% ..



4.2.6 Score per residue for model 6

- Molecule 1: Flavodoxin

Chain A:  93% ...



4.2.7 Score per residue for model 7

- Molecule 1: Flavodoxin

Chain A:  94% ..



4.2.8 Score per residue for model 8

- Molecule 1: Flavodoxin

Chain A:  93% ..



4.2.9 Score per residue for model 9

- Molecule 1: Flavodoxin

Chain A:  94% ..



4.2.10 Score per residue for model 10

- Molecule 1: Flavodoxin

Chain A:  96% ..



4.2.11 Score per residue for model 11

- Molecule 1: Flavodoxin

Chain A:  91% 6% .



4.2.12 Score per residue for model 12

- Molecule 1: Flavodoxin

Chain A:  96% ..



4.2.13 Score per residue for model 13

- Molecule 1: Flavodoxin


Chain A:  94% . .



4.2.14 Score per residue for model 14

- Molecule 1: Flavodoxin

Chain A:  94% . .



4.2.15 Score per residue for model 15

- Molecule 1: Flavodoxin


Chain A:  93% 5% .



4.2.16 Score per residue for model 16

- Molecule 1: Flavodoxin

Chain A:  95% . .



4.2.17 Score per residue for model 17 (medoid)

- Molecule 1: Flavodoxin

Chain A:  93% . . .



4.2.18 Score per residue for model 18

- Molecule 1: Flavodoxin

Chain A:  92% 5% •



4.2.19 Score per residue for model 19

- Molecule 1: Flavodoxin

Chain A:  91% 6% •



4.2.20 Score per residue for model 20

- Molecule 1: Flavodoxin

Chain A:  93% • •



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	v2.1
AMBER	refinement	v9

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2m6s_cs.str
Number of chemical shift lists	1
Total number of shifts	1764
Number of shifts mapped to atoms	1764
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	90%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.71±0.00	0±0/1149 (0.0±0.0%)	0.98±0.01	1±0/1567 (0.1±0.0%)
All	All	0.71	0/22980 (0.0%)	0.98	16/31340 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.1±0.2
All	All	0	1

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	118	ARG	NE-CZ-NH1	6.92	123.76	120.30	5	16

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	51	TYR	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1123	1074	1074	0±0
2	A	31	19	19	0±0
All	All	23080	21860	21860	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:44:TRP:CH2	1:A:52:VAL:HG21	0.50	2.41	11	4
1:A:44:TRP:CZ2	1:A:52:VAL:HG21	0.42	2.49	20	1
1:A:123:LEU:HD13	1:A:124:LEU:N	0.41	2.31	19	2

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/149 (97%)	131±3 (91±2%)	13±3 (9±2%)	1±1 (1±1%)	38	79
All	All	2880/2980 (97%)	2613 (91%)	252 (9%)	15 (1%)	38	79

All 6 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	119	VAL	6
1	A	115	SER	5
1	A	13	GLY	1
1	A	100	PHE	1
1	A	30	GLY	1
1	A	41	LEU	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/125 (98%)	120±1 (97±1%)	4±1 (3±1%)	55	91
All	All	2460/2500 (98%)	2390 (97%)	70 (3%)	55	91

All 24 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	53	LEU	16
1	A	41	LEU	6
1	A	40	GLU	6
1	A	89	VAL	5
1	A	37	GLU	4
1	A	82	GLN	4
1	A	119	VAL	4
1	A	112	GLN	3
1	A	99	ASN	3
1	A	14	ASN	2
1	A	51	TYR	2
1	A	86	ARG	2
1	A	15	SER	2
1	A	123	LEU	1
1	A	48	GLN	1
1	A	64	LEU	1
1	A	135	THR	1
1	A	11	MET	1
1	A	84	ASN	1
1	A	43	ASP	1
1	A	71	LEU	1
1	A	59	THR	1
1	A	148	LEU	1
1	A	45	LEU	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	FMN	A	201	-	31,33,33	1.92±0.05	2±0 (5±1%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	FMN	A	201	-	32,50,50	2.18±0.07	1±0 (3±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	A	201	-	-	0±0,18,18,18	0±0,3,3,3

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	A	201	FMN	C4A-N5	6.66	1.43	1.33	14	20
2	A	201	FMN	C5A-N5	5.81	1.44	1.35	20	17

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	A	201	FMN	C2-N1-C10	11.58	123.27	113.39	11	20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 90% for the well-defined parts and 90% for the entire structure.

7.1 Chemical shift list 1

File name: 2m6s_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1764
Number of shifts mapped to atoms	1764
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	15

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	148	-0.57 ± 0.13	Should be applied
$^{13}\text{C}_\beta$	134	0.35 ± 0.13	None needed (< 0.5 ppm)
$^{13}\text{C}'$	148	-0.13 ± 0.12	None needed (< 0.5 ppm)
^{15}N	139	-0.99 ± 0.50	None needed (imprecise)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 90%, i.e. 1534 atoms were assigned a chemical shift out of a possible 1696. 25 out of 28 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	707/709 (100%)	281/282 (100%)	290/290 (100%)	136/137 (99%)
Sidechain	747/843 (89%)	451/489 (92%)	281/330 (85%)	15/24 (62%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	80/144 (56%)	41/76 (54%)	35/63 (56%)	4/5 (80%)
Overall	1534/1696 (90%)	773/847 (91%)	606/683 (89%)	155/166 (93%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 90%, i.e. 1555 atoms were assigned a chemical shift out of a possible 1734. 25 out of 28 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	722/729 (99%)	287/290 (99%)	296/298 (99%)	139/141 (99%)
Sidechain	753/861 (87%)	455/500 (91%)	283/336 (84%)	15/25 (60%)
Aromatic	80/144 (56%)	41/76 (54%)	35/63 (56%)	4/5 (80%)
Overall	1555/1734 (90%)	783/866 (90%)	614/697 (88%)	158/171 (92%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	71	LEU	CA	158.32	66.36 – 44.96	48.0
1	A	86	ARG	NE	113.85	92.63 – 76.73	18.3
1	A	118	ARG	NE	113.68	92.63 – 76.73	18.2
1	A	148	LEU	HG	-0.67	3.16 – -0.14	-6.6
1	A	148	LEU	HD13	-0.95	2.16 – -0.64	-6.1
1	A	148	LEU	HD12	-0.95	2.16 – -0.64	-6.1
1	A	148	LEU	HD11	-0.95	2.16 – -0.64	-6.1
1	A	83	PRO	HD2	1.60	5.45 – 1.85	-5.7
1	A	110	LEU	HG	-0.24	3.16 – -0.14	-5.3
1	A	92	LEU	CG	32.86	32.55 – 21.05	5.3
1	A	123	LEU	HB3	-0.34	3.34 – -0.26	-5.2
1	A	145	GLY	N	129.70	129.07 – 90.27	5.2
1	A	75	ILE	HD13	-0.79	2.13 – -0.77	-5.1
1	A	75	ILE	HD12	-0.79	2.13 – -0.77	-5.1
1	A	75	ILE	HD11	-0.79	2.13 – -0.77	-5.1

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-

defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

