



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 08:49 PM GMT

PDB ID : 1M76  
Title : Crystal Structure of the S137C Mutant of L-3-HYDROXYACYL-COA Dehydrogenase in Complex with NAD and Acetoacetyl-COA  
Authors : Barycki, J.J.; Banaszak, L.J.  
Deposited on : 2002-07-18  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	
1	B	302	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5182 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3-HYDROXYACYL-COA DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	291	Total	C	N	O	S	0	0	0
			2243	1435	375	424	9			
1	B	291	Total	C	N	O	S	0	0	0
			2243	1435	375	424	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	137	CYS	SER	ENGINEERED	UNP Q16836
B	137	CYS	SER	ENGINEERED	UNP Q16836

- Molecule 2 is ACETOACETYL-COENZYME A (three-letter code: CAA) (formula: C<sub>25</sub>H<sub>40</sub>N<sub>7</sub>O<sub>18</sub>P<sub>3</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P S	3	0
			54	25	7	18	3 1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	S	3	0
			54	25	7	18	3	1		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	O	P		0	0
			44	21	7	14	2			
3	B	1	Total	C	N	O	P		0	0
			44	21	7	14	2			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	223	Total	O	0	0
			223	223		
4	B	277	Total	O	0	0
			277	277		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: 3-HYDROXYACYL-COA DEHYDROGENASE



#### • Molecule 1: 3-HYDROXYACYL-COA DEHYDROGENASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.39 Å 87.99 Å 158.97 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.21 – 2.15 31.21 – 2.15	Depositor EDS
% Data completeness (in resolution range)	98.4 (31.21-2.15) 98.5 (31.21-2.15)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.29 (at 2.16 Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.208 , 0.257 0.214 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	22.2	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	2 of 38752 reflections (0.005%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5182	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CAA, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/2285	0.58	0/3085
1	B	0.34	0/2285	0.59	0/3085
All	All	0.33	0/4570	0.58	0/6170

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2243	0	2293	36	0
1	B	2243	0	2293	39	0
2	A	54	0	36	0	0
2	B	54	0	36	2	0
3	A	44	0	26	2	0
3	B	44	0	26	0	0
4	A	223	0	0	9	0
4	B	277	0	0	7	0
All	All	5182	0	4710	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:60:GLU:O	1:B:64:LYS:HD3	1.71	0.88
1:A:269:ASP:OD2	1:A:272:ASN:HB2	1.76	0.85
1:A:293:LYS:HG3	4:A:823:HOH:O	1.77	0.83
1:B:116:ASN:HD21	1:B:144:ILE:HA	1.46	0.80
1:A:173:LYS:HE3	1:A:182:PHE:CG	2.17	0.78
1:B:56:LYS:O	1:B:60:GLU:HG3	1.92	0.69
1:B:263:ASP:O	1:B:267:GLU:HG3	1.92	0.69
1:B:116:ASN:ND2	1:B:144:ILE:HA	2.10	0.65
1:A:59:GLU:HG3	1:A:85:LEU:HD11	1.79	0.64
1:A:70:PHE:HB3	1:A:73:ASN:O	1.99	0.62
1:B:269:ASP:OD2	1:B:272:ASN:HB2	2.01	0.61
1:A:273:PRO:HA	1:A:276:GLN:HG2	1.83	0.61
1:B:63:ARG:O	1:B:67:LYS:HG2	2.02	0.60
1:B:12:LYS:N	1:B:180:LYS:HE2	2.18	0.58
1:A:190:LYS:HD2	4:A:1003:HOH:O	2.04	0.57
1:B:221:LEU:HD22	1:B:232:ILE:HD11	1.89	0.55
1:B:173:LYS:HG2	4:B:860:HOH:O	2.07	0.55
1:B:22:GLY:O	1:B:27:GLY:HA3	2.06	0.55
1:A:221:LEU:HD12	1:A:224:ARG:HH21	1.72	0.54
1:B:300:LYS:HD2	4:B:1280:HOH:O	2.06	0.54
1:A:219:ILE:O	1:A:223:GLU:HG3	2.07	0.54
1:B:200:LYS:NZ	4:B:1282:HOH:O	2.38	0.54
1:A:150:ARG:HG2	1:A:153:ARG:HB2	1.90	0.53
1:A:78:ASP:O	1:A:82:GLU:HG3	2.08	0.53
1:A:22:GLY:O	1:A:27:GLY:HA3	2.08	0.53
1:A:296:GLU:HG3	1:A:300:LYS:HG2	1.91	0.52
1:B:64:LYS:N	1:B:64:LYS:HD2	2.24	0.52
1:A:150:ARG:O	1:A:150:ARG:HG2	2.10	0.51
1:B:63:ARG:HH11	1:B:63:ARG:HG2	1.75	0.51
1:B:64:LYS:N	1:B:64:LYS:CD	2.73	0.51
1:A:283:LYS:O	1:A:287:GLU:HG3	2.10	0.51
1:B:68:LYS:NZ	2:B:751:CAA:H3B	2.26	0.50
1:A:172:ILE:HD12	1:A:172:ILE:N	2.27	0.50
1:A:212:VAL:HB	1:A:213:PRO:HD3	1.94	0.50
1:B:150:ARG:HG2	1:B:150:ARG:O	2.11	0.50
1:A:99:HIS:HB2	4:A:1240:HOH:O	2.11	0.49
1:A:95:ALA:HB2	1:A:121:ARG:HH12	1.79	0.48
1:B:292:LYS:HD2	4:B:1158:HOH:O	2.14	0.47
1:B:68:LYS:HZ3	2:B:751:CAA:H3B	1.79	0.47
1:A:292:LYS:HD2	4:A:1067:HOH:O	2.15	0.47

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:63:ARG:NH1	1:B:63:ARG:HG2	2.30	0.46
1:B:283:LYS:O	1:B:287:GLU:HG3	2.16	0.46
1:B:150:ARG:HG2	1:B:153:ARG:HB2	1.98	0.46
1:B:243:PRO:HD2	4:B:805:HOH:O	2.14	0.46
1:B:138:SER:HB3	1:B:261:ILE:HG23	1.97	0.46
1:A:221:LEU:HD12	1:A:224:ARG:NH2	2.30	0.46
1:A:47:THR:OG1	1:A:50:ILE:HG13	2.16	0.46
1:A:53:LYS:HD3	4:A:1226:HOH:O	2.16	0.45
1:B:259:LYS:HE3	4:B:913:HOH:O	2.16	0.45
1:A:138:SER:HB3	1:A:261:ILE:HG23	1.99	0.44
1:A:140:GLN:HG3	1:A:201:ASP:OD2	2.17	0.44
1:A:67:LYS:HE2	4:A:1233:HOH:O	2.18	0.44
1:B:133:ALA:HA	1:B:155:ALA:O	2.18	0.44
1:B:99:HIS:O	1:B:127:ALA:HB2	2.18	0.43
1:B:12:LYS:HA	1:B:180:LYS:NZ	2.33	0.43
1:A:87:THR:O	1:A:87:THR:HG22	2.19	0.43
1:B:51:LEU:HD13	1:B:90:THR:HB	1.99	0.43
1:A:138:SER:HB2	1:A:265:TRP:NE1	2.34	0.42
1:A:103:LEU:HD12	1:A:131:ILE:O	2.19	0.42
1:A:24:GLY:HA3	3:A:350:NAD:O5B	2.18	0.42
1:A:137:CYS:SG	3:A:350:NAD:C6N	3.08	0.42
1:B:162:PRO:HG2	1:B:165:VAL:HB	2.02	0.42
1:B:173:LYS:HB2	1:B:177:THR:HB	2.02	0.41
1:A:243:PRO:HG3	4:B:1000:HOH:O	2.21	0.41
1:A:278:SER:HA	1:A:279:PRO:HD3	1.91	0.41
1:B:179:GLN:O	1:B:183:GLU:HG3	2.20	0.41
1:B:12:LYS:CA	1:B:180:LYS:HZ1	2.33	0.41
1:B:87:THR:O	1:B:87:THR:HG22	2.21	0.41
1:A:176:MET:HE1	4:A:1098:HOH:O	2.20	0.41
1:B:259:LYS:HE2	1:B:282:ASN:OD1	2.21	0.41
1:A:108:ILE:HD13	1:A:118:LEU:HD22	2.03	0.41
1:A:225:GLY:HA2	4:A:829:HOH:O	2.21	0.40
1:B:41:VAL:HG12	1:B:42:VAL:N	2.36	0.40
1:B:221:LEU:C	1:B:221:LEU:HD23	2.42	0.40
4:A:998:HOH:O	1:B:243:PRO:HG3	2.19	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/302 (96%)	278 (96%)	11 (4%)	0	100	100
1	B	289/302 (96%)	278 (96%)	11 (4%)	0	100	100
All	All	578/604 (96%)	556 (96%)	22 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/253 (97%)	244 (100%)	1 (0%)	93	96
1	B	245/253 (97%)	244 (100%)	1 (0%)	93	96
All	All	490/506 (97%)	488 (100%)	2 (0%)	93	96

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	214	TYR
1	B	214	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	32	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	116	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAD	A	350	-	38,48,48	1.99	8 (21%)	47,73,73	1.71	14 (29%)
2	CAA	A	351	-	45,56,56	1.96	11 (24%)	58,83,83	2.75	10 (17%)
3	NAD	B	750	-	38,48,48	2.02	7 (18%)	47,73,73	1.57	9 (19%)
2	CAA	B	751	-	45,56,56	1.95	11 (24%)	58,83,83	2.61	9 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	350	-	-	0/22/62/62	0/5/5/5
2	CAA	A	351	-	-	0/50/71/71	0/3/3/3
3	NAD	B	750	-	-	0/22/62/62	0/5/5/5
2	CAA	B	751	-	-	1/50/71/71	0/3/3/3

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	350	NAD	C6N-C5N	-2.92	1.32	1.38
3	B	750	NAD	C5B-C4B	-2.89	1.42	1.51
3	A	350	NAD	C5B-C4B	-2.86	1.42	1.51
3	B	750	NAD	C6N-C5N	-2.82	1.32	1.38
2	B	751	CAA	C7P-N8P	-2.62	1.40	1.46
2	B	751	CAA	C3P-N4P	-2.33	1.40	1.46
2	A	351	CAA	C3P-N4P	-2.31	1.40	1.46
2	A	351	CAA	C7P-N8P	-2.27	1.40	1.46
3	A	350	NAD	C3N-C7N	-2.24	1.47	1.50
3	A	350	NAD	C2A-N1A	2.15	1.38	1.33
2	A	351	CAA	O1-C1	2.29	1.24	1.21
3	B	750	NAD	C2A-N1A	2.37	1.38	1.33
2	A	351	CAA	O4B-C1B	2.47	1.44	1.41
2	B	751	CAA	C1-S1P	2.49	1.81	1.76
2	B	751	CAA	O4B-C1B	2.51	1.44	1.41
2	B	751	CAA	C6A-N6A	2.76	1.43	1.34
2	A	351	CAA	C6A-N6A	2.76	1.43	1.34
3	A	350	NAD	C6N-N1N	2.92	1.43	1.35
3	B	750	NAD	C6N-N1N	2.97	1.43	1.35
2	B	751	CAA	C5P-N4P	3.11	1.40	1.33
2	B	751	CAA	C8A-N7A	3.24	1.40	1.34
2	A	351	CAA	C8A-N7A	3.34	1.41	1.34
2	A	351	CAA	C5P-N4P	3.43	1.41	1.33
2	A	351	CAA	C9P-N8P	3.72	1.41	1.33
2	B	751	CAA	C9P-N8P	3.91	1.41	1.33
2	B	751	CAA	C2A-N1A	4.40	1.42	1.33
2	A	351	CAA	C2A-N1A	4.42	1.42	1.33
2	B	751	CAA	C4A-N3A	4.55	1.42	1.35
2	A	351	CAA	C4A-N3A	4.56	1.42	1.35
3	A	350	NAD	C5N-C4N	4.99	1.49	1.38
3	B	750	NAD	C5N-C4N	5.01	1.49	1.38
2	B	751	CAA	C2A-N3A	5.46	1.41	1.32
2	A	351	CAA	C2A-N3A	5.54	1.42	1.32
3	A	350	NAD	C2N-C3N	6.11	1.48	1.39
3	A	350	NAD	C4N-C3N	6.27	1.50	1.39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	750	NAD	C4N-C3N	6.38	1.50	1.39
3	B	750	NAD	C2N-C3N	6.43	1.48	1.39

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	351	CAA	N3A-C2A-N1A	-17.25	115.68	128.89
2	B	751	CAA	N3A-C2A-N1A	-16.62	116.17	128.89
2	A	351	CAA	O1-C1-S1P	-4.83	119.00	122.83
3	A	350	NAD	N3A-C2A-N1A	-4.13	125.73	128.89
3	B	750	NAD	O4D-C4D-C5D	-3.57	96.55	109.32
3	B	750	NAD	N3A-C2A-N1A	-3.49	126.22	128.89
3	A	350	NAD	O4D-C4D-C5D	-3.23	97.78	109.32
2	A	351	CAA	P2A-O3A-P1A	-3.21	123.73	132.73
3	B	750	NAD	O4D-C1D-N1N	-2.70	105.16	108.13
2	B	751	CAA	P2A-O3A-P1A	-2.57	125.50	132.73
3	A	350	NAD	C1B-N9A-C4A	-2.55	123.10	126.94
3	A	350	NAD	O3D-C3D-C4D	-2.36	103.98	111.05
3	B	750	NAD	O7N-C7N-C3N	-2.36	117.01	119.59
3	A	350	NAD	O7N-C7N-C3N	-2.35	117.03	119.59
3	B	750	NAD	C2N-C3N-C4N	-2.13	115.92	118.29
3	A	350	NAD	C5N-C4N-C3N	-2.10	117.69	120.33
2	B	751	CAA	O2A-P1A-O5B	2.05	118.79	108.46
2	A	351	CAA	O2A-P1A-O5B	2.05	118.82	108.46
3	A	350	NAD	O3-PA-O5B	2.06	108.41	102.94
3	A	350	NAD	C6N-C5N-C4N	2.07	122.58	119.44
2	A	351	CAA	O4B-C4B-C3B	2.10	109.70	104.86
3	A	350	NAD	O2N-PN-O5D	2.21	119.63	108.46
2	B	751	CAA	C2B-C1B-N9A	2.28	117.77	114.29
3	A	350	NAD	C5D-C4D-C3D	2.36	124.57	115.21
3	A	350	NAD	C4D-O4D-C1D	2.38	112.34	109.72
2	B	751	CAA	C3B-C2B-C1B	2.46	105.87	99.98
2	A	351	CAA	C3B-C2B-C1B	2.48	105.93	99.98
2	B	751	CAA	C2A-N1A-C6A	2.53	123.29	118.77
3	B	750	NAD	C5D-C4D-C3D	2.55	125.32	115.21
2	B	751	CAA	O3B-P3B-O7A	2.58	113.56	107.11
2	A	351	CAA	O3B-P3B-O7A	2.65	113.72	107.11
3	B	750	NAD	C4D-O4D-C1D	2.68	112.66	109.72
2	A	351	CAA	C2A-N1A-C6A	2.68	123.56	118.77
3	A	350	NAD	C4A-C5A-N7A	2.70	111.96	109.48
3	B	750	NAD	C4A-C5A-N7A	2.84	112.09	109.48
3	A	350	NAD	C2B-C1B-N9A	2.89	118.70	114.29

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	751	CAA	O4B-C1B-N9A	3.17	114.73	108.10
2	A	351	CAA	O4B-C1B-N9A	3.52	115.47	108.10
2	B	751	CAA	C2-C1-S1P	4.30	117.79	113.50
2	A	351	CAA	C2-C1-S1P	4.36	117.85	113.50
3	B	750	NAD	C3N-C7N-N7N	4.39	122.63	117.82
3	A	350	NAD	C3N-C7N-N7N	4.66	122.91	117.82

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	751	CAA	O3-C3-C2-C1

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	350	NAD	2	0
2	B	751	CAA	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	291/302 (96%)	0.44	22 (7%) 17 23	12, 21, 45, 57	0
1	B	291/302 (96%)	0.12	6 (2%) 67 74	12, 18, 35, 49	0
All	All	582/604 (96%)	0.28	28 (4%) 34 45	12, 19, 40, 57	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	302	LYS	5.4
1	A	66	ALA	5.4
1	A	73	ASN	4.6
1	A	68	LYS	4.5
1	A	69	LYS	4.1
1	A	74	PRO	4.1
1	A	70	PHE	4.1
1	A	71	ALA	3.8
1	B	268	MET	3.8
1	B	68	LYS	3.7
1	A	75	LYS	3.6
1	A	13	ILE	3.6
1	B	302	LYS	3.6
1	B	14	ILE	3.1
1	B	13	ILE	3.1
1	A	268	MET	2.9
1	A	65	VAL	2.9
1	A	64	LYS	2.7
1	A	271	GLU	2.6
1	A	128	GLU	2.5
1	B	12	LYS	2.4
1	A	159	PHE	2.4
1	A	211	LEU	2.3
1	A	105	VAL	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	30	ILE	2.3
1	A	49	ASP	2.1
1	A	72	GLU	2.1
1	A	12	LYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CAA	A	351	54/54	0.75	0.25	2.00	12,52,80,81	3
2	CAA	B	751	54/54	0.75	0.27	1.84	15,53,83,84	3
3	NAD	B	750	44/44	0.95	0.12	-0.69	11,16,20,21	0
3	NAD	A	350	44/44	0.95	0.13	-0.74	14,17,25,27	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.