



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2017 – 05:53 PM EST

PDB ID : 5M7L  
Title : Blastochloris viridis photosynthetic reaction center synchrotron structure  
Authors : Sharma, A.S.; Johansson, L.; Dunevall, E.; Wahlgren, W.Y.; Neutze, R.; Kato, G.  
Deposited on : 2016-10-28  
Resolution : 3.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1408 (3.80-3.40)
Clashscore	102246	1010 (3.74-3.46)
Ramachandran outliers	100387	1007 (3.76-3.44)
Sidechain outliers	100360	1007 (3.76-3.44)
RSRZ outliers	91569	1003 (3.78-3.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	356	<div> <div>81%</div> <div>12%</div> <div>7%</div> </div>
2	B	274	<div> <div>82%</div> <div>16%</div> <div>••</div> </div>
3	C	324	<div> <div>85%</div> <div>14%</div> </div>
4	D	258	<div> <div>4%</div> <div>67%</div> <div>23%</div> <div>•• 6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
11	MQ7	C	404	-	-	-	X
12	NS5	C	405	-	-	-	X
13	OTP	C	406	-	-	-	X
14	PO4	C	408	-	-	-	X
6	DGA	A	405	-	-	-	X
9	MPG	B	305	-	-	-	X
9	MPG	B	306	-	-	-	X
9	MPG	C	407	-	-	-	X

## 2 Entry composition

There are 14 unique types of molecules in this entry. The entry contains 9890 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Photosynthetic reaction center cytochrome c subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	332	Total	C	N	O	S	0	0	0
			2598	1637	465	478	18			

- Molecule 2 is a protein called Reaction center protein L chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	273	Total	C	N	O	S	0	2	0
			2170	1458	350	355	7			


- Molecule 3 is a protein called Reaction center protein M chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	323	Total	C	N	O	S	0	0	0
			2546	1696	417	422	11			

- Molecule 4 is a protein called Reaction center protein H chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	243	Total	C	N	O	S	0	0	0
			1771	1140	297	332	2			

- Molecule 5 is HEME C (three-letter code: HEC) (formula:  $C_{34}H_{34}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 6 is DIACYL GLYCEROL (three-letter code: DGA) (formula: C<sub>39</sub>H<sub>76</sub>O<sub>5</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			37	33	4		

- Molecule 7 is BACTERIOCHLOROPHYLL A (three-letter code: BCL) (formula: C<sub>55</sub>H<sub>74</sub>MgN<sub>4</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	B	1	Total	C	Mg	N	O	0	0
			65	54	1	4	6		
7	B	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
7	B	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
7	C	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		

- Molecule 8 is BACTERIOPHEOPHYTIN B (three-letter code: BPB) (formula: C<sub>55</sub>H<sub>74</sub>N<sub>4</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	1	Total	C	N	O	0	0
			65	55	4	6		
8	C	1	Total	C	N	O	0	0
			61	51	4	6		

- Molecule 9 is [(Z)-octadec-9-enyl] (2R)-2,3-bis(oxidanyl)propanoate (three-letter code: MPG) (formula: C<sub>21</sub>H<sub>40</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			25	21	4		

Continued on next page...


*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			25	21	4		
9	C	1	Total	C		0	0
			17	17			

- Molecule 10 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	C	1	Total	Fe	0	0
			1	1		


- Molecule 11 is MENAQUINONE-7 (three-letter code: MQ7) (formula: C<sub>46</sub>H<sub>64</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	C	1	Total	C	O	0	0
			48	46	2		


- Molecule 12 is 15-cis-1,2-dihydroneurosporene (three-letter code: NS5) (formula: C<sub>40</sub>H<sub>60</sub>).






Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	C	1	Total	C	0	0
			40	40		

- Molecule 13 is (2E,6E,10E,14E,18E,22E,26E)-3,7,11,15,19,23,27,31-OCTAMETHYLD OTRIACONTA-2,6,10,14,18,22,26,30-OCTAENYL TRIHYDROGEN DIPHOSPHATE (three-letter code: OTP) (formula:  $C_{40}H_{68}O_7P_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
13	C	1	Total	C	O	0	0
			41	40	1		

- Molecule 14 is PHOSPHATE ION (three-letter code: PO4) (formula:  $O_4P$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
14	C	1	Total	O	P	0	0
			5	4	1		
14	C	1	Total	O	P	0	0
			5	4	1		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Photosynthetic reaction center cytochrome c subunit




- Molecule 2: Reaction center protein L chain




- Molecule 3: Reaction center protein M chain



- Molecule 4: Reaction center protein H chain





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.60Å 82.90Å 382.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.50 – 3.60 50.50 – 3.60	Depositor EDS
% Data completeness (in resolution range)	99.4 (50.50-3.60) 99.5 (50.50-3.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.55 (at 3.57Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.242 , 0.280 0.242 , 0.280	Depositor DCC
$R_{free}$ test set	1094 reflections (4.97%)	DCC
Wilson B-factor (Å <sup>2</sup> )	73.9	Xtriage
Anisotropy	0.638	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 57.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	9890	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BCL, MPG, BPB, PO4, DGA, FE2, MQ7, HEC, OTP, FME, NS5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.28	0/2665	0.47	0/3633
2	B	0.29	0/2263	0.49	1/3089 (0.0%)
3	C	0.28	0/2650	0.46	0/3629
4	D	0.46	0/1804	0.79	8/2485 (0.3%)
All	All	0.32	0/9382	0.55	9/12836 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
4	D	0	6
All	All	0	7

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	78	VAL	C-N-CD	-7.68	103.70	120.60
4	D	142	ASP	CB-CG-OD2	7.19	124.77	118.30
4	D	185	LEU	CB-CG-CD2	6.27	121.65	111.00
4	D	192	ARG	NE-CZ-NH2	-6.11	117.25	120.30
4	D	183	LEU	CA-CB-CG	5.79	128.63	115.30

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	58	THR	Peptide
4	D	125	ASP	Peptide
4	D	140	ALA	Peptide
4	D	184	GLU	Peptide
4	D	192	ARG	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2598	0	2576	40	0
2	B	2170	0	2100	43	0
3	C	2546	0	2430	40	0
4	D	1771	0	1656	68	0
5	A	172	0	128	22	0
6	A	37	0	58	2	0
7	B	197	0	218	6	0
7	C	66	0	74	1	0
8	B	65	0	74	1	0
8	C	61	0	63	4	0
9	B	50	0	80	3	0
9	C	17	0	31	1	0
10	C	1	0	0	0	0
11	C	48	0	64	0	0
12	C	40	0	60	4	0
13	C	41	0	65	1	0
14	C	10	0	0	0	0
All	All	9890	0	9677	194	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 194 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:187:VAL:HB	4:D:192:ARG:HD3	1.49	0.93
4:D:184:GLU:HG2	4:D:195:LEU:HA	1.57	0.86

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:187:VAL:HG12	4:D:192:ARG:HB2	1.56	0.85
2:B:214:GLN:NE2	3:C:19:VAL:O	2.15	0.79
4:D:192:ARG:CG	4:D:193:THR:HA	2.12	0.79

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/356 (93%)	313 (95%)	17 (5%)	0	100	100
2	B	273/274 (100%)	257 (94%)	12 (4%)	4 (2%)	13	57
3	C	321/324 (99%)	301 (94%)	18 (6%)	2 (1%)	30	74
4	D	239/258 (93%)	213 (89%)	22 (9%)	4 (2%)	11	55
All	All	1163/1212 (96%)	1084 (93%)	69 (6%)	10 (1%)	21	67

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	59	TRP
2	B	204	ASP
4	D	97	GLU
4	D	141	THR
4	D	84	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was



analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/297 (94%)	279 (100%)	1 (0%)	93	98
2	B	218/219 (100%)	217 (100%)	1 (0%)	92	97
3	C	247/250 (99%)	245 (99%)	2 (1%)	86	95
4	D	167/212 (79%)	165 (99%)	2 (1%)	78	92
All	All	912/978 (93%)	906 (99%)	6 (1%)	88	96

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	194	PHE
4	D	228	ASP
3	C	214	PHE
2	B	54	SER
4	D	142	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	11	GLN
3	C	16	HIS
3	C	45	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	FME	D	1	4	8,9,10	0.84	0	5,9,11	0.87	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FME	D	1	4	-	0/6/9/11	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 1 is monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	HEC	A	401	1	24,50,50	2.53	6 (25%)	19,82,82	1.93	6 (31%)
5	HEC	A	402	1	24,50,50	2.46	5 (20%)	19,82,82	1.90	6 (31%)
5	HEC	A	403	1	24,50,50	2.52	6 (25%)	19,82,82	1.72	4 (21%)
5	HEC	A	404	1	24,50,50	2.47	6 (25%)	19,82,82	1.68	3 (15%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	DGA	A	405	-	36,36,43	1.21	3 (8%)	38,38,45	3.09	6 (15%)
7	BCL	B	301	-	54,73,74	1.18	3 (5%)	53,113,115	1.19	4 (7%)
7	BCL	B	302	-	55,74,74	1.15	3 (5%)	55,115,115	1.25	7 (12%)
7	BCL	B	303	-	55,74,74	1.12	2 (3%)	55,115,115	1.28	7 (12%)
8	BPB	B	304	-	62,70,70	1.22	4 (6%)	64,101,101	1.62	8 (12%)
9	MPG	B	305	-	23,24,24	1.26	1 (4%)	20,25,25	1.88	2 (10%)
9	MPG	B	306	-	23,24,24	1.27	1 (4%)	20,25,25	1.57	2 (10%)
7	BCL	C	401	-	55,74,74	1.28	3 (5%)	55,115,115	1.31	6 (10%)
8	BPB	C	402	-	58,66,70	1.38	4 (6%)	58,96,101	1.59	9 (15%)
11	MQ7	C	404	-	49,49,49	1.61	10 (20%)	63,63,63	1.71	14 (22%)
12	NS5	C	405	-	39,39,39	2.28	17 (43%)	44,46,46	2.43	15 (34%)
13	OTP	C	406	-	40,40,48	0.73	0	47,47,61	2.02	18 (38%)
9	MPG	C	407	-	16,16,24	0.78	0	15,15,25	0.76	0
14	PO4	C	408	-	4,4,4	0.61	0	6,6,6	0.23	0
14	PO4	C	409	-	4,4,4	0.65	0	6,6,6	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEC	A	401	1	-	0/6/54/54	0/0/8/8
5	HEC	A	402	1	-	0/6/54/54	0/0/8/8
5	HEC	A	403	1	-	0/6/54/54	0/0/8/8
5	HEC	A	404	1	-	0/6/54/54	0/0/8/8
6	DGA	A	405	-	-	0/37/37/45	0/0/0/0
7	BCL	B	301	-	-	0/36/136/137	0/0/9/9
7	BCL	B	302	-	-	0/37/137/137	0/0/9/9
7	BCL	B	303	-	-	0/37/137/137	0/0/9/9
8	BPB	B	304	-	-	0/46/105/105	0/1/6/6
9	MPG	B	305	-	-	0/25/25/25	0/0/0/0
9	MPG	B	306	-	-	0/25/25/25	0/0/0/0
7	BCL	C	401	-	-	0/37/137/137	0/0/9/9
8	BPB	C	402	-	-	0/42/101/105	0/1/6/6
11	MQ7	C	404	-	-	0/41/61/61	0/2/2/2
12	NS5	C	405	-	-	0/43/43/43	0/0/0/0
13	OTP	C	406	-	-	0/45/45/55	0/0/0/0
9	MPG	C	407	-	-	0/14/14/25	0/0/0/0
14	PO4	C	408	-	-	0/0/0/0	0/0/0/0

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
14	PO4	C	409	-	-	0/0/0/0	0/0/0/0

The worst 5 of 74 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	403	HEC	C3C-C2C	-5.93	1.34	1.40
5	A	401	HEC	C3C-C2C	-5.90	1.34	1.40
5	A	404	HEC	C3C-C2C	-5.79	1.34	1.40
5	A	401	HEC	C3B-C2B	-5.70	1.34	1.40
5	A	402	HEC	C3B-C2B	-5.49	1.35	1.40

The worst 5 of 117 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	405	DGA	OG2-CG2-CG3	-13.22	78.07	107.89
6	A	405	DGA	OG2-CG2-CG1	-9.78	83.73	105.90
12	C	405	NS5	C29-C30-C31	-7.55	118.42	127.69
12	C	405	NS5	C18-C17-C15	-6.47	117.81	127.22
8	B	304	BPB	C2C-C3C-C4C	-6.15	101.66	107.24

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

16 monomers are involved in 43 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	401	HEC	4	0
5	A	402	HEC	6	0
5	A	403	HEC	5	0
5	A	404	HEC	7	0
6	A	405	DGA	2	0
7	B	301	BCL	2	0
7	B	302	BCL	1	0
7	B	303	BCL	3	0
8	B	304	BPB	1	0
9	B	305	MPG	2	0
9	B	306	MPG	1	0
7	C	401	BCL	1	0
8	C	402	BPB	4	0
12	C	405	NS5	4	0
13	C	406	OTP	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	C	407	MPG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	332/356 (93%)	-0.20	4 (1%) 81 69	48, 63, 86, 116	0
2	B	273/274 (99%)	-0.28	1 (0%) 93 88	52, 78, 111, 135	0
3	C	323/324 (99%)	-0.33	1 (0%) 94 90	51, 77, 116, 141	0
4	D	242/258 (93%)	0.44	11 (4%) 37 26	63, 122, 153, 235	0
All	All	1170/1212 (96%)	-0.12	17 (1%) 76 64	48, 76, 134, 235	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	84	GLU	6.1
4	D	83	PRO	4.9
4	D	192	ARG	2.9
4	D	104	THR	2.9
4	D	82	ARG	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	FME	D	1	10/11	0.86	0.38	-	77,81,114,124	0

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	DGA	A	405	37/44	0.59	0.54	10.82	68,91,112,114	0
9	MPG	B	306	25/25	0.64	0.58	9.04	76,94,112,116	0
9	MPG	C	407	17/25	0.43	0.90	5.82	100,112,133,134	0
12	NS5	C	405	40/40	0.76	0.51	5.59	65,75,95,96	0
13	OTP	C	406	41/49	0.78	0.38	4.10	71,78,106,108	0
9	MPG	B	305	25/25	0.73	0.43	3.68	61,93,124,125	0
11	MQ7	C	404	48/48	0.87	0.38	2.84	60,87,93,94	0
8	BPB	B	304	65/65	0.92	0.27	1.77	57,71,79,80	0
7	BCL	C	401	66/66	0.90	0.27	1.65	54,64,72,74	0
8	BPB	C	402	61/65	0.88	0.34	1.55	72,80,100,104	0
7	BCL	B	302	66/66	0.93	0.24	1.45	48,63,70,71	0
5	HEC	A	403	43/43	0.96	0.28	1.41	30,54,56,56	0
7	BCL	B	301	65/66	0.91	0.27	1.16	37,62,166,173	0
14	PO4	C	408	5/5	0.61	0.40	1.00	118,118,119,121	0
5	HEC	A	404	43/43	0.94	0.23	0.49	32,59,62,62	0
5	HEC	A	402	43/43	0.96	0.27	0.42	35,54,61,72	0
5	HEC	A	401	43/43	0.96	0.32	0.12	45,74,77,80	0
7	BCL	B	303	66/66	0.94	0.21	-0.38	56,64,74,79	0
10	FE2	C	403	1/1	0.94	0.17	-1.16	74,74,74,74	0
14	PO4	C	409	5/5	0.85	0.34	-	124,125,126,126	0

### 6.5 Other polymers ⓘ

There are no such residues in this entry.