



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:45 PM GMT

PDB ID : 4MMR
Title : Crystal Structure of Prefusion-stabilized RSV F Variant Cav1 at pH 9.5
Authors : Stewart-Jones, G.B.E.; McLellan, J.S.; Joyce, M.G.; Sastry, M.; Yang, Y.;
Graham, B.S.; Kwong, P.D.
Deposited on : 2013-09-09
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1114 (3.14-3.06)
Clashscore	102246	1222 (3.14-3.06)
Ramachandran outliers	100387	1174 (3.14-3.06)
Sidechain outliers	100360	1174 (3.14-3.06)
RSRZ outliers	91569	1119 (3.14-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	82	<div> <div>15%</div> <div>59%</div> <div>23%</div> <div>11%</div> <div>• 5%</div> </div>
2	B	414	<div> <div>2%</div> <div>65%</div> <div>20%</div> <div>5%</div> <div>• 9%</div> </div>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3523 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fusion glycoprotein F2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	78	Total	C	N	O	S	0	0	0
			612	385	100	124	3			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	ALA	PRO	ENGINEERED MUTATION	UNP P03420

- Molecule 2 is a protein called Fusion glycoprotein F1 fused with Fibrin trimerization domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	377	Total	C	N	O	S	0	0	0
			2911	1841	480	572	18			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	190	PHE	SER	ENGINEERED MUTATION	UNP P03420
B	207	LEU	VAL	ENGINEERED MUTATION	UNP P03420
B	379	VAL	ILE	ENGINEERED MUTATION	UNP P03420
B	447	VAL	MET	ENGINEERED MUTATION	UNP P03420
B	514	SER	-	LINKER	UNP P03420
B	515	ALA	-	LINKER	UNP P03420
B	516	ILE	-	LINKER	UNP P03420
B	517	GLY	-	LINKER	UNP P03420
B	539	LEU	PHE	VIRIANT	UNP P10104
B	545	GLY	-	EXPRESSION TAG	UNP P10104
B	546	GLY	-	EXPRESSION TAG	UNP P10104
B	547	LEU	-	EXPRESSION TAG	UNP P10104
B	548	VAL	-	EXPRESSION TAG	UNP P10104
B	549	PRO	-	EXPRESSION TAG	UNP P10104

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	550	ARG	-	EXPRESSION TAG	UNP P10104


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fusion glycoprotein F2



- Molecule 2: Fusion glycoprotein F1 fused with Fibrin trimerization domain



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 3 2	Depositor
Cell constants a, b, c, α , β , γ	170.85Å 170.85Å 170.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.66 – 3.10 47.39 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.1 (45.66-3.10) 99.1 (47.39-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 3.12Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.219 , 0.264 0.209 , 0.248	Depositor DCC
R_{free} test set	796 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	98.4	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 82.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 15885 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3523	wwPDB-VP
Average B, all atoms (Å ²)	124.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/619	0.75	0/836
2	B	0.60	0/2955	0.81	2/4007 (0.0%)
All	All	0.60	0/3574	0.80	2/4843 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	291	ILE	CB-CA-C	-5.80	100.00	111.60
2	B	334	LEU	CA-CB-CG	5.15	127.15	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	182	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	612	0	620	18	0
2	B	2911	0	2947	51	1
All	All	3523	0	3567	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:361:GLN:O	2:B:362:SER:OG	2.10	0.69
2:B:506:ILE:HG22	2:B:507:ARG:N	2.09	0.68
1:A:70:ASN:N	2:B:212:CYS:SG	2.68	0.66
2:B:397:THR:OG1	2:B:487:GLU:HB2	1.94	0.66
1:A:97:MET:SD	1:A:98:GLN:HG3	2.36	0.66
1:A:63:ASN:N	1:A:63:ASN:OD1	2.31	0.62
2:B:229:ARG:HH21	2:B:253:THR:HG23	1.67	0.59
2:B:178:VAL:HG12	2:B:178:VAL:O	2.02	0.59
2:B:267:THR:OG1	2:B:268:ASN:N	2.36	0.58
2:B:279:GLN:OE1	2:B:279:GLN:N	2.35	0.58
2:B:429:ARG:HB2	2:B:429:ARG:CZ	2.34	0.58
2:B:167:ILE:HG23	2:B:189:THR:HG21	1.85	0.58
2:B:427:LYS:HD3	2:B:428:ASN:N	2.21	0.55
1:A:78:LEU:CD2	2:B:220:VAL:HG11	2.37	0.55
2:B:245:THR:HA	2:B:246:PRO:C	2.28	0.54
1:A:97:MET:HB3	1:A:98:GLN:HE21	1.72	0.53
2:B:433:LYS:HE3	2:B:435:PHE:CE1	2.44	0.53
2:B:478:TYR:O	2:B:480:PRO:HD3	2.08	0.53
2:B:334:LEU:HD23	2:B:334:LEU:C	2.29	0.52
1:A:26:GLN:OE1	1:A:26:GLN:O	2.26	0.52
1:A:69:CYS:SG	1:A:70:ASN:N	2.83	0.51
2:B:506:ILE:CG2	2:B:507:ARG:N	2.75	0.50
2:B:202:GLN:NE2	2:B:202:GLN:O	2.45	0.49
2:B:258:LEU:HA	2:B:261:ILE:HD12	1.95	0.49
1:A:82:GLU:HG2	2:B:224:GLN:HA	1.94	0.49
1:A:35:SER:HB2	1:A:36:THR:HG23	1.94	0.49
2:B:293:LYS:O	2:B:296:VAL:HG12	2.14	0.48
1:A:80:LYS:HD3	1:A:81:GLN:N	2.28	0.48
2:B:368:ASP:OD1	2:B:368:ASP:C	2.52	0.48
1:A:76:VAL:HA	1:A:79:ILE:HG22	1.95	0.47
2:B:163:GLU:CD	2:B:182:SER:H	2.18	0.47
2:B:267:THR:HG23	2:B:270:GLN:HG3	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:293:LYS:O	2:B:294:GLU:C	2.53	0.47
2:B:240:ASN:O	2:B:241:ALA:C	2.53	0.47
2:B:202:GLN:C	2:B:202:GLN:HE21	2.19	0.46
1:A:78:LEU:HD22	2:B:220:VAL:HG11	1.96	0.46
2:B:138:LEU:HB3	2:B:141:LEU:HD12	1.98	0.46
2:B:223:PHE:HA	2:B:226:LYS:HB2	1.97	0.46
2:B:408:THR:OG1	2:B:411:GLY:N	2.49	0.46
2:B:398:SER:HA	2:B:485:SER:O	2.16	0.46
2:B:428:ASN:HB2	2:B:429:ARG:NH1	2.31	0.45
1:A:71:GLY:N	2:B:212:CYS:SG	2.90	0.45
2:B:198:TYR:C	2:B:198:TYR:CD1	2.89	0.45
2:B:293:LYS:O	2:B:295:GLU:N	2.50	0.45
1:A:83:LEU:HA	1:A:86:TYR:HB3	1.99	0.45
2:B:318:THR:O	2:B:339:ARG:NH1	2.51	0.44
2:B:254:ASN:H	2:B:254:ASN:ND2	2.16	0.43
2:B:210:GLN:HA	2:B:212:CYS:HA	2.01	0.43
1:A:86:TYR:C	1:A:86:TYR:CD1	2.92	0.43
2:B:503:LEU:O	2:B:507:ARG:HG3	2.19	0.43
2:B:491:SER:OG	2:B:494:GLN:HB2	2.18	0.42
1:A:63:ASN:C	1:A:64:ILE:HG12	2.40	0.42
2:B:497:GLU:HG2	2:B:498:LYS:N	2.34	0.42
2:B:183:ASN:OD1	2:B:185:VAL:HG22	2.20	0.42
2:B:507:ARG:HB2	2:B:507:ARG:CZ	2.49	0.42
1:A:73:ASP:HB3	2:B:214:ILE:HB	2.00	0.42
2:B:254:ASN:O	2:B:258:LEU:HD12	2.20	0.42
1:A:61:LEU:O	2:B:295:GLU:CB	2.68	0.42
2:B:254:ASN:O	2:B:255:SER:C	2.58	0.41
2:B:183:ASN:HD21	2:B:185:VAL:HG13	1.85	0.40
2:B:512:LEU:HD23	2:B:513:LEU:HD22	2.03	0.40
2:B:152:VAL:O	2:B:153:ALA:C	2.59	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:213:SER:OG	2:B:215:SER:O[14_444]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	76/82 (93%)	56 (74%)	15 (20%)	5 (7%)	1	9
2	B	375/414 (91%)	306 (82%)	57 (15%)	12 (3%)	5	26
All	All	451/496 (91%)	362 (80%)	72 (16%)	17 (4%)	4	22

All (17) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	64	ILE
2	B	202	GLN
2	B	294	GLU
1	A	74	ALA
1	A	83	LEU
2	B	206	ILE
2	B	213	SER
2	B	267	THR
2	B	510	ASP
1	A	97	MET
2	B	172	LEU
2	B	255	SER
2	B	382	CYS
2	B	241	ALA
1	A	55	SER
2	B	355	ALA
2	B	217	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	70/73 (96%)	52 (74%)	18 (26%)	0	2
2	B	345/373 (92%)	294 (85%)	51 (15%)	4	16
All	All	415/446 (93%)	346 (83%)	69 (17%)	3	12

All (69) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	29	THR
1	A	35	SER
1	A	58	THR
1	A	63	ASN
1	A	66	GLU
1	A	68	LYS
1	A	69	CYS
1	A	72	THR
1	A	75	LYS
1	A	77	LYS
1	A	79	ILE
1	A	80	LYS
1	A	81	GLN
1	A	83	LEU
1	A	88	ASN
1	A	97	MET
1	A	98	GLN
2	B	144	VAL
2	B	165	ASN
2	B	168	LYS
2	B	172	LEU
2	B	176	LYS
2	B	180	SER
2	B	183	ASN
2	B	187	VAL
2	B	189	THR
2	B	190	PHE
2	B	201	LYS
2	B	202	GLN
2	B	204	LEU
2	B	206	ILE
2	B	207	LEU
2	B	210	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	216	ASN
2	B	219	THR
2	B	259	SER
2	B	260	LEU
2	B	267	THR
2	B	279	GLN
2	B	291	ILE
2	B	292	ILE
2	B	311	THR
2	B	319	SER
2	B	327	LYS
2	B	334	LEU
2	B	348	SER
2	B	354	GLN
2	B	356	GLU
2	B	359	LYS
2	B	361	GLN
2	B	365	VAL
2	B	380	ASN
2	B	381	LEU
2	B	384	VAL
2	B	398	SER
2	B	423	THR
2	B	427	LYS
2	B	429	ARG
2	B	449	THR
2	B	466	SER
2	B	472	GLU
2	B	481	LEU
2	B	485	SER
2	B	501	GLN
2	B	503	LEU
2	B	506	ILE
2	B	510	ASP
2	B	513	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	GLN
1	A	98	GLN
2	B	202	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	254	ASN
2	B	277	ASN
2	B	354	GLN
2	B	380	ASN
2	B	501	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	78/82 (95%)	0.45	12 (15%) 3 1	82, 134, 237, 245	0
2	B	377/414 (91%)	-0.32	9 (2%) 62 39	68, 105, 185, 306	0
All	All	455/496 (91%)	-0.19	21 (4%) 36 17	68, 107, 203, 306	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	211	SER	7.3
1	A	67	ASN	5.9
1	A	70	ASN	4.9
1	A	99	SER	4.4
2	B	212	CYS	4.2
1	A	26	GLN	4.0
2	B	208	ASN	4.0
1	A	101	PRO	3.6
1	A	68	LYS	3.5
1	A	102	ALA	3.5
1	A	66	GLU	3.3
1	A	73	ASP	3.0
2	B	511	GLU	2.9
2	B	213	SER	2.7
2	B	363	ASN	2.7
2	B	205	PRO	2.5
1	A	69	CYS	2.5
2	B	204	LEU	2.5
1	A	103	THR	2.4
1	A	64	ILE	2.2
2	B	209	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.