



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 13, 2016 – 01:37 PM EST

PDB ID : 5MN6  
Title : S. aureus FtsZ 12-316 F138A GDP Closed form (3FCm)  
Authors : Wagstaff, J.M.; Tsim, M.; Kureisaite-Ciziene, D.; Lowe, J.  
Deposited on : 2016-12-12  
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028442  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028442


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1124 (3.24-3.16)
Clashscore	102246	1024 (3.22-3.18)
Ramachandran outliers	100387	1004 (3.22-3.18)
Sidechain outliers	100360	1003 (3.22-3.18)
RSRZ outliers	91569	1129 (3.24-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	305	 77% 18% . .
1	B	305	 5% 79% 19% ..

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4374 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cell division protein FtsZ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	296	Total	C	N	O	S	0	0	0
			2136	1321	368	435	12			
1	B	303	Total	C	N	O	S	0	0	0
			2182	1348	376	446	12			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	138	ALA	PHE	engineered mutation	UNP P0A031
B	138	ALA	PHE	engineered mutation	UNP P0A031

- Molecule 2 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula:  $C_{10}H_{15}N_5O_{11}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	41.12Å 68.15Å 207.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.20 41.32 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.5 (30.00-3.20) 99.7 (41.32-3.20)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.88 (at 3.18Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, $R_{free}$	0.216 , 0.300 0.221 , 0.216	Depositor DCC
$R_{free}$ test set	518 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	105.0	Xtriage
Anisotropy	0.432	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 66.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4374	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	115.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	0/2150	0.78	0/2900
1	B	0.51	0/2197	0.75	0/2966
All	All	0.52	0/4347	0.76	0/5866

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2136	0	2165	45	2
1	B	2182	0	2209	39	0
2	A	28	0	12	2	0
2	B	28	0	12	2	0
All	All	4374	0	4398	84	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:24:ASN:O	1:A:28:ASN:OD1	1.53	1.27
1:B:65:LEU:HD21	1:B:84:GLU:OE1	1.48	1.10
1:B:60:GLN:NE2	1:B:68:GLY:HA2	1.74	1.03
1:B:47:GLY:N	1:B:68:GLY:O	1.99	0.94
1:A:45:THR:O	1:A:69:LEU:O	1.93	0.87
1:A:45:THR:HG21	1:A:71:ALA:HB2	1.59	0.85
1:B:71:ALA:HB1	1:B:77:ILE:HB	1.59	0.84
1:B:60:GLN:HE21	1:B:68:GLY:HA2	1.39	0.83
1:A:45:THR:OG1	1:A:71:ALA:N	2.13	0.81
1:A:46:ASP:N	1:A:70:GLY:HA2	1.99	0.77
1:A:168:ARG:CZ	1:A:248:PRO:O	2.33	0.76
1:A:185:GLU:OE2	1:A:188:ASN:ND2	2.17	0.75
1:A:258:GLN:HE22	1:A:314:GLY:HA3	1.53	0.74
1:B:258:GLN:HE22	1:B:314:GLY:HA3	1.51	0.74
1:A:50:LEU:HD22	1:A:58:LYS:HB3	1.70	0.74
1:B:50:LEU:HD22	1:B:58:LYS:HB3	1.71	0.72
1:A:71:ALA:O	1:A:77:ILE:HD13	1.90	0.72
1:A:136:PHE:HB2	1:A:139:GLU:HG3	1.76	0.67
1:B:71:ALA:O	1:B:77:ILE:HD12	1.95	0.67
1:B:136:PHE:HB2	1:B:139:GLU:HG3	1.78	0.64
1:B:45:THR:O	1:B:69:LEU:O	2.14	0.63
1:A:45:THR:C	1:A:70:GLY:HA2	2.19	0.63
1:B:45:THR:C	1:B:69:LEU:O	2.37	0.62
1:A:60:GLN:HE21	1:A:68:GLY:HA2	1.64	0.62
1:A:134:ARG:HB2	1:A:165:PRO:HA	1.82	0.61
1:B:264:ILE:HD13	1:B:275:ALA:HA	1.82	0.60
1:B:134:ARG:HB2	1:B:165:PRO:HA	1.82	0.60
1:B:258:GLN:NE2	1:B:314:GLY:HA3	2.16	0.60
1:A:185:GLU:CD	1:A:188:ASN:HD21	2.04	0.60
1:B:164:ILE:HG21	1:B:189:VAL:HG11	1.84	0.59
1:A:14:LEU:HB2	1:A:38:VAL:HG22	1.84	0.59
1:A:258:GLN:NE2	1:A:314:GLY:HA3	2.17	0.59
1:A:168:ARG:NH1	1:A:250:LEU:O	2.36	0.59
1:A:154:MET:O	1:A:158:VAL:HG22	2.03	0.59
1:B:14:LEU:HB2	1:B:38:VAL:HG22	1.85	0.58
1:B:154:MET:O	1:B:158:VAL:HG22	2.04	0.58
1:B:93:ILE:HB	1:B:124:MET:CE	2.36	0.56
1:A:93:ILE:HB	1:A:124:MET:CE	2.36	0.56
1:B:65:LEU:HG	1:B:84:GLU:OE2	2.06	0.56
1:B:46:ASP:HA	1:B:69:LEU:C	2.27	0.55
1:A:168:ARG:NH1	1:A:248:PRO:O	2.41	0.52
1:A:56:GLU:N	1:A:56:GLU:OE1	2.42	0.52

Continued on next page...



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:104:GLY:HA2	1:A:133:THR:HG22	1.92	0.52
1:A:135:PRO:O	1:A:144:GLN:NE2	2.43	0.52
1:B:200:LEU:HB3	1:B:209:LEU:HD13	1.92	0.52
1:A:188:ASN:OD1	1:A:189:VAL:N	2.44	0.51
1:B:104:GLY:HA2	1:B:133:THR:HG22	1.91	0.51
1:B:108:GLY:N	2:B:401:GDP:O3B	2.44	0.49
1:B:65:LEU:HD21	1:B:84:GLU:CD	2.27	0.49
1:A:60:GLN:NE2	1:A:68:GLY:HA2	2.28	0.49
1:A:185:GLU:HA	1:A:188:ASN:ND2	2.28	0.48
1:A:46:ASP:CA	1:A:70:GLY:HA2	2.42	0.48
1:A:185:GLU:HA	1:A:188:ASN:CG	2.35	0.48
1:B:230:VAL:HG22	1:B:307:VAL:HG22	1.95	0.47
1:B:15:LYS:HG2	1:B:39:GLU:HB2	1.96	0.47
1:A:28:ASN:N	1:A:28:ASN:OD1	2.46	0.47
1:A:200:LEU:HB3	1:A:209:LEU:HD13	1.97	0.47
1:A:15:LYS:HG2	1:A:39:GLU:HB2	1.96	0.46
1:B:75:PRO:HB3	1:B:111:THR:HB	1.97	0.45
1:A:225:LEU:HD22	1:A:251:GLU:H	1.81	0.45
1:A:45:THR:CB	1:A:71:ALA:N	2.79	0.45
1:B:14:LEU:HD22	1:B:98:MET:HB3	1.98	0.45
1:A:214:VAL:O	1:A:218:MET:HG2	2.17	0.45
1:B:181:GLU:HG2	1:B:184:LYS:HD2	1.99	0.44
1:B:258:GLN:HE22	1:B:314:GLY:CA	2.27	0.44
1:B:93:ILE:HB	1:B:124:MET:HE2	1.99	0.44
1:B:214:VAL:O	1:B:218:MET:HG2	2.16	0.44
1:A:64:LYS:HD2	1:A:84:GLU:CD	2.39	0.43
1:A:183:PHE:HB3	2:A:401:GDP:C6	2.53	0.43
1:A:237:ALA:HA	1:A:306:ILE:HD11	2.00	0.43
1:A:108:GLY:N	2:A:401:GDP:O3B	2.51	0.43
1:B:242:LYS:HE2	1:B:242:LYS:HB3	1.80	0.43
1:B:139:GLU:OE2	2:B:401:GDP:O2'	2.36	0.43
1:B:62:GLY:O	1:B:63:GLU:C	2.57	0.43
1:A:185:GLU:CD	1:A:188:ASN:ND2	2.69	0.42
1:A:45:THR:CG2	1:A:71:ALA:HB2	2.39	0.42
1:A:168:ARG:NE	1:A:248:PRO:O	2.51	0.42
1:A:61:ILE:HG22	1:A:85:SER:HB2	2.01	0.42
1:A:14:LEU:HD22	1:A:98:MET:HB3	2.02	0.42
1:B:208:ASN:C	1:B:209:LEU:HD12	2.40	0.41
1:A:210:ASP:O	1:A:213:ASP:HB2	2.21	0.41
1:B:188:ASN:OD1	1:B:191:ARG:NH2	2.54	0.40
1:B:212:ALA:O	1:B:216:THR:OG1	2.40	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:302:LEU:HD22	1:B:305:GLU:HG3	2.02	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:GLU:OE1	1:A:180:MET:CE[4_555]	1.60	0.60
1:A:84:GLU:CD	1:A:180:MET:CE[4_555]	2.12	0.08

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/305 (96%)	282 (97%)	10 (3%)	0	100	100
1	B	301/305 (99%)	279 (93%)	19 (6%)	3 (1%)	19	65
All	All	593/610 (97%)	561 (95%)	29 (5%)	3 (0%)	34	78

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	63	GLU
1	B	205	GLY
1	B	208	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/230 (97%)	213 (95%)	11 (5%)	31	72
1	B	229/230 (100%)	221 (96%)	8 (4%)	43	80
All	All	453/460 (98%)	434 (96%)	19 (4%)	36	75

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	ASN
1	A	37	ASN
1	A	56	GLU
1	A	64	LYS
1	A	74	ASN
1	A	133	THR
1	A	177	THR
1	A	180	MET
1	A	216	THR
1	A	223	SER
1	A	272	LEU
1	B	37	ASN
1	B	64	LYS
1	B	80	LYS
1	B	133	THR
1	B	180	MET
1	B	185	GLU
1	B	216	THR
1	B	272	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	GLN
1	A	258	GLN
1	B	60	GLN
1	B	258	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GDP	A	401	-	24,30,30	1.29	2 (8%)	26,47,47	2.17	6 (23%)
2	GDP	B	401	-	24,30,30	1.20	3 (12%)	26,47,47	2.30	7 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GDP	A	401	-	-	0/12/32/32	0/3/3/3
2	GDP	B	401	-	-	0/12/32/32	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	GDP	C2'-C1'	-2.24	1.50	1.53
2	B	401	GDP	C5-C4	3.09	1.47	1.40
2	B	401	GDP	C6-C5	3.50	1.48	1.41
2	A	401	GDP	C6-C5	3.62	1.48	1.41
2	A	401	GDP	C5-C4	3.74	1.48	1.40

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	GDP	C5-C6-N1	-5.53	116.29	123.52
2	B	401	GDP	C6-C5-C4	-4.31	115.93	120.86
2	B	401	GDP	C5-C6-N1	-4.01	118.28	123.52
2	B	401	GDP	N3-C2-N1	-4.01	122.10	127.56
2	B	401	GDP	C1'-N9-C4	-3.63	122.75	126.81
2	A	401	GDP	N3-C2-N1	-3.15	123.27	127.56
2	A	401	GDP	C6-C5-C4	-2.81	117.64	120.86
2	B	401	GDP	C2'-C1'-N9	-2.27	107.39	113.47
2	B	401	GDP	O2B-PB-O1B	2.03	117.25	110.63
2	A	401	GDP	O2B-PB-O1B	2.19	117.78	110.63
2	A	401	GDP	C4'-O4'-C1'	2.93	112.75	109.64
2	B	401	GDP	C6-N1-C2	6.40	123.39	115.88
2	A	401	GDP	C6-N1-C2	6.69	123.72	115.88

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	GDP	2	0
2	B	401	GDP	2	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	296/305 (97%)	-0.09	0 100 100	59, 101, 167, 253	0
1	B	303/305 (99%)	0.17	15 (4%) 32 19	68, 113, 172, 253	0
All	All	599/610 (98%)	0.04	15 (2%) 61 47	59, 109, 170, 253	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	204	SER	6.8
1	B	203	VAL	3.7
1	B	103	SER	3.2
1	B	128	THR	2.8
1	B	70	GLY	2.8
1	B	102	THR	2.6
1	B	205	GLY	2.5
1	B	82	ALA	2.5
1	B	89	ILE	2.3
1	B	105	MET	2.2
1	B	303	GLN	2.2
1	B	306	ILE	2.2
1	B	42	ALA	2.1
1	B	202	ALA	2.1
1	B	80	LYS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GDP	A	401	28/28	0.92	0.19	-0.41	75,89,111,112	0
2	GDP	B	401	28/28	0.94	0.15	-0.62	84,98,114,125	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.