



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 08:54 PM GMT

PDB ID : 1MO2  
Title : Thioesterase Domain from 6-Deoxyerythronolide Synthase (DEBS TE), pH 8.5  
Authors : Tsai, S.-C.; Lu, H.; Cane, D.E.; Khosla, C.; Stroud, R.M.  
Deposited on : 2002-09-05  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	 2% 40% 37% 7% • 15%
1	B	294	 3% 43% 33% 8% • 15%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3830 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Erythronolide synthase, modules 5 and 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	S	0	0	0
			1887	1192	329	357	9			
1	B	251	Total	C	N	O	S	0	0	0
			1887	1192	329	357	9			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	INSERTION	UNP Q03133
A	2	ALA	-	INSERTION	UNP Q03133
A	3	SER	-	INSERTION	UNP Q03133
A	284	SER	-	INSERTION	UNP Q03133
A	285	SER	-	INSERTION	UNP Q03133
A	286	VAL	-	INSERTION	UNP Q03133
A	287	ASP	-	INSERTION	UNP Q03133
A	288	LYS	-	INSERTION	UNP Q03133
A	289	LEU	-	INSERTION	UNP Q03133
A	290	ALA	-	INSERTION	UNP Q03133
A	291	ALA	-	INSERTION	UNP Q03133
A	292	ALA	-	INSERTION	UNP Q03133
A	293	LEU	-	INSERTION	UNP Q03133
A	294	GLU	-	INSERTION	UNP Q03133
B	1	MET	-	INSERTION	UNP Q03133
B	2	ALA	-	INSERTION	UNP Q03133
B	3	SER	-	INSERTION	UNP Q03133
B	284	SER	-	INSERTION	UNP Q03133
B	285	SER	-	INSERTION	UNP Q03133
B	286	VAL	-	INSERTION	UNP Q03133
B	287	ASP	-	INSERTION	UNP Q03133
B	288	LYS	-	INSERTION	UNP Q03133
B	289	LEU	-	INSERTION	UNP Q03133
B	290	ALA	-	INSERTION	UNP Q03133
B	291	ALA	-	INSERTION	UNP Q03133

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	292	ALA	-	INSERTION	UNP Q03133
B	293	LEU	-	INSERTION	UNP Q03133
B	294	GLU	-	INSERTION	UNP Q03133

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	29	Total O 29 29	0	0
2	B	27	Total O 27 27	0	0



- Molecule 1: Erythronolide synthase, modules 5 and 6



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.50Å 102.50Å 156.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.00 49.22 – 3.00	Depositor EDS
% Data completeness (in resolution range)	89.6 (30.00-3.00) 84.2 (49.22-3.00)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.36 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.1	Depositor
R, $R_{free}$	0.242 , 0.287 0.253 , 0.299	Depositor DCC
$R_{free}$ test set	1082 reflections (5.08%)	DCC
Wilson B-factor (Å <sup>2</sup> )	67.2	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 61.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 23833 reflections	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	3830	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/1939	0.74	2/2651 (0.1%)
1	B	0.46	0/1939	0.73	1/2651 (0.0%)
All	All	0.47	0/3878	0.74	3/5302 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	260	PHE	N-CA-C	7.11	130.19	111.00
1	A	259	HIS	N-CA-C	5.28	125.24	111.00
1	B	266	HIS	N-CA-C	-5.06	97.35	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1887	0	1810	124	1
1	B	1887	0	1810	111	1
2	A	29	0	0	3	0
2	B	27	0	0	4	0
All	All	3830	0	3620	231	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

The worst 5 of 231 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:232:GLU:HB3	1:A:233:PRO:HD2	1.26	1.08
1:A:71:ILE:HD13	1:A:136:PHE:HB2	1.37	1.06
1:B:63:ASP:HA	1:B:97:PRO:HA	1.52	0.91
1:A:170:VAL:HG12	1:A:171:TYR:H	1.32	0.91
1:B:232:GLU:CB	1:B:233:PRO:HD2	2.00	0.91

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:THR:N	1:B:259:HIS:N[2_564]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/294 (84%)	194 (78%)	32 (13%)	21 (8%)	1	5
1	B	247/294 (84%)	201 (81%)	25 (10%)	21 (8%)	1	5
All	All	494/588 (84%)	395 (80%)	57 (12%)	42 (8%)	1	5

5 of 42 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	80	SER
1	A	233	PRO
1	A	240	ASP
1	A	256	PRO
1	A	259	HIS



### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/224 (86%)	167 (87%)	25 (13%)	5	22
1	B	192/224 (86%)	169 (88%)	23 (12%)	6	26
All	All	384/448 (86%)	336 (88%)	48 (12%)	6	24

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	264	GLN
1	B	42	SER
1	B	262	MET
1	A	266	HIS
1	A	273	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	266	HIS
1	B	123	GLN
1	B	259	HIS
1	A	264	GLN
1	B	180	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	251/294 (85%)	-0.26	5 (1%) 68 39	34, 52, 94, 120	0
1	B	251/294 (85%)	-0.26	8 (3%) 51 23	35, 51, 91, 123	0
All	All	502/588 (85%)	-0.26	13 (2%) 59 29	34, 52, 94, 123	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	240	ASP	3.8
1	A	237	TRP	3.7
1	A	240	ASP	3.6
1	B	238	PRO	3.5
1	A	259	HIS	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.