



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:52 PM GMT

PDB ID : 4N1I  
Title : Crystal Structure of the alpha-L-arabinofuranosidase UmAbf62A from  
Ustilago maidys  
Authors : Siguier, B.; Dumon, C.; Mourey, L.; Tranier, S.  
Deposited on : 2013-10-04  
Resolution : 1.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1235 (1.10-0.90)
Clashscore	102246	1333 (1.10-0.90)
Ramachandran outliers	100387	1247 (1.10-0.90)
Sidechain outliers	100360	1246 (1.10-0.90)
RSRZ outliers	91569	1239 (1.10-0.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	402	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	403[A]	-	-	X	X
3	GOL	A	403[B]	-	-	-	X
3	GOL	A	403[C]	-	-	-	X
3	GOL	A	404	-	-	-	X
3	GOL	A	405[A]	-	-	-	X
3	GOL	A	405[B]	-	-	-	X
4	TRS	A	406	-	-	X	X
5	1PE	A	407	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3167 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alpha-L-arabinofuranosidase UmAbf62A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	312	Total	C	N	O	S	8	29	0
			2631	1684	436	503	8			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	312	GLY	-	EXPRESSION TAG	UNP Q4P6F4
A	313	LEU	-	EXPRESSION TAG	UNP Q4P6F4
A	314	GLU	-	EXPRESSION TAG	UNP Q4P6F4
A	315	GLN	-	EXPRESSION TAG	UNP Q4P6F4
A	316	LYS	-	EXPRESSION TAG	UNP Q4P6F4
A	317	LEU	-	EXPRESSION TAG	UNP Q4P6F4
A	318	ILE	-	EXPRESSION TAG	UNP Q4P6F4
A	319	SER	-	EXPRESSION TAG	UNP Q4P6F4
A	320	GLU	-	EXPRESSION TAG	UNP Q4P6F4
A	321	GLU	-	EXPRESSION TAG	UNP Q4P6F4
A	322	ASP	-	EXPRESSION TAG	UNP Q4P6F4
A	323	LEU	-	EXPRESSION TAG	UNP Q4P6F4
A	324	ASN	-	EXPRESSION TAG	UNP Q4P6F4
A	325	SER	-	EXPRESSION TAG	UNP Q4P6F4
A	326	ALA	-	EXPRESSION TAG	UNP Q4P6F4
A	327	VAL	-	EXPRESSION TAG	UNP Q4P6F4
A	328	ASP	-	EXPRESSION TAG	UNP Q4P6F4
A	329	HIS	-	EXPRESSION TAG	UNP Q4P6F4
A	330	HIS	-	EXPRESSION TAG	UNP Q4P6F4
A	331	HIS	-	EXPRESSION TAG	UNP Q4P6F4
A	332	HIS	-	EXPRESSION TAG	UNP Q4P6F4
A	333	HIS	-	EXPRESSION TAG	UNP Q4P6F4
A	334	HIS	-	EXPRESSION TAG	UNP Q4P6F4

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	1
			18	9	9		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	1
			12	6	6		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula:  $C_4H_{12}NO_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 5 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula:  $C_{10}H_{22}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	478	Total 478	O 478	0	0



- Molecule 1: alpha-L-arabinofuranosidase UmAbf62A





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.28Å 65.90Å 68.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.30 – 1.00 37.28 – 1.00	Depositor EDS
% Data completeness (in resolution range)	99.5 (37.30-1.00) 99.5 (37.28-1.00)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.65 (at 1.00Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.114 , 0.139 0.115 , 0.140	Depositor DCC
$R_{free}$ test set	7283 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	5.6	Xtriage
Anisotropy	0.276	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 43.4	EDS
Estimated twinning fraction	0.019 for -h,l,k	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 145654 reflections	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	3167	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, TRS, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.05	6/2745 (0.2%)	1.12	23/3741 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	254	SER	CB-OG	6.32	1.50	1.42
1	A	320	GLU	CD-OE2	-6.25	1.18	1.25
1	A	314	GLU	CD-OE2	5.91	1.32	1.25
1	A	58[A]	TYR	CG-CD1	-5.38	1.32	1.39
1	A	58[B]	TYR	CG-CD1	-5.38	1.32	1.39

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	17	ARG	NE-CZ-NH1	9.77	125.19	120.30
1	A	57[A]	LYS	CD-CE-NZ	9.59	133.75	111.70
1	A	57[B]	LYS	CD-CE-NZ	9.59	133.75	111.70
1	A	155	LEU	CB-CG-CD1	-8.48	96.58	111.00
1	A	17	ARG	NE-CZ-NH2	-7.53	116.54	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	266	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2631	0	2553	62	0
2	A	1	0	0	0	0
3	A	42	0	51	18	0
4	A	8	0	12	14	0
5	A	7	0	9	5	0
6	A	478	0	0	33	1
All	All	3167	0	2625	74	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 74 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215[A]:GLN:NE2	4:A:406:TRS:O2	1.65	1.28
3:A:403[C]:GOL:H2	6:A:1234:HOH:O	1.00	1.16
1:A:279[B]:ARG:NH1	6:A:1047:HOH:O	1.77	1.11
3:A:403[B]:GOL:H2	6:A:1234:HOH:O	0.82	0.99
1:A:220:ARG:HH21	4:A:406:TRS:H31	1.29	0.96

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:1260:HOH:O	6:A:1261:HOH:O[2_664]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/334 (102%)	326 (96%)	14 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/279 (103%)	287 (100%)	1 (0%)	94	74

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	294	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	HIS
1	A	68	HIS
1	A	121	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	GOL	A	402	-	5,5,5	1.89	2 (40%)	5,5,5	0.67	0
3	GOL	A	403[A]	-	5,5,5	0.70	0	5,5,5	0.32	0
3	GOL	A	403[B]	-	5,5,5	3.27	1 (20%)	5,5,5	1.64	1 (20%)
3	GOL	A	403[C]	-	5,5,5	2.33	1 (20%)	5,5,5	1.52	1 (20%)
3	GOL	A	404	-	5,5,5	0.63	0	5,5,5	1.62	1 (20%)
3	GOL	A	405[A]	-	5,5,5	1.20	0	5,5,5	1.41	0
3	GOL	A	405[B]	-	5,5,5	2.28	2 (40%)	5,5,5	2.13	3 (60%)
4	TRS	A	406	-	7,7,7	0.97	0	9,9,9	2.12	3 (33%)
5	1PE	A	407	-	6,6,15	0.63	0	5,5,14	2.17	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	402	-	-	0/4/4/4	0/0/0/0
3	GOL	A	403[A]	-	-	0/4/4/4	0/0/0/0
3	GOL	A	403[B]	-	-	0/4/4/4	0/0/0/0
3	GOL	A	403[C]	-	-	0/4/4/4	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	404	-	-	0/4/4/4	0/0/0/0
3	GOL	A	405[A]	-	-	0/4/4/4	0/0/0/0
3	GOL	A	405[B]	-	-	0/4/4/4	0/0/0/0
4	TRS	A	406	-	-	0/9/9/9	0/0/0/0
5	1PE	A	407	-	-	0/4/4/13	0/0/0/0

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	403[B]	GOL	O2-C2	-6.93	1.22	1.43
3	A	403[C]	GOL	O2-C2	-5.10	1.28	1.43
3	A	405[B]	GOL	O1-C1	-2.97	1.29	1.42
3	A	402	GOL	O1-C1	2.32	1.52	1.42
3	A	405[B]	GOL	O2-C2	3.09	1.52	1.43

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	406	TRS	C3-C-C1	-4.27	101.53	110.78
5	A	407	1PE	OH3-C22-C12	-3.83	92.80	110.43
3	A	405[B]	GOL	O3-C3-C2	-2.97	95.78	110.18
3	A	405[B]	GOL	C3-C2-C1	-2.29	102.16	111.12
4	A	406	TRS	C2-C-C1	2.09	115.31	110.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 36 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	GOL	1	0
3	A	403[A]	GOL	8	0
3	A	403[B]	GOL	3	0
3	A	403[C]	GOL	3	0
3	A	404	GOL	1	0
3	A	405[A]	GOL	1	0
3	A	405[B]	GOL	1	0
4	A	406	TRS	14	0
5	A	407	1PE	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	312/334 (93%)	-0.16	8 (2%) 59 49	3, 6, 17, 39	2 (0%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	318	ILE	5.9
1	A	313	LEU	3.8
1	A	319	SER	3.5
1	A	293	PRO	3.3
1	A	9	SER	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	A	405[A]	6/6	0.88	0.32	46.56	4,9,19,26	6

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	405[B]	6/6	0.88	0.32	42.42	6,10,13,19	6
3	GOL	A	403[B]	6/6	0.92	0.23	22.07	3,5,5,6	6
5	1PE	A	407	7/16	0.89	0.36	14.57	20,22,35,37	0
3	GOL	A	403[C]	6/6	0.92	0.23	12.71	2,4,6,6	6
3	GOL	A	403[A]	6/6	0.92	0.23	11.06	2,2,3,4	6
3	GOL	A	404	6/6	0.95	0.19	9.63	7,18,33,38	0
4	TRS	A	406	8/8	0.87	0.20	5.33	7,16,24,29	8
3	GOL	A	402	6/6	0.95	0.13	2.81	11,13,17,18	0
2	CA	A	401	1/1	1.00	0.04	-2.74	2,2,2,2	0

## 6.5 Other polymers

There are no such residues in this entry.