



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:55 PM GMT

PDB ID : 4N76  
Title : Structure of Thermus thermophilus Argonaute bound to guide DNA and cleaved target DNA with Mn<sup>2+</sup>  
Authors : Sheng, G.; Zhao, H.; Wang, J.; Rao, Y.; Wang, Y.  
Deposited on : 2013-10-15  
Resolution : 2.89 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	685	<div> <div>7%</div> <div>69%</div> <div>22%</div> <div>6%</div> </div>
1	B	685	<div> <div>7%</div> <div>72%</div> <div>23%</div> <div>• •</div> </div>
2	C	21	<div> <div>29%</div> <div>19%</div> <div>52%</div> </div>
2	E	21	<div> <div>33%</div> <div>19%</div> <div>48%</div> </div>
3	D	10	<div> <div>80%</div> <div>20%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	10	 60% 40%

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10818 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Argonaute.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	641	Total	C	N	O	S	4	0	0
			4803	3069	895	834	5			
1	B	667	Total	C	N	O	S	0	0	0
			5115	3273	959	877	6			

- Molecule 2 is a DNA chain called 5'-D(P\*TP\*GP\*AP\*GP\*GP\*TP\*AP\*GP\*TP\*AP\*GP\*GP\*TP\*TP\*GP\*TP\*AP\*TP\*AP\*GP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	10	Total	C	N	O	P	0	0	0
			212	100	41	61	10			
2	E	11	Total	C	N	O	P	0	0	0
			234	110	46	67	11			


- Molecule 3 is a DNA chain called 5'-D(P\*TP\*AP\*CP\*TP\*AP\*CP\*CP\*TP\*CP\*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	10	Total	C	N	O	P	0	0	0
			201	96	33	62	10			
3	F	10	Total	C	N	O	P	0	0	0
			201	96	33	62	10			

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	3	Total	Mn	0	0
			3	3		
4	C	1	Total	Mn	0	0
			1	1		

- Molecule 5 is THYMIDINE-5'-MONOPHOSPHATE (three-letter code: DT) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>2</sub>O<sub>8</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			21	10	2	8	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	8	Total	O	0	0
			8	8		
6	B	14	Total	O	0	0
			14	14		
6	C	1	Total	O	0	0
			1	1		
6	E	3	Total	O	0	0
			3	3		
6	F	1	Total	O	0	0
			1	1		





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.92Å 117.42Å 160.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.37 – 2.89 49.37 – 2.89	Depositor EDS
% Data completeness (in resolution range)	95.0 (49.37-2.89) 94.7 (49.37-2.89)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.44 (at 2.91Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.219 , 0.269 0.222 , 0.271	Depositor DCC
$R_{free}$ test set	2294 reflections (5.07%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.6	Xtriage
Anisotropy	0.444	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 8.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	0 of 45264 reflections	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	10818	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.26	0/4910	0.46	1/6677 (0.0%)
1	B	0.27	0/5235	0.45	2/7115 (0.0%)
2	C	0.82	1/238 (0.4%)	1.21	1/365 (0.3%)
2	E	0.79	1/263 (0.4%)	1.18	1/404 (0.2%)
3	D	0.86	1/223 (0.4%)	1.18	0/339
3	F	0.92	1/223 (0.4%)	1.22	1/339 (0.3%)
All	All	0.35	4/11092 (0.0%)	0.57	6/15239 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	DT	OP3-P	-10.96	1.48	1.61
2	E	1	DT	OP3-P	-10.72	1.48	1.61
3	F	10	DT	OP3-P	-10.30	1.48	1.61
3	D	10	DT	OP3-P	-10.28	1.48	1.61

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	DT	OP1-P-OP2	-8.20	107.30	119.60
2	C	1	DT	OP1-P-OP2	-7.34	108.59	119.60
1	B	143	PRO	N-CA-CB	5.92	110.41	103.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	143	PRO	N-CA-CB	5.83	110.29	103.30
1	B	356	GLY	N-CA-C	5.35	126.47	113.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	437	VAL	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4803	0	4723	91	0
1	B	5115	0	5099	89	0
2	C	212	0	114	2	0
2	E	234	0	125	3	0
3	D	201	0	114	2	0
3	F	201	0	114	2	0
4	B	3	0	0	0	0
4	C	1	0	0	0	0
5	B	21	0	13	5	0
6	A	8	0	0	1	0
6	B	14	0	0	2	0
6	C	1	0	0	0	0
6	E	3	0	0	0	0
6	F	1	0	0	0	0
All	All	10818	0	10302	179	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 179 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:639:THR:HG22	1:A:640:ARG:HE	1.46	0.81

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:344:THR:HG21	1:A:460:LEU:HD11	1.62	0.80
1:B:639:THR:HG22	1:B:640:ARG:HE	1.46	0.80
1:B:608:ARG:HB2	1:B:611:ARG:HG3	1.67	0.76
1:B:99:ASP:N	1:B:105:GLU:OE1	2.18	0.74

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	633/685 (92%)	580 (92%)	47 (7%)	6 (1%)	21	57
1	B	655/685 (96%)	613 (94%)	36 (6%)	6 (1%)	21	57
All	All	1288/1370 (94%)	1193 (93%)	83 (6%)	12 (1%)	21	57

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	483	GLU
1	B	438	PRO
1	B	37	PRO
1	B	356	GLY
1	B	482	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	447/549 (81%)	409 (92%)	38 (8%)	13	37
1	B	492/549 (90%)	449 (91%)	43 (9%)	13	36
All	All	939/1098 (86%)	858 (91%)	81 (9%)	13	36

5 of 81 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	640	ARG
1	B	159	ASP
1	B	592	LEU
1	A	678	ASP
1	B	46	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	500	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	DT	B	704	-	17,22,22	1.24	3 (17%)	22,33,33	3.51	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	DT	B	704	-	-	0/6/22/22	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	704	DT	C6-C5	-2.18	1.34	1.40
5	B	704	DT	C6-N1	2.15	1.38	1.35
5	B	704	DT	C4-N3	2.84	1.38	1.33

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	704	DT	C5-C4-N3	-8.72	115.42	125.14
5	B	704	DT	OP2-P-OP3	2.98	118.74	107.38
5	B	704	DT	C4-N3-C2	13.24	126.69	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	704	DT	5	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	641/685 (93%)	0.54	47 (7%)	18 12	3, 25, 103, 143	1 (0%)
1	B	667/685 (97%)	0.51	46 (6%)	20 14	1, 27, 82, 113	0
2	C	10/21 (47%)	-0.15	0	100 100	22, 26, 65, 83	0
2	E	11/21 (52%)	-0.07	0	100 100	16, 24, 73, 100	0
3	D	10/10 (100%)	-0.11	0	100 100	27, 66, 90, 91	0
3	F	10/10 (100%)	-0.27	0	100 100	26, 56, 76, 78	0
All	All	1349/1432 (94%)	0.51	93 (6%)	20 14	1, 26, 92, 143	1 (0%)

The worst 5 of 93 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	233	LEU	10.4
1	A	232	ARG	8.6
1	B	19	LEU	6.3
1	A	220	GLY	6.2
1	A	258	THR	6.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	DT	B	704	21/21	0.95	0.28	-0.13	47,64,75,84	0
4	MN	B	701	1/1	0.96	0.14	-2.08	50,50,50,50	0
4	MN	B	702	1/1	0.79	0.08	-2.10	63,63,63,63	1
4	MN	B	703	1/1	0.79	0.09	-2.19	81,81,81,81	0
4	MN	C	101	1/1	0.98	0.15	-2.23	53,53,53,53	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.