



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 24, 2017 – 11:26 PM EST

PDB ID : 1N97
Title : Crystal Structure of CYP175A1 from *Thermus thermophilus* strain HB27
Authors : Yano, J.K.; Blasco, F.; Li, H.; Schmid, R.D.; Henne, A.; Poulos, T.L.
Deposited on : 2002-11-22
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028442
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028442


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	389	<div> <div>6%</div> <div> <div></div> <div>79%</div> <div>18%</div> <div>..</div> </div> </div>
1	B	389	<div> <div>4%</div> <div> <div></div> <div>79%</div> <div>17%</div> <div>..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SRT	A	601	X	-	-	X
2	SRT	B	602	X	-	-	X
4	EDO	A	715	-	-	-	X
4	EDO	A	717	-	-	-	X
4	EDO	B	704	-	-	-	X
4	EDO	B	707	-	-	-	X
4	EDO	B	708	-	-	-	X
4	EDO	B	709	-	-	-	X

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 6832 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYP175A1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	385	Total	C	N	O	S	0	0	0
			3052	1972	557	518	5			
1	B	385	Total	C	N	O	S	27	0	0
			3074	1985	560	524	5			

- Molecule 2 is S,R MESO-TARTARIC ACID (three-letter code: SRT) (formula: C₄H₆O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	4	6		
2	B	1	Total	C	O	0	0
			10	4	6		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	206	Total O 206 206	0	0
5	B	322	Total O 322 322	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CYP175A1



• Molecule 1: CYP175A1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.54Å 90.70Å 74.52Å 90.00° 99.68° 90.00°	Depositor
Resolution (Å)	19.99 – 1.80 19.99 – 1.80	Depositor EDS
% Data completeness (in resolution range)	92.0 (19.99-1.80) 92.0 (19.99-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.92 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.182 , 0.222 0.190 , 0.228	Depositor DCC
R_{free} test set	3639 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.246	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 59.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6832	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, EDO, SRT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.13	5/3136 (0.2%)	1.02	8/4263 (0.2%)
1	B	1.38	8/3158 (0.3%)	1.19	28/4288 (0.7%)
All	All	1.26	13/6294 (0.2%)	1.11	36/8551 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	10	TRP	C-N	-12.85	1.09	1.34
1	B	13	LEU	C-N	10.64	1.58	1.34
1	B	313	GLU	CD-OE2	-10.45	1.14	1.25
1	B	313	GLU	CD-OE1	10.29	1.36	1.25
1	B	106	TYR	CE2-CZ	-7.53	1.28	1.38

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	334	ARG	NE-CZ-NH2	-10.02	115.29	120.30
1	B	334	ARG	NE-CZ-NH1	9.14	124.87	120.30
1	B	187	ARG	NE-CZ-NH1	8.97	124.78	120.30
1	B	162	ASP	CB-CG-OD2	8.88	126.29	118.30
1	B	355	ARG	NE-CZ-NH1	8.65	124.62	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	10	TRP	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3052	0	3082	60	0
1	B	3074	0	3119	34	1
2	A	10	0	5	1	0
2	B	10	0	4	0	0
3	A	43	0	30	1	0
3	B	43	0	30	1	0
4	A	28	0	42	3	0
4	B	44	0	66	9	0
5	A	206	0	0	3	1
5	B	322	0	0	6	1
All	All	6832	0	6378	95	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 95 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:LEU:HD21	5:A:734:HOH:O	1.73	0.87
1:A:245:LYS:NZ	1:A:249:GLU:OE2	2.13	0.81
1:A:280:LEU:HD23	1:A:280:LEU:H	1.52	0.74
1:A:74:LEU:HB2	1:A:164:ILE:CD1	2.17	0.74
1:B:74:LEU:HD12	1:B:161:LEU:HG	1.70	0.73

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:748:HOH:O	5:B:1016:HOH:O[2_645]	1.70	0.50
1:B:252:GLU:OE1	5:A:867:HOH:O[2_656]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	383/389 (98%)	375 (98%)	7 (2%)	1 (0%)	46	29
1	B	383/389 (98%)	368 (96%)	11 (3%)	4 (1%)	19	5
All	All	766/778 (98%)	743 (97%)	18 (2%)	5 (1%)	26	11

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	14	LYS
1	A	208	ARG
1	B	6	LEU
1	B	12	TYR
1	B	282	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	305/319 (96%)	293 (96%)	12 (4%)	39	21
1	B	310/319 (97%)	299 (96%)	11 (4%)	43	25

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	615/638 (96%)	592 (96%)	23 (4%)	41	23

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	279	LEU
1	B	6	LEU
1	B	318	GLU
1	A	325	ARG
1	B	15	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	333	GLN
1	B	239	HIS
1	A	370	GLN
1	A	300	GLN
1	B	67	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SRT	A	601	-	3,9,9	0.73	0	6,12,12	2.10	2 (33%)
3	HEM	A	602	1	24,50,50	2.98	8 (33%)	16,82,82	2.04	7 (43%)
4	EDO	A	705	-	3,3,3	0.55	0	2,2,2	0.47	0
4	EDO	A	713	-	3,3,3	0.41	0	2,2,2	0.43	0
4	EDO	A	714	-	3,3,3	0.36	0	2,2,2	0.45	0
4	EDO	A	715	-	3,3,3	0.52	0	2,2,2	0.31	0
4	EDO	A	717	-	3,3,3	0.49	0	2,2,2	0.31	0
4	EDO	A	718	-	3,3,3	0.47	0	2,2,2	0.58	0
4	EDO	A	719	-	3,3,3	0.39	0	2,2,2	0.77	0
2	SRT	B	602	-	3,9,9	0.29	0	6,12,12	3.03	4 (66%)
3	HEM	B	603	1	24,50,50	2.33	9 (37%)	16,82,82	2.18	7 (43%)
4	EDO	B	701	-	3,3,3	0.77	0	2,2,2	0.20	0
4	EDO	B	702	-	3,3,3	0.84	0	2,2,2	0.34	0
4	EDO	B	703	-	3,3,3	0.30	0	2,2,2	0.59	0
4	EDO	B	704	-	3,3,3	0.64	0	2,2,2	0.05	0
4	EDO	B	706	-	3,3,3	0.43	0	2,2,2	0.34	0
4	EDO	B	707	-	3,3,3	0.73	0	2,2,2	0.47	0
4	EDO	B	708	-	3,3,3	0.55	0	2,2,2	0.44	0
4	EDO	B	709	-	3,3,3	0.50	0	2,2,2	0.34	0
4	EDO	B	710	-	3,3,3	0.44	0	2,2,2	0.28	0
4	EDO	B	711	-	3,3,3	0.60	0	2,2,2	0.94	0
4	EDO	B	712	-	3,3,3	0.60	0	2,2,2	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SRT	A	601	-	2/2/4/4	0/4/12/12	0/0/0/0
3	HEM	A	602	1	-	0/6/54/54	0/0/8/8
4	EDO	A	705	-	-	0/1/1/1	0/0/0/0
4	EDO	A	713	-	-	0/1/1/1	0/0/0/0
4	EDO	A	714	-	-	0/1/1/1	0/0/0/0
4	EDO	A	715	-	-	0/1/1/1	0/0/0/0
4	EDO	A	717	-	-	0/1/1/1	0/0/0/0
4	EDO	A	718	-	-	0/1/1/1	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	719	-	-	0/1/1/1	0/0/0/0
2	SRT	B	602	-	2/2/4/4	0/4/12/12	0/0/0/0
3	HEM	B	603	1	-	0/6/54/54	0/0/8/8
4	EDO	B	701	-	-	0/1/1/1	0/0/0/0
4	EDO	B	702	-	-	0/1/1/1	0/0/0/0
4	EDO	B	703	-	-	0/1/1/1	0/0/0/0
4	EDO	B	704	-	-	0/1/1/1	0/0/0/0
4	EDO	B	706	-	-	0/1/1/1	0/0/0/0
4	EDO	B	707	-	-	0/1/1/1	0/0/0/0
4	EDO	B	708	-	-	0/1/1/1	0/0/0/0
4	EDO	B	709	-	-	0/1/1/1	0/0/0/0
4	EDO	B	710	-	-	0/1/1/1	0/0/0/0
4	EDO	B	711	-	-	0/1/1/1	0/0/0/0
4	EDO	B	712	-	-	0/1/1/1	0/0/0/0

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	HEM	C3C-C2C	-6.97	1.31	1.40
3	B	603	HEM	C3C-C2C	-5.64	1.33	1.40
3	A	602	HEM	C3B-C2B	-5.27	1.33	1.40
3	B	603	HEM	C3B-C2B	-4.84	1.34	1.40
3	B	603	HEM	C4D-ND	-2.55	1.33	1.36

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	602	SRT	O2-C2-C3	-6.14	91.14	108.67
3	B	603	HEM	CBD-CAD-C3D	-4.00	105.46	112.47
3	A	602	HEM	CAD-CBD-CGD	-3.58	105.81	112.78
3	A	602	HEM	CBD-CAD-C3D	-3.52	106.29	112.47
2	A	601	SRT	O2-C2-C3	-3.31	99.20	108.67

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	602	SRT	C2
2	B	602	SRT	C3
2	A	601	SRT	C2
2	A	601	SRT	C3

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	SRT	1	0
3	A	602	HEM	1	0
4	A	714	EDO	1	0
4	A	715	EDO	1	0
4	A	717	EDO	1	0
3	B	603	HEM	1	0
4	B	703	EDO	1	0
4	B	707	EDO	3	0
4	B	708	EDO	1	0
4	B	709	EDO	1	0
4	B	710	EDO	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	385/389 (98%)	0.30	24 (6%)	24 19	20, 39, 60, 73	25 (6%)
1	B	382/389 (98%)	0.03	17 (4%)	37 31	15, 27, 55, 82	54 (14%)
All	All	767/778 (98%)	0.16	41 (5%)	30 25	15, 33, 58, 82	79 (10%)

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	6	LEU	8.8
1	A	40	ARG	7.8
1	A	6	LEU	4.4
1	B	8	GLU	4.3
1	A	41	PHE	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	EDO	B	709	4/4	0.72	0.43	30.89	55,64,69,71	0
4	EDO	B	707	4/4	0.92	0.33	14.22	40,44,50,50	0
4	EDO	B	708	4/4	0.62	0.46	10.14	60,63,68,68	0
4	EDO	A	717	4/4	0.76	0.36	8.48	62,68,68,69	0
4	EDO	B	704	4/4	0.68	0.22	7.61	58,64,67,70	0
2	SRT	B	602	10/10	0.93	0.17	6.08	32,36,41,41	0
2	SRT	A	601	10/10	0.85	0.22	5.18	41,49,55,56	0
4	EDO	A	715	4/4	0.72	0.20	3.95	67,70,70,72	0
4	EDO	B	711	4/4	0.92	0.14	1.68	33,40,43,45	0
4	EDO	B	706	4/4	0.89	0.20	1.65	39,48,49,55	0
4	EDO	B	702	4/4	0.92	0.12	1.31	33,38,39,42	0
4	EDO	B	703	4/4	0.91	0.13	1.25	45,47,50,53	0
4	EDO	B	710	4/4	0.96	0.12	0.54	38,45,46,53	0
4	EDO	A	718	4/4	0.95	0.11	0.53	45,48,50,54	0
4	EDO	A	705	4/4	0.86	0.16	0.40	50,51,54,59	0
4	EDO	A	713	4/4	0.79	0.16	0.12	69,69,69,69	0
4	EDO	B	701	4/4	0.94	0.09	-0.34	33,37,39,41	0
3	HEM	A	602	43/43	0.98	0.06	-1.08	18,24,32,37	0
3	HEM	B	603	43/43	0.99	0.06	-1.45	11,15,24,30	0
4	EDO	A	719	4/4	0.79	0.31	-	58,66,66,71	0
4	EDO	A	714	4/4	0.70	0.25	-	69,70,70,71	0
4	EDO	B	712	4/4	0.86	0.27	-	60,62,62,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.