



Full wwPDB NMR Structure Validation Report ⓘ

Jan 10, 2017 – 04:55 PM EST

PDB ID : 2NAZ
Title : The solution NMR structure of the C-terminal effector domain of BfmR from *Acinetobacter baumannii*
Authors : Olson, A.L.; Thompson, R.J.; Cavanagh, J.; Feldmann, E.A.; Bobay, B.G.
Deposited on : 2016-01-15

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20028442
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 85%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	118	

2 Ensemble composition and analysis ⓘ

This entry contains 10 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:136-A:188, A:204-A:213, A:218-A:221, A:228-A:232 (72)	0.10	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 3, 4, 5, 7, 10
2	2, 9
Single-model clusters	6; 8

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1752 atoms, of which 873 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Transcriptional regulatory protein RstA.

Mol	Chain	Residues	Atoms					Trace
1	A	109	Total	C	H	N	O	0
			1752	547	873	158	174	

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	129	MET	-	INITIATING METHIONINE	UNP D0C9D6
A	239	LEU	-	EXPRESSION TAG	UNP D0C9D6
A	240	GLU	-	EXPRESSION TAG	UNP D0C9D6
A	241	HIS	-	EXPRESSION TAG	UNP D0C9D6
A	242	HIS	-	EXPRESSION TAG	UNP D0C9D6
A	243	HIS	-	EXPRESSION TAG	UNP D0C9D6
A	244	HIS	-	EXPRESSION TAG	UNP D0C9D6
A	245	HIS	-	EXPRESSION TAG	UNP D0C9D6
A	246	HIS	-	EXPRESSION TAG	UNP D0C9D6

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Transcriptional regulatory protein RstA




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: Transcriptional regulatory protein RstA



4.2.2 Score per residue for model 2


- Molecule 1: Transcriptional regulatory protein RstA






4.2.3 Score per residue for model 3

- Molecule 1: Transcriptional regulatory protein RstA




4.2.4 Score per residue for model 4

- Molecule 1: Transcriptional regulatory protein RstA




4.2.5 Score per residue for model 5 (medoid)

- Molecule 1: Transcriptional regulatory protein RstA




4.2.6 Score per residue for model 6

- Molecule 1: Transcriptional regulatory protein RstA




4.2.7 Score per residue for model 7

- Molecule 1: Transcriptional regulatory protein RstA




4.2.8 Score per residue for model 8

- Molecule 1: Transcriptional regulatory protein RstA




4.2.9 Score per residue for model 9

- Molecule 1: Transcriptional regulatory protein RstA



4.2.10 Score per residue for model 10

- Molecule 1: Transcriptional regulatory protein RstA



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
SPARKY	structure solution	
SPARKY	structure solution	
SPARKY	structure solution	
SPARKY	structure solution	
AMBER	refinement	12.0

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2naz_cs.cif
Number of chemical shift lists	1
Total number of shifts	1202
Number of shifts mapped to atoms	1202
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	85%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.77±0.01	0±0/592 (0.0±0.0%)	1.37±0.02	6±1/799 (0.8±0.1%)
All	All	0.77	0/5920 (0.0%)	1.37	60/7990 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	3.0±0.8
All	All	0	30

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	149	ARG	NE-CZ-NH1	7.79	124.19	120.30	7	2
1	A	181	ARG	NE-CZ-NH1	7.14	123.87	120.30	1	6
1	A	165	TYR	CB-CG-CD2	-6.81	116.91	121.00	1	10
1	A	177	ARG	NE-CZ-NH1	6.58	123.59	120.30	5	7
1	A	220	ARG	NE-CZ-NH1	6.55	123.58	120.30	5	10
1	A	181	ARG	NE-CZ-NH2	-6.10	117.25	120.30	1	3
1	A	229	TYR	CB-CG-CD2	-6.09	117.35	121.00	2	3
1	A	179	LEU	CB-CG-CD2	5.99	121.19	111.00	6	7
1	A	208	ARG	NE-CZ-NH1	5.79	123.19	120.30	7	2
1	A	206	ARG	NE-CZ-NH1	5.64	123.12	120.30	2	3
1	A	136	ARG	NE-CZ-NH1	5.43	123.02	120.30	8	4
1	A	187	ARG	NE-CZ-NH1	5.42	123.01	120.30	10	1
1	A	207	ILE	CA-CB-CG1	5.28	121.04	111.00	6	2

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	229	TYR	Sidechain	10
1	A	156	GLU	Peptide	9
1	A	221	ILE	Peptide	7
1	A	181	ARG	Sidechain	1
1	A	136	ARG	Sidechain	1
1	A	177	ARG	Sidechain	1
1	A	165	TYR	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	582	590	590	13±2
All	All	5820	5900	5900	134

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:184:ILE:H	1:A:184:ILE:HD12	0.59	1.57	7	6
1:A:184:ILE:HD12	1:A:184:ILE:H	0.58	1.56	8	4
1:A:139:PHE:CZ	1:A:172:ALA:HB1	0.57	2.35	10	10
1:A:174:ASN:CB	1:A:179:LEU:HD11	0.56	2.31	6	6
1:A:160:PHE:CE1	1:A:210:LYS:HE3	0.55	2.36	6	8
1:A:139:PHE:CE2	1:A:172:ALA:HB1	0.54	2.37	10	10
1:A:171:LEU:HB3	1:A:231:PHE:HB2	0.54	1.79	10	4
1:A:160:PHE:CE2	1:A:210:LYS:HE3	0.52	2.39	8	1
1:A:144:ILE:HD11	1:A:168:LEU:HD13	0.52	1.80	8	10
1:A:153:LEU:HD13	1:A:211:ILE:HA	0.52	1.81	8	10
1:A:179:LEU:CB	1:A:184:ILE:HD11	0.49	2.37	8	9
1:A:144:ILE:HD11	1:A:168:LEU:CD1	0.49	2.38	8	9
1:A:144:ILE:CD1	1:A:168:LEU:HD22	0.49	2.38	1	9
1:A:181:ARG:HA	1:A:184:ILE:HD13	0.49	1.85	6	1
1:A:144:ILE:HD11	1:A:168:LEU:HD22	0.47	1.85	1	7

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:171:LEU:HD21	1:A:229:TYR:HB2	0.45	1.88	6	2
1:A:160:PHE:CE1	1:A:210:LYS:HE2	0.45	2.47	10	1
1:A:139:PHE:CD1	1:A:142:LEU:HD23	0.45	2.47	4	10
1:A:137:ILE:HG22	1:A:139:PHE:CE2	0.45	2.47	6	9
1:A:160:PHE:CE1	1:A:210:LYS:CE	0.43	3.01	10	1
1:A:174:ASN:HB3	1:A:179:LEU:HD11	0.43	1.90	8	1
1:A:170:LEU:HD11	1:A:187:ARG:HG3	0.42	1.90	9	2
1:A:179:LEU:HB2	1:A:184:ILE:HD11	0.42	1.90	4	1
1:A:146:ASN:HB2	1:A:169:TRP:CZ3	0.41	2.50	6	1
1:A:204:ILE:HG13	1:A:229:TYR:CZ	0.41	2.51	5	1
1:A:179:LEU:C	1:A:184:ILE:HD11	0.40	2.36	1	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	72/118 (61%)	62±1 (85±1%)	9±1 (12±1%)	2±1 (3±1%)	10	44
All	All	720/1180 (61%)	615 (85%)	85 (12%)	20 (3%)	10	44

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	181	ARG	9
1	A	180	SER	6
1	A	159	ASP	4
1	A	208	ARG	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	63/105 (60%)	53±1 (84±2%)	10±1 (16±2%)	6	43
All	All	630/1050 (60%)	527 (84%)	103 (16%)	6	43

All 16 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	185	PHE	10
1	A	168	LEU	10
1	A	161	THR	10
1	A	139	PHE	10
1	A	179	LEU	10
1	A	219	LYS	9
1	A	167	LEU	8
1	A	188	LEU	8
1	A	183	ASP	7
1	A	152	THR	5
1	A	170	LEU	5
1	A	187	ARG	4
1	A	145	ASP	3
1	A	220	ARG	2
1	A	136	ARG	1
1	A	141	ASP	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 85% for the well-defined parts and 76% for the entire structure.

7.1 Chemical shift list 1

File name: 2naz_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1202
Number of shifts mapped to atoms	1202
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	96	-0.45 ± 0.09	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	90	-0.07 ± 0.16	None needed (< 0.5 ppm)
$^{13}\text{C}'$	95	-0.13 ± 0.13	None needed (< 0.5 ppm)
^{15}N	96	-1.41 ± 0.15	Should be applied

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 85%, i.e. 785 atoms were assigned a chemical shift out of a possible 928. 14 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	350/356 (98%)	141/142 (99%)	139/144 (97%)	70/70 (100%)
Sidechain	390/508 (77%)	237/295 (80%)	153/184 (83%)	0/29 (0%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	45/64 (70%)	30/34 (88%)	15/29 (52%)	0/1 (0%)
Overall	785/928 (85%)	408/471 (87%)	307/357 (86%)	70/100 (70%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 76%, i.e. 1062 atoms were assigned a chemical shift out of a possible 1392. 17 out of 18 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	481/539 (89%)	194/215 (90%)	191/218 (88%)	96/106 (91%)
Sidechain	536/781 (69%)	328/454 (72%)	208/279 (75%)	0/48 (0%)
Aromatic	45/72 (62%)	30/38 (79%)	15/33 (45%)	0/1 (0%)
Overall	1062/1392 (76%)	552/707 (78%)	414/530 (78%)	96/155 (62%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	146	ASN	HB2	0.71	4.36 – 1.26	-6.8
1	A	173	SER	HB2	2.12	5.18 – 2.58	-6.8

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

