



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:06 PM GMT

PDB ID : 1NJY
Title : THYMINE-THYMINE MISMATCH AT THE POLYMERASE ACTIVE SITE
Authors : Johnson, S.J.; Beese, L.S.
Deposited on : 2003-01-02
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	11	
2	C	16	
3	A	580	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SUC	A	901	-	-	-	X
4	SUC	A	902	-	-	-	X
7	TTP	A	20	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5593 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA PRIMER STRAND.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	11	Total	C	N	O	P	0	0	0
			222	107	40	65	10			


- Molecule 2 is a DNA chain called DNA TEMPLATE STRAND.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	13	Total	C	N	O	P	0	0	0
			265	127	50	76	12			

- Molecule 3 is a protein called DNA POLYMERASE I.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	580	Total	C	N	O	S	0	0	0
			4650	2956	807	870	17			

- Molecule 4 is SUGAR (SUCROSE) (three-letter code: SUC) (formula: C₁₂H₂₂O₁₁).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			23	12	11		
4	A	1	Total	C	O	0	0
			23	12	11		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		

- Molecule 7 is THYMIDINE-5'-TRIPHOSPHATE (three-letter code: TTP) (formula: C₁₀H₁₇N₂O₁₄P₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	P	0	0
			13	10	3		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	351	Total O 351 351	0	0
8	B	11	Total O 11 11	0	0
8	C	24	Total O 24 24	0	0

- Molecule 1: DNA PRIMER STRAND



DG
DA
DC
G3

G13
C14
A15

Category	Item	Value	Color
Category 1	K706	6332	Red
	A707	9533	Green
	V708	1534	Green
	H709	4535	Red
	F710	7536	Red
	G711	2537	Green
	L712	8538	Green
	I716	3547	Red
	K718	9548	Green
	V719	7549	Red
Category 2	L725	7550	Red
	M726	8551	Red
	N727	7552	Red
	S728	4553	Green
	R729	7554	Green
	F730	3558	Red
	E731	2562	Green
	E734	4565	Green
	R738	7566	Green
	Q755	7571	Green
Category 3	L767	1574	Green
	R779	9584	Green
	F786	7587	Red
	R789	7588	Green
	M792	6610	Green
	L796	7625	Green
	D802	7626	Green
	A813	7627	Green
	E817	6633	Green
	A822	7636	Green
Category 4	A836	7662	Green
	V848	8677	Green
	P849	7678	Red
	E850	7679	Green
	Q854	7689	Green
	A876	7690	Green
	K876	7691	Red
		7692	Green
		8696	Green
		7704	Green
Category 5	A297	1297	Red
	K298	2298	Green
	L303	4303	Red
	A304	7304	Red
	D305	8305	Red
	R306	2306	Green
	M311	3311	Green
	G334	4334	Green
	P348	7348	Green
	D408	2408	Green
Category 6	Q418	3418	Green
	Y429	4429	Green
	G430	7430	Green
	K431	8431	Red
	G432	4432	Green
	A433	7433	Red
	K434	8434	Red
	R435	2435	Green
	A436	3436	Green
	L457	4457	Green
Category 7	F458	7458	Green
	R459	2459	Green
	P460	3460	Green
	S483	4483	Green
	I484	7484	Green
	L485	8485	Green
	A486	2486	Green
	E487	3487	Green
	M488	4488	Green
	E489	7489	Green
Category 8	E501	2501	Green
	K505	3505	Green
	E509	4509	Red
	V514	7514	Green
	E515	8515	Green
	Q516	2516	Green
	R517	3517	Green
	I518	4518	Green
	Y519	7519	Red
	E520	8520	Green
Category 9	E525	2525	Green
	I528	3528	Green
	N529	4529	Green
	S530	7530	Green
	E531	8531	Green

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.98Å 93.72Å 106.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.33 – 2.00 45.33 – 2.00	Depositor EDS
% Data completeness (in resolution range)	90.7 (45.33-2.00) 95.7 (45.33-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.17 (at 2.00Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.209 , 0.243 0.209 , 0.243	Depositor DCC
R_{free} test set	2713 reflections (5.05%)	DCC
Wilson B-factor (Å ²)	24.7	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 50.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 56729 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5593	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, TTP, SUC, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.31	0/248	0.76	0/381
2	C	0.33	0/297	0.66	0/457
3	A	0.33	0/4734	0.57	0/6398
All	All	0.33	0/5279	0.58	0/7236

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	222	0	126	9	0
2	C	265	0	148	3	0
3	A	4650	0	4698	54	0
4	A	46	0	44	1	0
5	A	10	0	0	0	0
6	A	1	0	0	0	0
7	A	13	0	0	1	0
8	A	351	0	0	2	0
8	B	11	0	0	0	0
8	C	24	0	0	0	0
All	All	5593	0	5016	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:789:ARG:HA	3:A:792:MET:HE3	1.44	0.99
3:A:534:LEU:HD11	3:A:574:ILE:HD13	1.70	0.72
3:A:306:ARG:HH11	3:A:306:ARG:HB3	1.54	0.72
3:A:306:ARG:HB3	3:A:306:ARG:NH1	2.05	0.72
3:A:691:GLN:HG3	3:A:738:ARG:HH22	1.56	0.71
1:B:25:DA:H2''	1:B:26:DC:H5'	1.73	0.71
1:B:19:DG:H2''	1:B:20:DC:H5'	1.73	0.69
3:A:728:SER:HB3	3:A:731:GLU:HG3	1.76	0.68
2:C:13:DG:H2''	2:C:14:DC:H5'	1.76	0.67
2:C:3:DG:H2'	2:C:3:DG:N3	2.09	0.65
3:A:536:VAL:O	3:A:540:GLU:HB2	1.97	0.64
1:B:19:DG:H2''	1:B:20:DC:C5'	2.29	0.61
3:A:584:GLN:O	3:A:589:GLU:HG3	2.01	0.61
3:A:459:ARG:HB3	3:A:460:PRO:HD3	1.84	0.59
3:A:408:ASP:HB2	4:A:901:SUC:O1'	2.05	0.57
3:A:516:GLN:O	3:A:520:GLU:HG3	2.04	0.57
1:B:22:DA:H2''	1:B:23:DT:H5'	1.87	0.57
3:A:558:ALA:O	3:A:562:GLU:HG2	2.05	0.57
3:A:501:GLU:O	3:A:505:LYS:HG2	2.05	0.56
3:A:728:SER:HB3	3:A:731:GLU:CG	2.36	0.56
3:A:706:LYS:HE3	7:A:20:TTP:O1B	2.06	0.55
3:A:689:ILE:O	3:A:738:ARG:NH1	2.40	0.55
2:C:14:DC:H2''	2:C:15:DA:O5'	2.06	0.55
1:B:25:DA:H2''	1:B:26:DC:C5'	2.37	0.55
3:A:813:ALA:O	3:A:817:GLU:HG3	2.07	0.54
1:B:22:DA:H2''	1:B:23:DT:C5'	2.37	0.53
3:A:610:LEU:C	3:A:610:LEU:HD23	2.31	0.51
3:A:306:ARG:HH11	3:A:306:ARG:CB	2.22	0.51
3:A:434:LYS:N	3:A:434:LYS:HD3	2.27	0.50
3:A:519:TYR:CD1	3:A:525:GLU:HA	2.48	0.49
1:B:22:DA:H1'	1:B:23:DT:H5''	1.92	0.49
3:A:587:TYR:CE2	3:A:627:PRO:HD3	2.48	0.48
3:A:483:SER:O	3:A:487:GLU:HG3	2.14	0.48
3:A:691:GLN:CG	3:A:738:ARG:HH22	2.26	0.47
3:A:725:LEU:O	3:A:727:ILE:HG23	2.14	0.47
3:A:518:ILE:HD12	3:A:528:ILE:HD13	1.97	0.46
3:A:692:VAL:HB	3:A:696:GLU:HB2	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:565:ALA:HA	3:A:571:VAL:HB	1.98	0.44
3:A:677:ARG:HB2	3:A:679:LEU:HG	1.99	0.44
3:A:633:GLY:O	3:A:636:ILE:HG12	2.18	0.44
3:A:435:ARG:O	3:A:436:ALA:HB2	2.18	0.44
3:A:565:ALA:HB3	3:A:566:PRO:HD3	2.00	0.43
3:A:530:SER:HB3	3:A:533:GLN:HB2	2.00	0.43
3:A:551:LYS:HG3	3:A:551:LYS:O	2.17	0.43
3:A:662:LEU:HD13	3:A:796:ILE:HD11	2.01	0.43
1:B:28:DT:O4'	3:A:625:ASN:HA	2.18	0.43
3:A:734:GLU:O	3:A:738:ARG:HG2	2.17	0.43
3:A:708:VAL:O	3:A:712:ILE:HG12	2.19	0.43
3:A:514:VAL:O	3:A:518:ILE:HG13	2.19	0.42
3:A:457:LEU:O	3:A:460:PRO:HD2	2.20	0.42
3:A:755:GLN:HG2	8:A:2619:HOH:O	2.18	0.42
3:A:515:GLU:HG2	3:A:519:TYR:CE2	2.55	0.42
3:A:457:LEU:C	3:A:460:PRO:HD2	2.40	0.42
3:A:767:LEU:HG	3:A:802:ASP:HB3	2.01	0.42
3:A:485:LEU:O	3:A:489:GLU:HG3	2.20	0.41
3:A:850:GLU:OE2	3:A:854:GLN:HG3	2.20	0.41
3:A:418:GLN:HA	8:A:2575:HOH:O	2.20	0.41
1:B:20:DC:H2''	1:B:21:DG:C8	2.56	0.41
3:A:822:ALA:CB	3:A:836:ALA:HB2	2.50	0.41
3:A:848:VAL:HB	3:A:849:PRO:HD3	2.03	0.41
3:A:789:ARG:CA	3:A:792:MET:HE3	2.33	0.40
3:A:704:GLN:O	3:A:708:VAL:HG23	2.20	0.40
3:A:304:ALA:CB	3:A:311:MET:HE1	2.51	0.40
3:A:429:TYR:O	3:A:435:ARG:HA	2.21	0.40
3:A:334:GLY:HA2	3:A:348:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	A	578/580 (100%)	555 (96%)	20 (4%)	3 (0%)	34 26

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	298	LYS
3	A	551	LYS
3	A	628	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	A	495/496 (100%)	492 (99%)	3 (1%)	90 93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	434	LYS
3	A	554	TYR
3	A	779	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
7	TTP	A	20	-	8,12,30	1.27	0	15,20,47	1.18	2 (13%)
4	SUC	A	901	-	24,24,24	1.56	3 (12%)	36,36,36	0.97	2 (5%)
4	SUC	A	902	-	24,24,24	1.62	3 (12%)	36,36,36	0.95	2 (5%)
5	SO4	A	911	-	4,4,4	0.22	0	6,6,6	0.07	0
5	SO4	A	912	-	4,4,4	0.19	0	6,6,6	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	TTP	A	20	-	-	0/12/12/34	0/0/0/2
4	SUC	A	901	-	-	0/12/51/51	0/2/2/2
4	SUC	A	902	-	-	0/12/51/51	0/2/2/2
5	SO4	A	911	-	-	0/0/0/0	0/0/0/0
5	SO4	A	912	-	-	0/0/0/0	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	902	SUC	C4-C5	2.16	1.57	1.53
4	A	901	SUC	C4-C5	2.32	1.58	1.53
4	A	902	SUC	O5-C1	2.54	1.48	1.41
4	A	901	SUC	O5-C1	2.60	1.48	1.41
4	A	901	SUC	C3-C2	5.63	1.67	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	902	SUC	C3-C2	5.77	1.67	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	901	SUC	C1-C2-C3	-2.21	105.61	109.97
4	A	902	SUC	C1-C2-C3	-2.06	105.92	109.97
4	A	901	SUC	C1-O5-C5	2.00	117.64	113.75
7	A	20	TTP	O2A-PA-O1A	2.09	117.32	110.58
4	A	902	SUC	C1-O5-C5	2.23	118.07	113.75
7	A	20	TTP	O2G-PG-O1G	2.69	119.25	110.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	20	TTP	1	0
4	A	901	SUC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	11/11 (100%)	0.74	2 (18%) 2 2	37, 41, 71, 73	0
2	C	13/16 (81%)	1.20	4 (30%) 1 1	30, 42, 82, 86	0
3	A	580/580 (100%)	0.29	27 (4%) 35 37	14, 26, 48, 59	0
All	All	604/607 (99%)	0.32	33 (5%) 29 30	14, 27, 49, 86	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	552	THR	6.2
3	A	433	ALA	6.1
3	A	549	LYS	6.0
3	A	550	THR	6.0
3	A	551	LYS	5.9
2	C	15	DA	5.9
3	A	297	ALA	5.1
2	C	13	DG	4.7
2	C	14	DC	4.2
3	A	553	GLY	4.1
3	A	719	TYR	3.8
3	A	520	GLU	3.8
3	A	303	LEU	3.6
3	A	434	LYS	3.5
3	A	305	ASP	3.4
1	B	19	DG	3.4
3	A	710	PHE	3.1
3	A	431	LYS	2.9
3	A	509	GLU	2.9
3	A	716	ILE	2.8
3	A	532	LYS	2.7
3	A	729	ARG	2.4
3	A	306	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	20	DC	2.3
3	A	691	GLN	2.3
3	A	730	LYS	2.3
3	A	786	PHE	2.2
3	A	519	TYR	2.2
3	A	525	GLU	2.2
3	A	304	ALA	2.2
3	A	298	LYS	2.1
3	A	547	LEU	2.1
2	C	3	DG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	SUC	A	902	23/23	0.70	0.35	6.67	58,59,60,60	0
4	SUC	A	901	23/23	0.77	0.22	3.08	47,48,49,49	0
7	TTP	A	20	13/29	0.72	0.27	2.38	78,78,79,79	0
5	SO4	A	912	5/5	0.93	0.13	-0.52	57,57,57,57	0
5	SO4	A	911	5/5	0.97	0.15	-	65,65,65,65	0
6	MG	A	920	1/1	0.97	0.07	-	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.