



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:01 PM GMT

PDB ID : 4NJX
Title : Human histidine triad nucleotide-binding protein 2 (hHINT2) in P41212 space group at 2.83 Å
Authors : Dolot, R.M.; Włodarczyk, A.; Bujacz, G.D.; Nawrot, B.
Deposited on : 2013-11-11
Resolution : 2.83 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3170 (2.88-2.80)
Clashscore	102246	3658 (2.88-2.80)
Ramachandran outliers	100387	3591 (2.88-2.80)
Sidechain outliers	100360	3594 (2.88-2.80)
RSRZ outliers	91569	3184 (2.88-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	172	<div> <div>2%</div> <div>57%</div> <div>10%</div> <div>33%</div> </div>
1	B	172	<div> <div>%</div> <div>56%</div> <div>8%</div> <div>34%</div> </div>
1	C	172	<div> <div>%</div> <div>46%</div> <div>9%</div> <div>41%</div> </div>
1	D	172	<div> <div>48%</div> <div>9%</div> <div>42%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3428 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histidine triad nucleotide-binding protein 2, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	116	Total	C	N	O	S	0	1	0
			907	582	161	163	1			
1	B	113	Total	C	N	O	S	0	1	0
			889	571	157	160	1			
1	C	101	Total	C	N	O	S	0	0	0
			790	507	140	142	1			
1	D	100	Total	C	N	O	S	0	0	0
			784	504	139	140	1			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	MET	-	INITIATING METHIONINE	UNP Q9BX68
A	-7	SER	-	EXPRESSION TAG	UNP Q9BX68
A	-6	HIS	-	EXPRESSION TAG	UNP Q9BX68
A	-5	HIS	-	EXPRESSION TAG	UNP Q9BX68
A	-4	HIS	-	EXPRESSION TAG	UNP Q9BX68
A	-3	HIS	-	EXPRESSION TAG	UNP Q9BX68
A	-2	HIS	-	EXPRESSION TAG	UNP Q9BX68
A	-1	HIS	-	EXPRESSION TAG	UNP Q9BX68
A	0	SER	-	EXPRESSION TAG	UNP Q9BX68
B	-8	MET	-	INITIATING METHIONINE	UNP Q9BX68
B	-7	SER	-	EXPRESSION TAG	UNP Q9BX68
B	-6	HIS	-	EXPRESSION TAG	UNP Q9BX68
B	-5	HIS	-	EXPRESSION TAG	UNP Q9BX68
B	-4	HIS	-	EXPRESSION TAG	UNP Q9BX68
B	-3	HIS	-	EXPRESSION TAG	UNP Q9BX68
B	-2	HIS	-	EXPRESSION TAG	UNP Q9BX68
B	-1	HIS	-	EXPRESSION TAG	UNP Q9BX68
B	0	SER	-	EXPRESSION TAG	UNP Q9BX68
C	-8	MET	-	INITIATING METHIONINE	UNP Q9BX68
C	-7	SER	-	EXPRESSION TAG	UNP Q9BX68
C	-6	HIS	-	EXPRESSION TAG	UNP Q9BX68

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-5	HIS	-	EXPRESSION TAG	UNP Q9BX68
C	-4	HIS	-	EXPRESSION TAG	UNP Q9BX68
C	-3	HIS	-	EXPRESSION TAG	UNP Q9BX68
C	-2	HIS	-	EXPRESSION TAG	UNP Q9BX68
C	-1	HIS	-	EXPRESSION TAG	UNP Q9BX68
C	0	SER	-	EXPRESSION TAG	UNP Q9BX68
D	-8	MET	-	INITIATING METHIONINE	UNP Q9BX68
D	-7	SER	-	EXPRESSION TAG	UNP Q9BX68
D	-6	HIS	-	EXPRESSION TAG	UNP Q9BX68
D	-5	HIS	-	EXPRESSION TAG	UNP Q9BX68
D	-4	HIS	-	EXPRESSION TAG	UNP Q9BX68
D	-3	HIS	-	EXPRESSION TAG	UNP Q9BX68
D	-2	HIS	-	EXPRESSION TAG	UNP Q9BX68
D	-1	HIS	-	EXPRESSION TAG	UNP Q9BX68
D	0	SER	-	EXPRESSION TAG	UNP Q9BX68


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	19	Total O 19 19	0	0
2	B	15	Total O 15 15	0	0
2	C	11	Total O 11 11	0	0
2	D	13	Total O 13 13	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histidine triad nucleotide-binding protein 2, mitochondrial




- Molecule 1: Histidine triad nucleotide-binding protein 2, mitochondrial



- Molecule 1: Histidine triad nucleotide-binding protein 2, mitochondrial



- Molecule 1: Histidine triad nucleotide-binding protein 2, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	76.38 Å 76.38 Å 133.25 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.96 – 2.83 41.96 – 2.83	Depositor EDS
% Data completeness (in resolution range)	99.7 (41.96-2.83) 99.8 (41.96-2.83)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.89 (at 2.86 Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.178 , 0.257 0.180 , 0.257	Depositor DCC
R_{free} test set	496 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	48.9	Xtriage
Anisotropy	0.343	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 48.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	2 of 9888 reflections (0.020%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3428	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.30 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.3387e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/929	0.84	1/1261 (0.1%)
1	B	0.66	0/910	0.90	4/1236 (0.3%)
1	C	0.66	0/809	0.90	2/1099 (0.2%)
1	D	0.61	0/803	0.88	3/1091 (0.3%)
All	All	0.64	0/3451	0.88	10/4687 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	154	GLY	N-CA-C	7.62	132.14	113.10
1	A	132	ARG	CG-CD-NE	6.82	126.12	111.80
1	D	132	ARG	CG-CD-NE	6.72	125.90	111.80
1	C	132	ARG	CG-CD-NE	6.60	125.66	111.80
1	D	67	ASP	CB-CG-OD2	6.41	124.07	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	162	PRO	Peptide
1	C	65	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	907	0	921	16	0
1	B	889	0	904	13	1
1	C	790	0	799	19	1
1	D	784	0	794	16	0
2	A	19	0	0	1	1
2	B	15	0	0	0	1
2	C	11	0	0	1	0
2	D	13	0	0	0	0
All	All	3428	0	3418	46	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 46 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:66:ALA:O	1:C:67:ASP:HB2	1.80	0.80
1:A:136:ASN:ND2	1:C:132:ARG:HH11	1.84	0.75
1:A:132:ARG:HH11	1:C:136:ASN:ND2	1.85	0.74
1:A:109[A]:GLN:CG	1:C:109:GLN:HE22	2.02	0.72
1:B:162:PRO:HA	1:D:162:PRO:O	1.90	0.71

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:129:ASP:OD2	2:A:204:HOH:O[4_564]	1.88	0.32
1:C:65:PRO:CG	2:B:215:HOH:O[7_656]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	115/172 (67%)	111 (96%)	4 (4%)	0	100	100
1	B	112/172 (65%)	105 (94%)	4 (4%)	3 (3%)	6	20
1	C	99/172 (58%)	93 (94%)	4 (4%)	2 (2%)	9	29
1	D	98/172 (57%)	93 (95%)	4 (4%)	1 (1%)	19	50
All	All	424/688 (62%)	402 (95%)	16 (4%)	6 (1%)	14	40

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	64	LEU
1	C	162	PRO
1	B	162	PRO
1	B	154	GLY
1	D	65	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	96/129 (74%)	94 (98%)	2 (2%)	61	88
1	B	95/129 (74%)	91 (96%)	4 (4%)	36	69
1	C	84/129 (65%)	77 (92%)	7 (8%)	14	36
1	D	83/129 (64%)	81 (98%)	2 (2%)	57	86
All	All	358/516 (69%)	343 (96%)	15 (4%)	36	69

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	63	SER
1	C	65	PRO
1	C	145	VAL
1	B	139	LYS
1	C	140	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	84	GLN
1	C	109	GLN
1	D	109	GLN
1	B	136	ASN
1	D	108	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	116/172 (67%)	-0.39	4 (3%) 49 38	32, 48, 70, 134	0
1	B	113/172 (65%)	-0.65	1 (0%) 85 80	33, 47, 65, 101	0
1	C	101/172 (58%)	-0.46	1 (0%) 84 78	34, 47, 70, 113	0
1	D	100/172 (58%)	-0.69	0 100 100	31, 46, 61, 93	0
All	All	430/688 (62%)	-0.55	6 (1%) 78 71	31, 47, 68, 134	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	49	GLY	4.0
1	A	48	PRO	3.9
1	C	63	SER	3.9
1	A	50	GLY	3.5
1	A	51	ALA	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.