



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:56 AM GMT

PDB ID : 3NMT  
Title : Crystal structure of pyrabactin bound abscisic acid receptor PYL2 mutant A93F in complex with type 2C protein phosphatase HAB1  
Authors : Zhou, X.E.; Melcher, K.; Ng, L.-M.; Soon, F.-F.; Xu, Y.; Suino-Powell, K.M.; Kovach, A.; Li, J.; Yong, E.-L.; Xu, H.E.  
Deposited on : 2010-06-22  
Resolution : 2.56 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3324 (2.60-2.52)
Clashscore	102246	3729 (2.60-2.52)
Ramachandran outliers	100387	3673 (2.60-2.52)
Sidechain outliers	100360	3673 (2.60-2.52)
RSRZ outliers	91569	3333 (2.60-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	178	<div> <div>3%</div> <div>67%</div> <div>28%</div> <div>• •</div> </div>
2	B	341	<div> <div>6%</div> <div>58%</div> <div>27%</div> <div>• 13%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PYV	A	900	-	-	-	X
4	MG	B	100	-	-	-	X
4	MG	B	101	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3868 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Absciscic acid receptor PYL2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	176	Total	C	N	O	S	0	0	0
			1401	882	243	272	4			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	GLY	-	EXPRESSION TAG	UNP O80992
A	13	SER	-	EXPRESSION TAG	UNP O80992
A	93	PHE	ALA	ENGINEERED MUTATION	UNP O80992


- Molecule 2 is a protein called Protein phosphatase 2C 16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	296	Total	C	N	O	S	0	0	0
			2305	1453	413	423	16			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	171	GLY	-	EXPRESSION TAG	UNP Q9CAJ0

- Molecule 3 is 4-BROMO-N-(PYRIDIN-2-YLMETHYL)NAPHTHALENE-1-SULFONAMIDE (three-letter code: PYV) (formula: C<sub>16</sub>H<sub>13</sub>BrN<sub>2</sub>O<sub>2</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	Br	C	N	O	S	0	0
			22	1	16	2	2	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Mg	0	0
			2	2		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	50	Total	O	0	0
			50	50		
6	B	78	Total	O	0	0
			78	78		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Absciscic acid receptor PYL2



- Molecule 2: Protein phosphatase 2C 16



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.02Å 66.88Å 144.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.85 – 2.56 29.85 – 2.56	Depositor EDS
% Data completeness (in resolution range)	99.5 (29.85-2.56) 99.5 (29.85-2.56)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.57 (at 2.57Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, $R_{free}$	0.224 , 0.266 0.224 , 0.264	Depositor DCC
$R_{free}$ test set	1431 reflections (7.87%)	DCC
Wilson B-factor (Å <sup>2</sup> )	41.2	Xtriage
Anisotropy	0.961	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 49.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 19612 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3868	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.86% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SO4, PYV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.90	1/1430 (0.1%)	0.76	0/1942
2	B	0.76	0/2348	0.76	1/3165 (0.0%)
All	All	0.82	1/3778 (0.0%)	0.76	1/5107 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	35	CYS	CB-SG	-6.45	1.71	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	243	ASP	CB-CG-OD1	5.54	123.29	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1401	0	1386	38	0
2	B	2305	0	2306	64	0
3	A	22	0	13	0	0
4	B	2	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	10	0	0	0	0
6	A	50	0	0	3	0
6	B	78	0	0	5	0
All	All	3868	0	3705	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:460:ALA:HB1	2:B:461:PRO:HA	1.47	0.95
2:B:353:ARG:HH12	2:B:418:MET:HE1	1.45	0.80
2:B:394:LEU:HD13	6:B:27:HOH:O	1.84	0.76
2:B:301:ILE:HG23	2:B:320:VAL:HG12	1.70	0.74
1:A:176:LYS:HA	1:A:176:LYS:HE3	1.74	0.69
2:B:410:ILE:HD12	2:B:412:GLU:HB3	1.75	0.68
1:A:76:ASP:HB2	6:A:192:HOH:O	1.92	0.68
2:B:187:ILE:HD11	2:B:466:ARG:HB3	1.76	0.66
2:B:394:LEU:HD23	2:B:409:VAL:HG11	1.79	0.65
2:B:310:VAL:HA	2:B:312:SER:HA	1.80	0.63
2:B:190:TRP:CZ3	2:B:470:ILE:HG22	2.34	0.62
1:A:121:LEU:HD22	1:A:151:VAL:HG21	1.83	0.61
2:B:250:VAL:HG23	2:B:325:VAL:HG22	1.83	0.60
2:B:323:GLU:HB2	2:B:403:ARG:HB2	1.82	0.60
2:B:287:ASP:O	2:B:291:THR:HG23	2.01	0.59
2:B:302:GLU:HG3	2:B:403:ARG:HD3	1.86	0.57
2:B:428:ILE:HD12	6:B:59:HOH:O	2.04	0.57
2:B:342:SER:OG	2:B:415:VAL:HG22	2.05	0.57
2:B:192:THR:HG22	2:B:497:ILE:HG13	1.86	0.56
1:A:120:ARG:HH21	1:A:157:ASN:HD21	1.53	0.56
1:A:153:ILE:HD12	1:A:159:GLU:HA	1.88	0.55
2:B:460:ALA:HB1	2:B:461:PRO:CA	2.27	0.55
1:A:67:VAL:HG13	1:A:85:VAL:HG13	1.89	0.54
2:B:206:PHE:O	2:B:255:ARG:HG3	2.07	0.54
1:A:38:LEU:HD23	1:A:148:SER:HB3	1.91	0.53
2:B:376:GLU:OE1	2:B:381:LYS:HD3	2.08	0.53
2:B:262:LEU:O	2:B:266:ILE:HG12	2.09	0.53
2:B:286:TRP:O	2:B:289:VAL:HG12	2.09	0.53
2:B:286:TRP:HA	2:B:289:VAL:HG12	1.91	0.52
2:B:215:LEU:O	2:B:233:HIS:HA	2.09	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:ASN:HD22	1:A:135:ASN:C	2.13	0.52
1:A:93:PHE:HA	1:A:119:HIS:NE2	2.26	0.51
2:B:495:SER:O	2:B:496:ILE:HD13	2.10	0.51
2:B:211:HIS:H	2:B:237:HIS:CD2	2.28	0.51
1:A:65:HIS:HB3	2:B:324:THR:HG21	1.93	0.51
2:B:198:ASN:HD22	2:B:198:ASN:N	2.09	0.51
2:B:409:VAL:HG23	2:B:409:VAL:O	2.10	0.51
2:B:233:HIS:HB2	6:B:75:HOH:O	2.10	0.50
1:A:110:LEU:HD23	1:A:111:SER:N	2.26	0.50
2:B:374:ARG:HH21	2:B:374:ARG:HG3	1.77	0.50
2:B:192:THR:H	2:B:475:GLN:HE22	1.59	0.50
2:B:334:VAL:HG22	2:B:339:ILE:HG12	1.92	0.50
1:A:78:ASP:HB3	6:A:230:HOH:O	2.11	0.49
2:B:350:VAL:HG11	2:B:440:ASN:HB3	1.93	0.49
1:A:120:ARG:NH2	1:A:157:ASN:HD21	2.11	0.49
1:A:39:ILE:HD11	1:A:167:ASP:HA	1.95	0.48
2:B:250:VAL:CG2	2:B:325:VAL:HG22	2.42	0.48
1:A:79:VAL:HG22	1:A:103:VAL:HG23	1.95	0.48
2:B:425:GLU:OE2	2:B:505:ARG:NH2	2.47	0.48
1:A:59:ASN:HA	1:A:61:GLU:OE2	2.12	0.48
2:B:219:MET:HE3	6:B:16:HOH:O	2.14	0.48
2:B:249:LYS:HB3	2:B:321:ALA:HB2	1.96	0.47
2:B:190:TRP:CZ3	2:B:208:VAL:HG11	2.50	0.47
1:A:110:LEU:C	1:A:110:LEU:HD23	2.35	0.47
1:A:108:ARG:HH11	1:A:108:ARG:HG2	1.79	0.46
2:B:211:HIS:H	2:B:237:HIS:HD2	1.64	0.46
1:A:35:CYS:HB2	1:A:153:ILE:HD11	1.98	0.46
2:B:290:PHE:HZ	2:B:333:LEU:HD13	1.80	0.46
2:B:366:PRO:HD3	2:B:396:MET:HA	1.97	0.46
1:A:62:ARG:HH12	1:A:184:ALA:HB1	1.81	0.45
2:B:439:ASN:O	2:B:443:VAL:HG23	2.16	0.45
2:B:188:PRO:HD3	2:B:237:HIS:NE2	2.32	0.45
1:A:135:ASN:HD22	1:A:136:GLN:N	2.14	0.45
1:A:19:GLU:HB2	1:A:20:PRO:HD3	1.99	0.45
2:B:259:HIS:CD2	2:B:259:HIS:H	2.35	0.44
2:B:464:ALA:O	2:B:468:LYS:HE2	2.18	0.44
1:A:91:LEU:CD1	2:B:393:VAL:HG13	2.47	0.44
1:A:35:CYS:HB2	1:A:153:ILE:CD1	2.48	0.44
1:A:178:GLY:O	1:A:182:THR:HB	2.18	0.43
1:A:121:LEU:HD22	1:A:151:VAL:CG2	2.48	0.43
2:B:352:PHE:CD1	2:B:444:CYS:HB3	2.52	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:432:ASP:OD1	2:B:432:ASP:N	2.49	0.43
2:B:233:HIS:N	6:B:75:HOH:O	2.52	0.43
2:B:210:PRO:HA	2:B:237:HIS:HB3	2.00	0.43
1:A:91:LEU:HD22	1:A:91:LEU:N	2.32	0.43
1:A:125:LYS:HB2	1:A:150:THR:HG22	2.00	0.43
2:B:460:ALA:CB	2:B:461:PRO:HA	2.33	0.43
2:B:234:LEU:N	2:B:234:LEU:HD13	2.34	0.42
2:B:341:VAL:O	2:B:415:VAL:HA	2.20	0.42
2:B:461:PRO:HB2	2:B:462:PRO:HD2	2.02	0.42
2:B:496:ILE:C	2:B:497:ILE:HD12	2.40	0.42
1:A:60:PRO:HB2	1:A:64:LYS:HD2	2.00	0.42
2:B:352:PHE:CG	2:B:444:CYS:HB3	2.54	0.42
2:B:447:ALA:O	2:B:451:ILE:HG12	2.19	0.42
2:B:217:ILE:HG23	2:B:233:HIS:N	2.34	0.42
1:A:134:LEU:HD12	1:A:135:ASN:N	2.35	0.42
1:A:175:GLN:O	1:A:179:VAL:HG23	2.20	0.42
1:A:153:ILE:CD1	1:A:159:GLU:HA	2.50	0.41
1:A:64:LYS:O	1:A:67:VAL:HG23	2.20	0.41
1:A:38:LEU:HD13	1:A:38:LEU:C	2.40	0.41
2:B:338:HIS:NE2	2:B:419:PRO:HB3	2.35	0.41
1:A:66:PHE:O	1:A:87:VAL:HG13	2.19	0.41
2:B:331:VAL:HG22	2:B:342:SER:HB2	2.02	0.41
1:A:154:PRO:HG2	2:B:385:TRP:CH2	2.56	0.41
2:B:244:GLY:HA2	2:B:327:SER:HA	2.03	0.40
2:B:217:ILE:HD12	2:B:217:ILE:N	2.36	0.40
1:A:171:LYS:O	1:A:175:GLN:HG3	2.21	0.40
2:B:376:GLU:HA	2:B:380:GLY:O	2.21	0.40
1:A:55:ARG:HG3	6:A:233:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	174/178 (98%)	163 (94%)	9 (5%)	2 (1%)	17	33
2	B	288/341 (84%)	269 (93%)	16 (6%)	3 (1%)	19	36
All	All	462/519 (89%)	432 (94%)	25 (5%)	5 (1%)	17	33

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	462	PRO
1	A	95	THR
2	B	463	LEU
2	B	460	ALA
1	A	156	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	162/163 (99%)	149 (92%)	13 (8%)	15	27
2	B	247/285 (87%)	231 (94%)	16 (6%)	21	38
All	All	409/448 (91%)	380 (93%)	29 (7%)	18	34

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	27	GLN
1	A	59	ASN
1	A	61	GLU
1	A	62	ARG
1	A	72	LEU
1	A	94	SER
1	A	97	THR
1	A	100	LEU
1	A	119	HIS
1	A	135	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	172	LEU
1	A	176	LYS
2	B	213	LEU
2	B	217	ILE
2	B	234	LEU
2	B	235	THR
2	B	268	ARG
2	B	325	VAL
2	B	336	SER
2	B	369	GLU
2	B	386	GLN
2	B	407	PRO
2	B	422	ARG
2	B	434	LEU
2	B	450	ARG
2	B	458	ASN
2	B	466	ARG
2	B	504	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	ASN
1	A	123	ASN
1	A	135	ASN
1	A	157	ASN
1	A	187	HIS
2	B	198	ASN
2	B	259	HIS
2	B	285	GLN
2	B	439	ASN
2	B	475	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PYV	A	900	-	24,24,24	4.65	6 (25%)	34,34,34	1.75	6 (17%)
5	SO4	B	512	-	4,4,4	0.40	0	6,6,6	0.84	0
5	SO4	B	513	-	4,4,4	0.43	0	6,6,6	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PYV	A	900	-	-	0/12/12/12	0/3/3/3
5	SO4	B	512	-	-	0/0/0/0	0/0/0/0
5	SO4	B	513	-	-	0/0/0/0	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	900	PYV	BR-C1	-15.23	1.49	1.90
3	A	900	PYV	C4-S	-15.15	1.61	1.77
3	A	900	PYV	C11-N1	-2.08	1.45	1.47
3	A	900	PYV	C5-C4	2.44	1.40	1.37
3	A	900	PYV	C3-C2	3.16	1.49	1.43
3	A	900	PYV	C4-C3	4.65	1.50	1.43

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	A	900	PYV	O2-S-O1	-6.17	111.35	119.54
3	A	900	PYV	C5-C4-C3	-3.45	117.97	121.04
3	A	900	PYV	C14-C13-N2	-2.45	119.34	123.44
3	A	900	PYV	C13-N2-C12	2.33	120.56	117.36
3	A	900	PYV	C1-C2-C3	2.70	120.22	117.60
3	A	900	PYV	C4-S-N1	3.73	114.14	106.55

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	176/178 (98%)	-0.08	6 (3%) 49 46	21, 38, 82, 102	0
2	B	296/341 (86%)	0.19	19 (6%) 23 20	23, 45, 112, 145	0
All	All	472/519 (90%)	0.09	25 (5%) 30 28	21, 42, 102, 145	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	269	ILE	9.5
1	A	29	GLU	5.3
1	A	136	GLN	5.2
2	B	464	ALA	5.1
2	B	313	SER	4.1
1	A	137	ASP	3.9
2	B	506	LYS	3.3
2	B	234	LEU	3.3
2	B	463	LEU	3.3
1	A	118	GLU	3.2
2	B	320	VAL	3.2
2	B	266	ILE	2.8
2	B	430	ALA	2.6
2	B	216	PRO	2.6
1	A	28	PHE	2.5
2	B	270	LYS	2.3
1	A	117	GLY	2.3
2	B	214	LYS	2.3
2	B	314	ASP	2.3
2	B	303	GLY	2.3
2	B	219	MET	2.3
2	B	372	TYR	2.3
2	B	369	GLU	2.2
2	B	233	HIS	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	312	SER	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	PYV	A	900	22/22	0.62	0.38	7.50	98,113,116,117	0
4	MG	B	100	1/1	0.93	0.34	5.12	35,35,35,35	0
4	MG	B	101	1/1	0.94	0.27	3.16	17,17,17,17	0
5	SO4	B	512	5/5	0.99	0.08	-1.15	26,27,29,31	0
5	SO4	B	513	5/5	0.99	0.07	-	46,47,48,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.