



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:04 PM GMT

PDB ID : 4NMD  
Title : Crystal structure of proline utilization A (PutA) from *Geobacter sulfurreducens* PCA reduced with dithionite  
Authors : Singh, H.; Tanner, J.J.  
Deposited on : 2013-11-14  
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	8664 (2.00-1.96)
Clashscore	102246	9905 (2.00-1.96)
Ramachandran outliers	100387	9792 (2.00-1.96)
Sidechain outliers	100360	9791 (2.00-1.96)
RSRZ outliers	91569	8679 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1005	<div> <div>6%</div> <div>93%</div> <div>• •</div> </div>
1	B	1005	<div> <div>2%</div> <div>91%</div> <div>6% •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	1102	-	-	-	X
3	EDO	A	1104	-	-	-	X
3	EDO	A	1105	-	-	-	X
3	EDO	A	1107	-	-	-	X
3	EDO	A	1108	-	-	-	X
3	EDO	B	1102	-	-	-	X
3	EDO	B	1104	-	-	-	X
3	EDO	B	1106	-	-	-	X
3	EDO	B	1107	-	-	-	X
3	EDO	B	1108	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 16213 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proline dehydrogenase and Delta-1-pyrroline-5-carboxylate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	979	Total	C	N	O	S	0	3	0
			7531	4806	1302	1384	39			
1	B	980	Total	C	N	O	S	0	3	0
			7601	4852	1311	1399	39			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	EXPRESSION TAG	UNP Q746X3
B	0	SER	-	EXPRESSION TAG	UNP Q746X3

- Molecule 2 is DIHYDROFLAVINE-ADENINE DINUCLEOTIDE (three-letter code: FDA) (formula:  $C_{27}H_{35}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	450	Total	O	0	0
			450	450		
4	B	469	Total	O	0	0
			469	469		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proline dehydrogenase and Delta-1-pyrroline-5-carboxylate dehydrogenase



- Molecule 1: Proline dehydrogenase and Delta-1-pyrroline-5-carboxylate dehydrogenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.17Å 151.36Å 175.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.09 – 1.98 49.09 – 1.98	Depositor EDS
% Data completeness (in resolution range)	97.9 (49.09-1.98) 97.8 (49.09-1.98)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.33 (at 1.98Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.165 , 0.199 0.174 , 0.206	Depositor DCC
$R_{free}$ test set	8673 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.1	Xtriage
Anisotropy	0.516	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 46.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 172734 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	16213	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.69% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FDA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.35	0/7710	0.49	0/10459
1	B	0.35	0/7781	0.50	0/10546
All	All	0.35	0/15491	0.49	0/21005

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7531	0	7302	23	0
1	B	7601	0	7432	32	0
2	A	53	0	33	2	0
2	B	53	0	33	1	0
3	A	28	0	42	1	0
3	B	28	0	42	0	0
4	A	450	0	0	0	0
4	B	469	0	0	2	0
All	All	16213	0	14884	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:28:LEU:HD22	1:B:39:MET:HG3	1.76	0.68
1:A:384:VAL:HG22	1:A:403:ILE:HD11	1.76	0.66
1:A:406:TYR:HB2	2:A:1101:FDA:HM72	1.78	0.65
1:B:579:GLN:HE21	1:B:583:LYS:HE3	1.64	0.63
1:A:636:GLU:HB2	3:A:1105:EDO:H22	1.83	0.60
1:B:810:ILE:HD11	1:B:897:VAL:HG11	1.84	0.59
1:B:406:TYR:HB2	2:B:1101:FDA:HM72	1.84	0.58
1:A:13:ARG:NH1	1:A:16:GLU:OE1	2.34	0.57
1:B:448:GLU:OE1	1:B:456:ARG:NH2	2.24	0.52
1:B:284:LYS:NZ	1:B:288:ASP:OD2	2.42	0.52
1:B:143:VAL:HG13	1:B:406:TYR:HA	1.92	0.52
1:A:17:PHE:HE1	1:A:396:ILE:HA	1.75	0.51
1:A:82:MET:SD	1:A:104:LEU:HD21	2.50	0.51
1:B:793[A]:CYS:SG	4:B:1668:HOH:O	2.44	0.50
1:B:484:VAL:HB	1:B:604:TYR:CG	2.47	0.50
1:A:388:MET:O	1:A:389:ALA:C	2.48	0.50
1:A:390:GLU:O	1:A:394:LYS:N	2.38	0.48
1:A:579:GLN:HG3	1:A:583:LYS:HE3	1.96	0.47
1:A:143:VAL:HG13	1:A:406:TYR:HA	1.96	0.47
1:B:566:ASP:OD2	1:B:568:ARG:HB3	2.15	0.47
1:A:579:GLN:HE21	1:A:583:LYS:HG3	1.79	0.47
1:B:204:PRO:HG3	1:B:243:ILE:HG23	1.98	0.46
1:B:266:ARG:HG2	4:B:1539:HOH:O	2.16	0.46
1:B:579:GLN:NE2	1:B:583:LYS:HE3	2.32	0.45
1:A:328:THR:HG21	1:B:1003:VAL:HG11	1.99	0.45
1:B:314:THR:O	1:B:318:LYS:HG2	2.17	0.44
1:A:484:VAL:HB	1:A:604:TYR:CG	2.53	0.44
1:B:579:GLN:HE21	1:B:583:LYS:HG3	1.81	0.44
1:A:28:LEU:O	1:A:36:GLY:HA2	2.18	0.44
1:B:648:ALA:HB1	1:B:726:LEU:HD22	1.99	0.44
1:A:172:LYS:HB2	1:A:237:LEU:HD23	1.99	0.43
1:A:765:ALA:HA	1:A:798:ARG:O	2.18	0.43
1:B:115:MET:O	1:B:118:GLN:HB3	2.18	0.43
1:B:851:GLY:HA3	1:B:885:ILE:HD13	2.00	0.43
1:B:30:ASN:O	1:B:36:GLY:HA3	2.18	0.43
1:B:120:ILE:HG22	1:B:122:GLY:H	1.82	0.43
1:B:172:LYS:HB2	1:B:237:LEU:HD23	1.99	0.43
1:A:113:GLU:O	1:A:117:ARG:HG3	2.18	0.43
1:A:637:LEU:HD21	1:A:997:GLU:HG3	2.02	0.42
1:B:513:PHE:HB3	1:B:706:PHE:HB3	2.01	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1101:FDA:N1	2:A:1101:FDA:O2'	2.49	0.42
1:A:28:LEU:HD23	1:A:35:MET:SD	2.59	0.42
1:B:185:GLY:O	1:B:193:GLY:HA3	2.19	0.42
1:B:377:GLU:HB3	1:B:402:ARG:CZ	2.50	0.41
1:B:83:PRO:HD2	1:B:86:MET:HG3	2.02	0.41
1:B:484:VAL:HB	1:B:604:TYR:CD1	2.56	0.41
1:B:806:TYR:O	1:B:810:ILE:HG12	2.21	0.41
1:B:731:GLY:O	1:B:760:MET:HA	2.21	0.41
1:A:342:ARG:HB2	1:A:369:MET:SD	2.62	0.40
1:B:128:ALA:O	1:B:132:LEU:HG	2.20	0.40
1:A:810:ILE:HD11	1:A:897:VAL:HG11	2.03	0.40
1:A:729:PHE:O	1:A:758:SER:HA	2.21	0.40
1:A:433:LEU:HD23	1:A:433:LEU:HA	1.85	0.40
1:B:852:LYS:NZ	1:B:860:GLU:OE2	2.45	0.40
1:B:729:PHE:O	1:B:758:SER:HA	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	974/1005 (97%)	947 (97%)	26 (3%)	1 (0%)	56	51
1	B	977/1005 (97%)	956 (98%)	21 (2%)	0	100	100
All	All	1951/2010 (97%)	1903 (98%)	47 (2%)	1 (0%)	56	51

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	79	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	753/821 (92%)	742 (98%)	11 (2%)	72	73
1	B	770/821 (94%)	759 (99%)	11 (1%)	74	75
All	All	1523/1642 (93%)	1501 (99%)	22 (1%)	72	75

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	104	LEU
1	A	177	SER
1	A	278	TYR
1	A	388	MET
1	A	393	ARG
1	A	403	ILE
1	A	439	GLU
1	A	585	LEU
1	A	891	PHE
1	A	936	PHE
1	A	1003	VAL
1	B	20	SER
1	B	79	GLU
1	B	278	TYR
1	B	412	MET
1	B	520	ARG
1	B	527	THR
1	B	585	LEU
1	B	759	GLU
1	B	891	PHE
1	B	936	PHE
1	B	993	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	579	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FDA	A	1101	-	48,58,58	2.73	15 (31%)	54,89,89	2.21	11 (20%)
3	EDO	A	1102	-	3,3,3	0.44	0	2,2,2	0.56	0
3	EDO	A	1103	-	3,3,3	0.48	0	2,2,2	0.40	0
3	EDO	A	1104	-	3,3,3	0.45	0	2,2,2	0.45	0
3	EDO	A	1105	-	3,3,3	0.47	0	2,2,2	0.49	0
3	EDO	A	1106	-	3,3,3	0.45	0	2,2,2	0.51	0
3	EDO	A	1107	-	3,3,3	0.49	0	2,2,2	0.41	0
3	EDO	A	1108	-	3,3,3	0.45	0	2,2,2	0.31	0
2	FDA	B	1101	-	48,58,58	2.70	17 (35%)	54,89,89	2.07	11 (20%)
3	EDO	B	1102	-	3,3,3	0.48	0	2,2,2	0.49	0
3	EDO	B	1103	-	3,3,3	0.46	0	2,2,2	0.64	0
3	EDO	B	1104	-	3,3,3	0.49	0	2,2,2	0.33	0
3	EDO	B	1105	-	3,3,3	0.43	0	2,2,2	0.55	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	B	1106	-	3,3,3	0.52	0	2,2,2	0.45	0
3	EDO	B	1107	-	3,3,3	0.54	0	2,2,2	0.15	0
3	EDO	B	1108	-	3,3,3	0.50	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FDA	A	1101	-	-	0/30/50/50	0/6/6/6
3	EDO	A	1102	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1103	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1104	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1105	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1106	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1107	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1108	-	-	0/1/1/1	0/0/0/0
2	FDA	B	1101	-	-	0/30/50/50	0/6/6/6
3	EDO	B	1102	-	-	0/1/1/1	0/0/0/0
3	EDO	B	1103	-	-	0/1/1/1	0/0/0/0
3	EDO	B	1104	-	-	0/1/1/1	0/0/0/0
3	EDO	B	1105	-	-	0/1/1/1	0/0/0/0
3	EDO	B	1106	-	-	0/1/1/1	0/0/0/0
3	EDO	B	1107	-	-	0/1/1/1	0/0/0/0
3	EDO	B	1108	-	-	0/1/1/1	0/0/0/0

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1101	FDA	C9A-C5X	-3.41	1.35	1.42
2	A	1101	FDA	C9A-C5X	-2.87	1.36	1.42
2	B	1101	FDA	PA-O2A	-2.41	1.44	1.54
2	A	1101	FDA	PA-O2A	-2.36	1.44	1.54
2	B	1101	FDA	C4X-C10	-2.25	1.36	1.41
2	B	1101	FDA	PA-O5B	-2.10	1.49	1.59
2	B	1101	FDA	C2B-C3B	-2.09	1.47	1.53
2	A	1101	FDA	O3B-C3B	-2.06	1.38	1.43
2	A	1101	FDA	C2A-N3A	2.50	1.36	1.32
2	B	1101	FDA	C2A-N3A	2.74	1.37	1.32
2	B	1101	FDA	C6A-N6A	3.01	1.44	1.34
2	A	1101	FDA	C6A-N6A	3.09	1.44	1.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1101	FDA	C10-N10	3.25	1.42	1.39
2	B	1101	FDA	C9-C9A	3.32	1.48	1.40
2	A	1101	FDA	C9-C9A	3.58	1.48	1.40
2	A	1101	FDA	C4-C4X	3.86	1.49	1.41
2	A	1101	FDA	C4-N3	4.04	1.40	1.33
2	B	1101	FDA	C4-N3	4.18	1.40	1.33
2	B	1101	FDA	C4-C4X	4.23	1.49	1.41
2	A	1101	FDA	C10-N10	4.50	1.44	1.39
2	B	1101	FDA	C9A-N10	4.71	1.45	1.38
2	A	1101	FDA	C9A-N10	4.89	1.45	1.38
2	B	1101	FDA	C6-C5X	4.93	1.49	1.41
2	A	1101	FDA	C6-C5X	5.11	1.49	1.41
2	B	1101	FDA	O4-C4	5.36	1.37	1.24
2	B	1101	FDA	C10-N1	5.45	1.44	1.35
2	A	1101	FDA	O4-C4	5.64	1.38	1.24
2	A	1101	FDA	C10-N1	5.86	1.45	1.35
2	B	1101	FDA	C5X-N5	6.50	1.45	1.35
2	A	1101	FDA	C5X-N5	6.60	1.45	1.35
2	A	1101	FDA	C4X-N5	7.25	1.44	1.33
2	B	1101	FDA	C4X-N5	7.51	1.45	1.33

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	FDA	N3A-C2A-N1A	-11.21	120.31	128.89
2	B	1101	FDA	N3A-C2A-N1A	-10.29	121.02	128.89
2	A	1101	FDA	C1B-N9A-C4A	-3.48	121.69	126.94
2	A	1101	FDA	C4X-C4-N3	-3.35	119.01	123.59
2	B	1101	FDA	C4X-C4-N3	-2.96	119.53	123.59
2	A	1101	FDA	P-O3P-PA	-2.86	124.69	132.73
2	B	1101	FDA	O2A-PA-O5B	-2.51	95.80	108.46
2	A	1101	FDA	O2P-P-O5'	-2.21	97.33	108.46
2	B	1101	FDA	C4A-C5A-N7A	-2.20	107.45	109.48
2	B	1101	FDA	C1B-N9A-C4A	-2.17	123.67	126.94
2	A	1101	FDA	C4A-C5A-N7A	-2.13	107.52	109.48
2	A	1101	FDA	O2A-PA-O3P	-2.01	95.96	105.09
2	B	1101	FDA	C9A-C5X-N5	-2.00	119.40	122.36
2	B	1101	FDA	C4-C4X-N5	2.05	121.20	118.72
2	A	1101	FDA	O5'-P-O1P	2.13	117.88	109.62
2	B	1101	FDA	C5X-C9A-N10	2.33	119.39	117.62
2	B	1101	FDA	O5'-P-O1P	2.43	119.03	109.62
2	A	1101	FDA	C5X-C9A-N10	3.03	119.92	117.62

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	FDA	C1'-N10-C9A	3.22	122.47	118.86
2	B	1101	FDA	C1'-N10-C9A	4.35	123.75	118.86
2	B	1101	FDA	C4-N3-C2	5.24	119.78	115.25
2	A	1101	FDA	C4-N3-C2	6.11	120.53	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1101	FDA	2	0
3	A	1105	EDO	1	0
2	B	1101	FDA	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	979/1005 (97%)	0.12	60 (6%) 25 28	12, 22, 55, 94	0
1	B	980/1005 (97%)	-0.10	16 (1%) 74 77	11, 21, 41, 74	0
All	All	1959/2010 (97%)	0.01	76 (3%) 43 47	11, 21, 47, 94	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	438	ALA	8.5
1	A	389	ALA	7.4
1	A	85	PHE	7.0
1	A	86	MET	6.3
1	A	392	VAL	6.0
1	A	83	PRO	5.4
1	A	41	TRP	5.3
1	A	84	ALA	5.1
1	A	80	GLN	5.1
1	A	104	LEU	5.0
1	A	78	ASN	5.0
1	B	472	LEU	5.0
1	A	463	ALA	4.9
1	A	82	MET	4.8
1	A	460	ALA	4.8
1	A	471	GLY	4.6
1	A	178	TRP	4.5
1	A	103	VAL	4.5
1	B	462	ALA	4.4
1	A	81	ASP	4.4
1	B	463	ALA	4.1
1	A	473	GLY	4.0
1	A	439	GLU	3.9
1	A	472	LEU	3.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	391	PRO	3.9
1	B	471	GLY	3.8
1	A	34	TRP	3.7
1	A	28	LEU	3.7
1	B	473	GLY	3.5
1	A	102	ALA	3.4
1	B	84	ALA	3.4
1	A	462	ALA	3.3
1	A	396	ILE	3.2
1	A	107	VAL	3.2
1	A	441	ALA	3.1
1	A	186	GLY	3.1
1	A	437	PHE	3.1
1	B	83	PRO	3.0
1	B	81	ASP	2.9
1	A	38	ALA	2.9
1	A	440	ASP	2.8
1	B	41	TRP	2.8
1	A	865	ALA	2.8
1	B	459	ALA	2.7
1	A	0	SER	2.7
1	B	3	ASN	2.6
1	A	1	MET	2.6
1	B	104	LEU	2.6
1	A	459	ALA	2.6
1	A	184	LYS	2.5
1	A	176	GLY	2.5
1	B	4	SER	2.5
1	A	2	LEU	2.5
1	A	79	GLU	2.5
1	A	393	ARG	2.5
1	A	188	PRO	2.4
1	B	80	GLN	2.4
1	A	190	LEU	2.4
1	A	189	GLY	2.4
1	A	45	ASN	2.3
1	A	390	GLU	2.2
1	B	178	TRP	2.2
1	A	44	GLN	2.2
1	A	400	ALA	2.2
1	A	177	SER	2.2
1	A	388	MET	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	30	ASN	2.1
1	A	27	SER	2.1
1	A	35	MET	2.1
1	A	36	GLY	2.1
1	A	183	GLY	2.1
1	A	429	ASN	2.0
1	A	173	LYS	2.0
1	A	866	GLY	2.0
1	A	26	PRO	2.0
1	B	461	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EDO	A	1104	4/4	0.97	0.15	6.21	23,26,32,36	0
3	EDO	A	1105	4/4	0.88	0.20	5.59	29,31,38,41	0
3	EDO	B	1104	4/4	0.98	0.10	4.35	23,31,31,32	0
3	EDO	B	1107	4/4	0.93	0.14	3.04	30,36,37,40	0
3	EDO	A	1107	4/4	0.94	0.18	3.03	17,28,33,35	0
3	EDO	A	1108	4/4	0.95	0.16	2.73	15,20,22,25	0
3	EDO	B	1102	4/4	0.96	0.14	2.41	21,22,24,26	0
3	EDO	B	1106	4/4	0.98	0.15	2.37	21,27,30,31	0
3	EDO	B	1108	4/4	0.95	0.13	2.23	24,26,26,26	0
3	EDO	A	1102	4/4	0.97	0.14	2.11	17,20,21,26	0
3	EDO	A	1106	4/4	0.95	0.12	1.59	21,35,36,40	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EDO	B	1103	4/4	0.95	0.11	0.87	20,20,23,27	0
3	EDO	A	1103	4/4	0.96	0.12	0.83	26,29,32,36	0
2	FDA	B	1101	53/53	0.97	0.10	0.58	14,18,24,29	0
2	FDA	A	1101	53/53	0.97	0.10	-0.18	8,19,28,32	0
3	EDO	B	1105	4/4	0.89	0.17	-	41,42,43,45	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.