



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 2, 2016 – 12:00 AM GMT

PDB ID : 7NN9  
Title : NATIVE INFLUENZA VIRUS NEURAMINIDASE SUBTYPE N9 (TERN)  
Authors : Varghese, J.N.; Colman, P.M.  
Deposited on : 1995-03-15  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MAN	A	471(C)	X	-	-	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3400 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NEURAMINIDASE N9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	388	3067	1914	538	592	23	31	0	0

- Molecule 2 is a polymer of unknown type called SUGAR (7-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	7	83	46	2	35	0	0

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Ca 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	221	Total 221	O 221	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: NEURAMINIDASE N9



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	182.80 Å   182.80 Å   182.80 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	6.00 – 2.00 5.92 – 2.19	Depositor EDS
% Data completeness (in resolution range)	(Not available) (6.00-2.00) 68.9 (5.92-2.19)	Depositor EDS
$R_{merge}$	0.01	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.152 , (Not available) 0.150 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	13.0	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 56.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>1</sup>	$\langle  L  \rangle = 0.37$ , $\langle L^2 \rangle = 0.19$	Xtriage
Outliers	0 of 17687 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3400	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.55	0/3150	0.80	2/4290 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	1	0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	364	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	A	364	ARG	NE-CZ-NH2	-5.12	117.74	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	471(C)	MAN	C1

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3067	0	2893	34	0
2	A	83	0	70	2	0
3	A	28	0	26	0	0
4	A	1	0	0	0	0
5	A	221	0	0	2	2
All	All	3400	0	2989	34	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87:LEU:H	1:A:233:HIS:HD2	1.31	0.78
1:A:292:ARG:HH21	1:A:294:ASN:ND2	1.83	0.75
1:A:98:HIS:HE1	1:A:419:ARG:HH11	1.34	0.74
1:A:168:THR:H	1:A:170:ASN:HD21	1.35	0.73
1:A:98:HIS:CE1	1:A:419:ARG:HH11	2.12	0.67
1:A:364:ARG:HD2	1:A:375:GLU:OE2	2.01	0.60
1:A:84:PHE:CE1	1:A:187:LYS:HD2	2.37	0.59
1:A:273:LYS:HG3	1:A:340:PRO:HD3	1.88	0.56
1:A:97:TRP:H	1:A:395:GLN:HE22	1.55	0.55
1:A:328:PRO:HG3	1:A:343:GLY:HA3	1.91	0.52
1:A:98:HIS:CE1	1:A:419:ARG:NH1	2.78	0.52
1:A:292:ARG:HH21	1:A:294:ASN:HD21	1.56	0.51
1:A:292:ARG:HE	1:A:294:ASN:HD22	1.59	0.50
1:A:422:PHE:CZ	1:A:446:MET:HG3	2.46	0.50
1:A:152:ARG:HG2	1:A:178:TRP:CD2	2.47	0.50
1:A:240:VAL:HG22	1:A:254:ILE:HD13	1.96	0.47
1:A:144:HIS:HD2	5:A:637:HOH:O	1.96	0.47
1:A:205:ILE:N	1:A:205:ILE:HD12	2.31	0.46
1:A:465:GLU:CD	1:A:465:GLU:H	2.20	0.45
1:A:221:ASN:HB3	1:A:244:GLY:HA2	1.98	0.45
1:A:220:ARG:HH21	2:A:469(A):NAG:H82	1.82	0.44
1:A:141:ARG:NH2	1:A:467:PHE:HA	2.33	0.44
1:A:87:LEU:H	1:A:233:HIS:CD2	2.22	0.44
1:A:131:PHE:CE1	1:A:163:LEU:HD12	2.54	0.43
1:A:170:ASN:H	1:A:170:ASN:HD22	1.66	0.43
1:A:367:SER:HB2	1:A:400:ASN:HD21	1.83	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:ILE:HG12	1:A:203:ALA:HB2	2.00	0.42
1:A:168:THR:H	1:A:170:ASN:ND2	2.09	0.42
1:A:318:CYS:SG	1:A:383:LEU:O	2.78	0.41
1:A:121:TYR:CG	1:A:228:SER:HA	2.56	0.41
1:A:278:CYS:HB3	1:A:289:CYS:HB3	2.02	0.41
1:A:220:ARG:NH2	2:A:469(A):NAG:H82	2.35	0.41
1:A:152:ARG:HG2	1:A:178:TRP:CG	2.56	0.40
1:A:144:HIS:CD2	5:A:637:HOH:O	2.74	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:775:HOH:O	5:A:777:HOH:O[48_555]	0.54	1.66
5:A:785:HOH:O	5:A:785:HOH:O[48_555]	1.35	0.85

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/388 (100%)	366 (95%)	20 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/341 (100%)	333 (98%)	8 (2%)	58	60

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	ASP
1	A	141	ARG
1	A	170	ASN
1	A	204	VAL
1	A	230	CYS
1	A	304	ARG
1	A	364	ARG
1	A	446	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN
1	A	98	HIS
1	A	144	HIS
1	A	170	ASN
1	A	221	ASN
1	A	233	HIS
1	A	294	ASN
1	A	344	ASN
1	A	346	ASN
1	A	392	GLN
1	A	395	GLN
1	A	400	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

7 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	469(A)	1,2	14,14,15	1.70	2 (14%)	15,19,21	1.22	2 (13%)
2	NAG	A	470(B)	2	14,14,15	1.30	2 (14%)	15,19,21	2.15	6 (40%)
2	MAN	A	471(C)	2	11,11,12	1.32	1 (9%)	14,15,17	1.52	3 (21%)
2	MAN	A	472(D)	2	11,11,12	1.38	1 (9%)	14,15,17	2.08	6 (42%)
2	MAN	A	473(E)	2	11,11,12	0.88	0	14,15,17	1.21	2 (14%)
2	MAN	A	474(F)	2	11,11,12	1.32	2 (18%)	14,15,17	1.44	2 (14%)
2	MAN	A	475(G)	2	11,11,12	1.90	2 (18%)	14,15,17	3.27	8 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	469(A)	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	470(B)	2	-	0/6/23/26	0/1/1/1
2	MAN	A	471(C)	2	1/1/4/5	0/2/19/22	0/1/1/1
2	MAN	A	472(D)	2	-	0/2/19/22	0/1/1/1
2	MAN	A	473(E)	2	-	0/2/19/22	0/1/1/1
2	MAN	A	474(F)	2	-	0/2/19/22	0/1/1/1
2	MAN	A	475(G)	2	-	0/2/19/22	1/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	475(G)	MAN	C4-C5	-4.96	1.42	1.53
2	A	471(C)	MAN	C4-C5	-3.41	1.45	1.53
2	A	472(D)	MAN	C4-C5	-3.31	1.46	1.53
2	A	475(G)	MAN	O5-C1	-2.39	1.39	1.43
2	A	470(B)	NAG	C1-C2	-2.30	1.49	1.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	470(B)	NAG	C2-N2	-2.27	1.42	1.46
2	A	474(F)	MAN	C4-C3	2.21	1.58	1.52
2	A	474(F)	MAN	O5-C5	2.29	1.48	1.43
2	A	469(A)	NAG	C4-C5	2.84	1.59	1.53
2	A	469(A)	NAG	C1-C2	4.44	1.58	1.52

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	475(G)	MAN	O2-C2-C1	-4.48	100.22	109.21
2	A	475(G)	MAN	C2-C3-C4	-4.05	104.16	111.04
2	A	470(B)	NAG	O7-C7-C8	-3.75	115.18	122.06
2	A	475(G)	MAN	O4-C4-C5	-3.21	100.74	109.24
2	A	475(G)	MAN	O3-C3-C2	-2.80	104.94	110.00
2	A	470(B)	NAG	C2-N2-C7	-2.72	119.54	123.04
2	A	471(C)	MAN	C1-C2-C3	-2.70	106.34	109.54
2	A	472(D)	MAN	O2-C2-C1	-2.67	103.85	109.21
2	A	471(C)	MAN	O3-C3-C4	-2.55	104.58	110.34
2	A	470(B)	NAG	C4-C3-C2	-2.53	107.30	111.23
2	A	470(B)	NAG	O6-C6-C5	-2.31	103.68	111.33
2	A	472(D)	MAN	C6-C5-C4	-2.31	107.31	113.02
2	A	473(E)	MAN	O4-C4-C3	-2.24	105.29	110.34
2	A	472(D)	MAN	C1-C2-C3	-2.22	106.91	109.54
2	A	472(D)	MAN	O6-C6-C5	-2.21	104.03	111.33
2	A	474(F)	MAN	C3-C4-C5	-2.18	106.40	110.20
2	A	469(A)	NAG	C4-C3-C2	-2.13	107.92	111.23
2	A	471(C)	MAN	O6-C6-C5	-2.02	104.66	111.33
2	A	472(D)	MAN	C3-C4-C5	-2.00	106.71	110.20
2	A	475(G)	MAN	O2-C2-C3	2.23	114.60	110.12
2	A	473(E)	MAN	C3-C4-C5	2.23	114.09	110.20
2	A	475(G)	MAN	O4-C4-C3	2.39	115.71	110.34
2	A	469(A)	NAG	C1-O5-C5	2.94	115.98	112.25
2	A	475(G)	MAN	C1-C2-C3	3.38	113.53	109.54
2	A	470(B)	NAG	C8-C7-N2	3.62	123.03	116.11
2	A	470(B)	NAG	C6-C5-C4	3.63	121.96	113.02
2	A	474(F)	MAN	C1-O5-C5	3.90	117.19	112.25
2	A	472(D)	MAN	C1-O5-C5	4.87	118.43	112.25
2	A	475(G)	MAN	C1-O5-C5	7.80	122.15	112.25

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	471(C)	MAN	C1

There are no torsion outliers.

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	475(G)	MAN	C1-C2-C3-C4-C5-O5

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	469(A)	NAG	2	0

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	476(A)	1	14,14,15	1.05	0	15,19,21	1.70	5 (33%)
3	NAG	A	477(A)	1	14,14,15	1.42	1 (7%)	15,19,21	2.32	6 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	476(A)	1	-	0/6/23/26	0/1/1/1
3	NAG	A	477(A)	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	477(A)	NAG	C1-C2	-4.63	1.46	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	477(A)	NAG	C2-N2-C7	-5.07	116.53	123.04
3	A	476(A)	NAG	C4-C3-C2	-3.49	105.80	111.23
3	A	476(A)	NAG	C6-C5-C4	-3.19	105.15	113.02
3	A	477(A)	NAG	C3-C4-C5	-2.87	105.19	110.20
3	A	477(A)	NAG	O7-C7-C8	-2.50	117.48	122.06
3	A	476(A)	NAG	C3-C2-N2	-2.40	104.80	110.56
3	A	476(A)	NAG	C2-N2-C7	-2.00	120.46	123.04
3	A	476(A)	NAG	O3-C3-C2	2.14	113.34	109.11
3	A	477(A)	NAG	C8-C7-N2	2.37	120.64	116.11
3	A	477(A)	NAG	C1-O5-C5	2.80	115.81	112.25
3	A	477(A)	NAG	C6-C5-C4	4.42	123.93	113.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	388/388 (100%)	-0.79	0 100 100	10, 17, 28, 38	8 (2%)

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	469(A)	14/15	0.95	0.12	1.51	21,22,25,25	0
2	MAN	A	471(C)	11/12	0.96	0.09	1.33	27,29,35,42	0
2	MAN	A	474(F)	11/12	0.96	0.09	0.08	23,24,25,25	0
2	MAN	A	472(D)	11/12	0.92	0.12	-	27,28,29,31	0
2	MAN	A	475(G)	11/12	0.70	0.36	-	48,52,55,55	0
2	MAN	A	473(E)	11/12	0.95	0.10	-	25,26,27,28	0
2	NAG	A	470(B)	14/15	0.95	0.10	-	21,23,26,27	0

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	CA	A	600	1/1	0.99	0.03	-2.21	23,23,23,23	0
3	NAG	A	477(A)	14/15	0.89	0.16	-	37,40,41,42	0
3	NAG	A	476(A)	14/15	0.93	0.12	-	28,30,32,33	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.