



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:38 AM GMT

PDB ID : 2NPX
Title : NADH BINDING SITE AND CATALYSIS OF NADH PEROXIDASE
Authors : Stehle, T.; Claiborne, A.; Schulz, G.E.
Deposited on : 1992-05-29
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	447	<div> <div></div> <div>79%</div> <div>17%</div> <div>..</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3927 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH PEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	447	3493	2225	573	683	12	0	0	0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	337	Total	O	0	0
			337	337		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NADH PEROXIDASE



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	77.20Å 134.50Å 145.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.40 33.48 – 2.41	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.40) 93.2 (33.48-2.41)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 2.42Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.159 , (Not available) 0.147 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	23.6	Xtriage
Anisotropy	0.405	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 76.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 27870 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3927	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: OCS, FAD, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/3547	1.55	48/4807 (1.0%)

There are no bond length outliers.

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	303	ARG	NE-CZ-NH1	15.02	127.81	120.30
1	A	59	ARG	NE-CZ-NH1	11.11	125.86	120.30
1	A	30	TRP	CD1-CG-CD2	9.05	113.54	106.30
1	A	374	TRP	CD1-CG-CD2	9.02	113.52	106.30
1	A	134	ARG	NE-CZ-NH2	-8.54	116.03	120.30
1	A	132	ARG	NE-CZ-NH1	8.00	124.30	120.30
1	A	69	ARG	NE-CZ-NH1	7.73	124.16	120.30
1	A	374	TRP	CE2-CD2-CG	-7.63	101.19	107.30
1	A	432	TRP	CD1-CG-CD2	7.61	112.39	106.30
1	A	136	TRP	CD1-CG-CD2	7.54	112.33	106.30
1	A	136	TRP	CE2-CD2-CG	-7.37	101.40	107.30
1	A	30	TRP	CE2-CD2-CG	-7.23	101.51	107.30
1	A	374	TRP	CB-CG-CD1	-7.14	117.71	127.00
1	A	250	TRP	CD1-CG-CD2	7.12	112.00	106.30
1	A	432	TRP	CE2-CD2-CG	-7.12	101.60	107.30
1	A	250	TRP	CE2-CD2-CG	-6.94	101.75	107.30
1	A	303	ARG	NE-CZ-NH2	-6.87	116.86	120.30
1	A	69	ARG	NE-CZ-NH2	-6.86	116.87	120.30
1	A	244	VAL	N-CA-CB	-6.73	96.69	111.50
1	A	279	VAL	N-CA-CB	-6.71	96.75	111.50
1	A	362	TYR	CB-CG-CD2	-6.56	117.06	121.00
1	A	242	VAL	N-CA-CB	-6.46	97.28	111.50
1	A	100	ARG	NE-CZ-NH2	-6.46	117.07	120.30
1	A	30	TRP	CG-CD1-NE1	-6.39	103.71	110.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	134	ARG	NE-CZ-NH1	6.38	123.49	120.30
1	A	59	ARG	NE-CZ-NH2	-6.16	117.22	120.30
1	A	189	LEU	CA-CB-CG	6.12	129.37	115.30
1	A	104	TYR	CB-CG-CD2	-6.09	117.35	121.00
1	A	47	TYR	CB-CG-CD2	-6.05	117.37	121.00
1	A	270	ARG	CB-CG-CD	-6.03	95.92	111.60
1	A	132	ARG	NE-CZ-NH2	-5.97	117.31	120.30
1	A	374	TRP	CG-CD1-NE1	-5.83	104.27	110.10
1	A	100	ARG	NE-CZ-NH1	5.71	123.15	120.30
1	A	294	VAL	CG1-CB-CG2	-5.59	101.95	110.90
1	A	374	TRP	CG-CD2-CE3	5.59	138.93	133.90
1	A	317	VAL	N-CA-CB	-5.52	99.35	111.50
1	A	188	TYR	CB-CG-CD1	-5.43	117.74	121.00
1	A	124	ASP	CA-C-N	-5.42	105.28	117.20
1	A	30	TRP	CG-CD2-CE3	5.39	138.75	133.90
1	A	136	TRP	CG-CD2-CE3	5.35	138.71	133.90
1	A	429	ASP	CB-CG-OD1	5.25	123.02	118.30
1	A	140	LEU	CA-CB-CG	5.21	127.29	115.30
1	A	391	LEU	CA-CB-CG	5.10	127.03	115.30
1	A	284	LEU	CA-CB-CG	5.08	127.00	115.30
1	A	60	TYR	CB-CG-CD2	-5.05	117.97	121.00
1	A	202	GLU	CA-CB-CG	-5.04	102.31	113.40
1	A	434	ILE	CA-CB-CG1	-5.03	101.44	111.00
1	A	222	ARG	NE-CZ-NH1	5.00	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3493	0	3489	41	0
2	A	53	0	31	2	0
3	A	44	0	26	2	0
4	A	337	0	0	7	0
All	All	3927	0	3546	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:VAL:HG22	1:A:305:GLN:HB3	1.73	0.69
1:A:87:HIS:HE1	1:A:275:ASP:OD2	1.84	0.61
1:A:45:GLN:HG3	4:A:1153:HOH:O	2.02	0.59
1:A:345:MET:HG2	4:A:1101:HOH:O	2.02	0.58
1:A:81:ALA:HB3	1:A:90:THR:HG23	1.87	0.57
1:A:268:TYR:HB2	1:A:270:ARG:HD2	1.87	0.55
1:A:88:GLN:HG2	1:A:103:ASN:OD1	2.08	0.53
1:A:192:GLU:HG3	1:A:357:THR:HG21	1.91	0.52
1:A:312:ASN:ND2	1:A:317:VAL:H	2.09	0.51
1:A:115:PRO:HB2	1:A:130:LEU:HG	1.93	0.50
1:A:216:ARG:HD3	4:A:987:HOH:O	2.12	0.49
1:A:208:ILE:HG22	1:A:210:THR:HG23	1.95	0.48
1:A:144:THR:HG22	1:A:145:VAL:HG13	1.96	0.48
1:A:39:PHE:CD1	1:A:134:ARG:HG2	2.49	0.47
1:A:388:GLY:HA2	1:A:408:ILE:HD11	1.98	0.46
1:A:312:ASN:HD21	1:A:318:LYS:N	2.13	0.45
1:A:23:HIS:HE1	4:A:933:HOH:O	1.99	0.45
1:A:277:PHE:CD1	1:A:277:PHE:N	2.84	0.45
1:A:181:LEU:HD13	4:A:1146:HOH:O	2.17	0.45
1:A:132:ARG:HD3	4:A:1095:HOH:O	2.16	0.44
1:A:270:ARG:NH2	1:A:316:PRO:HG3	2.32	0.44
1:A:42:OCS:OD2	2:A:449:FAD:C10	2.65	0.44
1:A:401:ILE:HD12	1:A:401:ILE:HA	1.70	0.44
1:A:113:ALA:HB1	1:A:244:VAL:HG13	2.00	0.43
1:A:76:ASN:O	1:A:94:LEU:HB2	2.18	0.43
1:A:141:LYS:O	1:A:144:THR:HB	2.18	0.43
1:A:380:ASP:O	1:A:384:THR:HA	2.19	0.42
1:A:312:ASN:HD21	1:A:318:LYS:H	1.68	0.42
1:A:85:LYS:O	1:A:87:HIS:HD2	2.03	0.42
1:A:359:VAL:HG12	1:A:372:LYS:HE3	2.01	0.42
1:A:372:LYS:HE2	1:A:372:LYS:HB3	1.88	0.41
1:A:83:GLN:HE22	1:A:90:THR:HG22	1.84	0.41
1:A:242:VAL:HG22	3:A:818:NAD:C5A	2.50	0.41
1:A:218:GLU:HB2	1:A:225:LYS:HB2	2.02	0.41
1:A:23:HIS:CE1	4:A:933:HOH:O	2.74	0.40
1:A:42:OCS:OD2	2:A:449:FAD:C4X	2.69	0.40
1:A:129:TYR:HD2	1:A:140:LEU:HD13	1.86	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:PHE:CD2	1:A:39:PHE:HB2	2.57	0.40
1:A:159:TYR:HB3	3:A:818:NAD:C5N	2.52	0.40
1:A:312:ASN:HD22	1:A:317:VAL:H	1.68	0.40
1:A:63:GLY:O	1:A:67:GLU:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	444/447 (99%)	428 (96%)	15 (3%)	1 (0%)	52	69

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	123	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	375/375 (100%)	343 (92%)	32 (8%)	13	20

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	LEU
1	A	21	ASN
1	A	59	ARG
1	A	86	GLU
1	A	90	THR
1	A	94	LEU
1	A	114	VAL
1	A	118	LEU
1	A	123	LYS
1	A	124	ASP
1	A	130	LEU
1	A	134	ARG
1	A	140	LEU
1	A	144	THR
1	A	148	GLU
1	A	154	VAL
1	A	174	LYS
1	A	189	LEU
1	A	205	ASN
1	A	222	ARG
1	A	227	VAL
1	A	231	ASN
1	A	237	LEU
1	A	242	VAL
1	A	244	VAL
1	A	262	LEU
1	A	270	ARG
1	A	291	ASP
1	A	317	VAL
1	A	345	MET
1	A	365	ASP
1	A	440	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	ASN
1	A	23	HIS
1	A	83	GLN
1	A	87	HIS
1	A	142	GLN
1	A	204	ASN
1	A	205	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	288	ASN
1	A	295	ASN
1	A	312	ASN
1	A	371	GLN
1	A	385	GLN
1	A	445	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	OCS	A	42	1	7,8,9	1.81	2 (28%)	7,11,13	6.95	5 (71%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCS	A	42	1	-	0/4/7/9	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	42	OCS	OD2-SG	-3.83	1.36	1.46
1	A	42	OCS	CB-SG	2.26	1.81	1.77

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	42	OCS	OD2-SG-OD1	-3.60	103.23	111.61
1	A	42	OCS	OD2-SG-OD3	-3.51	103.44	111.61
1	A	42	OCS	O-C-CA	-2.48	119.04	125.49
1	A	42	OCS	OD1-SG-CB	12.22	117.24	106.94
1	A	42	OCS	OD3-SG-CB	12.25	117.27	106.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	42	OCS	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	449	-	48,58,58	1.23	4 (8%)	54,89,89	1.81	8 (14%)
3	NAD	A	818	-	38,48,48	1.93	6 (15%)	47,73,73	1.74	10 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	449	-	-	0/30/50/50	0/6/6/6
3	NAD	A	818	-	-	0/22/62/62	0/5/5/5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	818	NAD	C3N-C7N	-2.74	1.46	1.50
3	A	818	NAD	PN-O2N	-2.35	1.44	1.54
2	A	449	FAD	P-O5'	-2.10	1.49	1.59
2	A	449	FAD	C4-C4X	2.36	1.46	1.41
3	A	818	NAD	O4D-C1D	2.55	1.44	1.41
3	A	818	NAD	C2N-C3N	2.59	1.42	1.39
2	A	449	FAD	O4B-C1B	3.46	1.45	1.41
2	A	449	FAD	C4-N3	3.49	1.39	1.33
3	A	818	NAD	C5N-C4N	5.90	1.51	1.38
3	A	818	NAD	C4N-C3N	6.86	1.51	1.39

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	449	FAD	C4X-C4-N3	-6.25	115.04	123.59
3	A	818	NAD	C5N-C4N-C3N	-4.10	115.18	120.33
2	A	449	FAD	C4X-C10-N10	-3.50	118.46	120.52
3	A	818	NAD	C2B-C1B-N9A	-3.12	109.52	114.29
2	A	449	FAD	C2B-C1B-N9A	-2.77	110.07	114.29
2	A	449	FAD	O4'-C4'-C5'	-2.60	104.53	110.19
3	A	818	NAD	C4B-O4B-C1B	-2.56	106.91	109.72
3	A	818	NAD	C4A-C5A-N7A	-2.47	107.20	109.48
3	A	818	NAD	O7N-C7N-C3N	-2.24	117.14	119.59
2	A	449	FAD	C4X-N5-C5X	2.01	119.08	116.76
3	A	818	NAD	O4B-C1B-N9A	2.12	112.53	108.10
2	A	449	FAD	O4B-C1B-N9A	2.49	113.30	108.10
3	A	818	NAD	C3N-C2N-N1N	2.61	123.37	120.36
2	A	449	FAD	P-O3P-PA	3.20	141.72	132.73
3	A	818	NAD	C3N-C7N-N7N	3.70	121.87	117.82
3	A	818	NAD	O4D-C1D-N1N	4.41	112.98	108.13
3	A	818	NAD	N3A-C2A-N1A	4.42	132.28	128.89
2	A	449	FAD	C4-N3-C2	7.13	121.41	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	449	FAD	2	0
3	A	818	NAD	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	446/447 (99%)	-0.87	3 (0%) 89 88	5, 16, 36, 57	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	220	ASP	3.1
1	A	447	ARG	2.6
1	A	124	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	OCS	A	42	9/10	0.99	0.08	-	9,14,16,19	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	FAD	A	449	53/53	0.98	0.08	-0.42	6,10,16,16	0
3	NAD	A	818	44/44	0.98	0.09	-0.76	18,21,23,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.