



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:05 PM GMT

PDB ID : 4NPP
Title : The GLIC-His10 wild-type structure in equilibrium between the open and locally-closed (LC) forms
Authors : Sauguet, L.; Shahsavar, A; Poitevin, F.; Huon, C.; Menny, A.; Nemecz, A.; Haouz, A.; Changeux, J.P.; Corringer, P.J.; Delarue, M.
Deposited on : 2013-11-22
Resolution : 3.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1005 (3.42-3.30)
Clashscore	102246	1076 (3.42-3.30)
Ramachandran outliers	100387	1059 (3.42-3.30)
Sidechain outliers	100360	1058 (3.42-3.30)
RSRZ outliers	91569	1010 (3.42-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	<div> <div>5%</div> <div>84%</div> <div>10%</div> <div>5%</div> </div>
1	B	329	<div> <div>5%</div> <div>84%</div> <div>11%</div> <div>5%</div> </div>
1	C	329	<div> <div>5%</div> <div>85%</div> <div>9%</div> <div>5%</div> </div>
1	D	329	<div> <div>6%</div> <div>84%</div> <div>10%</div> <div>5%</div> </div>
1	E	329	<div> <div>3%</div> <div>85%</div> <div>9%</div> <div>5%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 13465 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proton-gated ion channel.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	311	Total	C	N	O	S	0	21	0
			2692	1775	430	482	5			
1	B	311	Total	C	N	O	S	0	21	0
			2692	1775	430	482	5			
1	C	311	Total	C	N	O	S	0	21	0
			2692	1775	430	482	5			
1	D	311	Total	C	N	O	S	0	21	0
			2692	1775	430	482	5			
1	E	311	Total	C	N	O	S	0	21	0
			2692	1775	430	482	5			

There are 65 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ALA	-	EXPRESSION TAG	UNP Q7NDN8
A	318	GLY	-	EXPRESSION TAG	UNP Q7NDN8
A	319	GLY	-	EXPRESSION TAG	UNP Q7NDN8
A	320	HIS	-	EXPRESSION TAG	UNP Q7NDN8
A	321	HIS	-	EXPRESSION TAG	UNP Q7NDN8
A	322	HIS	-	EXPRESSION TAG	UNP Q7NDN8
A	323	HIS	-	EXPRESSION TAG	UNP Q7NDN8
A	324	HIS	-	EXPRESSION TAG	UNP Q7NDN8
A	325	HIS	-	EXPRESSION TAG	UNP Q7NDN8
A	326	HIS	-	EXPRESSION TAG	UNP Q7NDN8
A	327	HIS	-	EXPRESSION TAG	UNP Q7NDN8
A	328	HIS	-	EXPRESSION TAG	UNP Q7NDN8
A	329	HIS	-	EXPRESSION TAG	UNP Q7NDN8
B	1	ALA	-	EXPRESSION TAG	UNP Q7NDN8
B	318	GLY	-	EXPRESSION TAG	UNP Q7NDN8
B	319	GLY	-	EXPRESSION TAG	UNP Q7NDN8
B	320	HIS	-	EXPRESSION TAG	UNP Q7NDN8
B	321	HIS	-	EXPRESSION TAG	UNP Q7NDN8
B	322	HIS	-	EXPRESSION TAG	UNP Q7NDN8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	323	HIS	-	EXPRESSION TAG	UNP Q7NDN8
B	324	HIS	-	EXPRESSION TAG	UNP Q7NDN8
B	325	HIS	-	EXPRESSION TAG	UNP Q7NDN8
B	326	HIS	-	EXPRESSION TAG	UNP Q7NDN8
B	327	HIS	-	EXPRESSION TAG	UNP Q7NDN8
B	328	HIS	-	EXPRESSION TAG	UNP Q7NDN8
B	329	HIS	-	EXPRESSION TAG	UNP Q7NDN8
C	1	ALA	-	EXPRESSION TAG	UNP Q7NDN8
C	318	GLY	-	EXPRESSION TAG	UNP Q7NDN8
C	319	GLY	-	EXPRESSION TAG	UNP Q7NDN8
C	320	HIS	-	EXPRESSION TAG	UNP Q7NDN8
C	321	HIS	-	EXPRESSION TAG	UNP Q7NDN8
C	322	HIS	-	EXPRESSION TAG	UNP Q7NDN8
C	323	HIS	-	EXPRESSION TAG	UNP Q7NDN8
C	324	HIS	-	EXPRESSION TAG	UNP Q7NDN8
C	325	HIS	-	EXPRESSION TAG	UNP Q7NDN8
C	326	HIS	-	EXPRESSION TAG	UNP Q7NDN8
C	327	HIS	-	EXPRESSION TAG	UNP Q7NDN8
C	328	HIS	-	EXPRESSION TAG	UNP Q7NDN8
C	329	HIS	-	EXPRESSION TAG	UNP Q7NDN8
D	1	ALA	-	EXPRESSION TAG	UNP Q7NDN8
D	318	GLY	-	EXPRESSION TAG	UNP Q7NDN8
D	319	GLY	-	EXPRESSION TAG	UNP Q7NDN8
D	320	HIS	-	EXPRESSION TAG	UNP Q7NDN8
D	321	HIS	-	EXPRESSION TAG	UNP Q7NDN8
D	322	HIS	-	EXPRESSION TAG	UNP Q7NDN8
D	323	HIS	-	EXPRESSION TAG	UNP Q7NDN8
D	324	HIS	-	EXPRESSION TAG	UNP Q7NDN8
D	325	HIS	-	EXPRESSION TAG	UNP Q7NDN8
D	326	HIS	-	EXPRESSION TAG	UNP Q7NDN8
D	327	HIS	-	EXPRESSION TAG	UNP Q7NDN8
D	328	HIS	-	EXPRESSION TAG	UNP Q7NDN8
D	329	HIS	-	EXPRESSION TAG	UNP Q7NDN8
E	1	ALA	-	EXPRESSION TAG	UNP Q7NDN8
E	318	GLY	-	EXPRESSION TAG	UNP Q7NDN8
E	319	GLY	-	EXPRESSION TAG	UNP Q7NDN8
E	320	HIS	-	EXPRESSION TAG	UNP Q7NDN8
E	321	HIS	-	EXPRESSION TAG	UNP Q7NDN8
E	322	HIS	-	EXPRESSION TAG	UNP Q7NDN8
E	323	HIS	-	EXPRESSION TAG	UNP Q7NDN8
E	324	HIS	-	EXPRESSION TAG	UNP Q7NDN8
E	325	HIS	-	EXPRESSION TAG	UNP Q7NDN8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	326	HIS	-	EXPRESSION TAG	UNP Q7NDN8
E	327	HIS	-	EXPRESSION TAG	UNP Q7NDN8
E	328	HIS	-	EXPRESSION TAG	UNP Q7NDN8
E	329	HIS	-	EXPRESSION TAG	UNP Q7NDN8

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ni 1 1	0	0
2	A	1	Total Ni 1 1	0	0
2	D	1	Total Ni 1 1	0	0
2	C	1	Total Ni 1 1	0	0
2	E	1	Total Ni 1 1	0	0

- Molecule 1: Proton-gated ion channel





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	113.55Å 127.62Å 185.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.16 – 3.35 19.16 – 3.35	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.16-3.35) 99.8 (19.16-3.35)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.14 (at 3.36Å)	Xtriage
Refinement program	BUSTER 2.11.4	Depositor
R, R_{free}	0.227 , 0.248 0.235 , 0.258	Depositor DCC
R_{free} test set	1969 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	108.2	Xtriage
Anisotropy	0.454	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 65.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 39210 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	13465	wwPDB-VP
Average B, all atoms (Å ²)	116.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.46	0/2765	0.65	0/3782
1	B	0.44	0/2765	0.63	0/3782
1	C	0.44	0/2765	0.64	0/3782
1	D	0.45	0/2765	0.65	0/3782
1	E	0.43	0/2765	0.63	0/3782
All	All	0.44	0/13825	0.64	0/18910

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2692	0	2718	18	2
1	B	2692	0	2717	19	0
1	C	2692	0	2718	18	0
1	D	2692	0	2718	18	2
1	E	2692	0	2718	18	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1	0	0	0	0
2	E	1	0	0	0	0
All	All	13465	0	13589	80	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 80 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:201:ILE:HG12	1:B:242[B]:VAL:HG11	1.65	0.79
1:C:63:VAL:HG21	1:D:136:ASP:CG	2.10	0.72
1:B:201:ILE:HG12	1:B:242[B]:VAL:CG1	2.20	0.70
1:D:7:PRO:HG3	1:D:135:VAL:HG21	1.80	0.62
1:E:204:PRO:HB2	1:E:238[B]:PHE:CZ	2.35	0.62

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:ASP:OD2	1:D:145:ASP:OD2[3_555]	1.85	0.35
1:A:166:THR:CG2	1:D:97:ASP:OD2[3_555]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/329 (100%)	317 (96%)	13 (4%)	0	100	100
1	B	330/329 (100%)	319 (97%)	9 (3%)	2 (1%)	30	70
1	C	330/329 (100%)	318 (96%)	12 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	330/329 (100%)	323 (98%)	7 (2%)	0	100	100
1	E	330/329 (100%)	316 (96%)	14 (4%)	0	100	100
All	All	1650/1645 (100%)	1593 (96%)	55 (3%)	2 (0%)	56	89

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	244[A]	THR
1	B	244[B]	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/294 (102%)	288 (96%)	11 (4%)	41	76
1	B	299/294 (102%)	290 (97%)	9 (3%)	48	80
1	C	299/294 (102%)	289 (97%)	10 (3%)	45	79
1	D	299/294 (102%)	289 (97%)	10 (3%)	45	79
1	E	299/294 (102%)	291 (97%)	8 (3%)	52	82
All	All	1495/1470 (102%)	1447 (97%)	48 (3%)	45	80

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	42	PHE
1	C	170	LYS
1	E	166	THR
1	C	63	VAL
1	C	111	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	311/329 (94%)	-0.01	18 (5%) 26 26	66, 109, 154, 219	0
1	B	311/329 (94%)	0.06	17 (5%) 29 28	77, 119, 162, 222	0
1	C	311/329 (94%)	0.04	18 (5%) 26 26	75, 122, 165, 213	0
1	D	311/329 (94%)	-0.02	19 (6%) 25 25	72, 110, 150, 193	0
1	E	311/329 (94%)	-0.04	11 (3%) 48 47	71, 110, 153, 183	0
All	All	1555/1645 (94%)	0.01	83 (5%) 30 30	66, 114, 158, 222	0

The worst 5 of 83 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	59	SER	5.0
1	C	58	ARG	4.8
1	A	244[A]	THR	4.2
1	A	178	ASP	4.2
1	A	68	PRO	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NI	B	401	1/1	0.86	0.26	-	141,141,141,141	0
2	NI	A	401	1/1	0.94	0.15	-	135,135,135,135	0
2	NI	D	401	1/1	0.91	0.14	-	110,110,110,110	0
2	NI	C	401	1/1	0.85	0.10	-	116,116,116,116	0
2	NI	E	401	1/1	0.88	0.17	-	119,119,119,119	0

6.5 Other polymers

There are no such residues in this entry.