



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:03 AM GMT

PDB ID : 3NT6
Title : Structure of the Shewanella loihica PV-4 NADH-dependent persulfide reductase C43S/C531S Double Mutant
Authors : Sazinsky, M.H.; Crane, E.J.; Warner, M.D.; Lukose, V.; Lee, K.H.; Lopez, K.
Deposited on : 2010-07-02
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	574	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 21%, green 73%, grey 1%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> 73% 21% ... </div> </div>
1	B	574	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 74%, yellow 20%, orange 5%, red 1%, grey 1%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> 74% 20% ... </div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9342 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FAD-dependent pyridine nucleotide-disulphide oxidoreductase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	565	Total	C	N	O	S	Se	0	0	0
			4295	2694	762	821	2	16			
1	B	565	Total	C	N	O	S	Se	0	0	0
			4294	2694	763	820	2	15			


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	43	SER	CYS	ENGINEERED MUTATION	UNP A3QAV3
A	531	SER	CYS	ENGINEERED MUTATION	UNP A3QAV3
A	567	LEU	-	EXPRESSION TAG	UNP A3QAV3
A	568	GLU	-	EXPRESSION TAG	UNP A3QAV3
A	569	HIS	-	EXPRESSION TAG	UNP A3QAV3
A	570	HIS	-	EXPRESSION TAG	UNP A3QAV3
A	571	HIS	-	EXPRESSION TAG	UNP A3QAV3
A	572	HIS	-	EXPRESSION TAG	UNP A3QAV3
A	573	HIS	-	EXPRESSION TAG	UNP A3QAV3
A	574	HIS	-	EXPRESSION TAG	UNP A3QAV3
B	43	SER	CYS	ENGINEERED MUTATION	UNP A3QAV3
B	531	SER	CYS	ENGINEERED MUTATION	UNP A3QAV3
B	567	LEU	-	EXPRESSION TAG	UNP A3QAV3
B	568	GLU	-	EXPRESSION TAG	UNP A3QAV3
B	569	HIS	-	EXPRESSION TAG	UNP A3QAV3
B	570	HIS	-	EXPRESSION TAG	UNP A3QAV3
B	571	HIS	-	EXPRESSION TAG	UNP A3QAV3
B	572	HIS	-	EXPRESSION TAG	UNP A3QAV3
B	573	HIS	-	EXPRESSION TAG	UNP A3QAV3
B	574	HIS	-	EXPRESSION TAG	UNP A3QAV3

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Cl 1 1	0	0
2	A	1	Total Cl 1 1	0	0

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O P 53 27 9 15 2	0	0
3	B	1	Total C N O P 53 27 9 15 2	0	0

- Molecule 4 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
4	B	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	259	Total	O	0	0
			259	259		
5	B	290	Total	O	0	0
			290	290		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FAD-dependent pyridine nucleotide-disulphide oxidoreductase



- Molecule 1: FAD-dependent pyridine nucleotide-disulphide oxidoreductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	133.71 Å 133.71 Å 79.24 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	115.00 – 2.00 38.31 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.5 (115.00-2.00) 99.9 (38.31-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.85 (at 2.00 Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.122 , 0.155 0.178 , 0.207	Depositor DCC
R_{free} test set	5160 reflections (5.07%)	DCC
Wilson B-factor (Å ²)	35.8	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 40.0	EDS
Estimated twinning fraction	0.023 for H, K, L 0.517 for -H, H+K, -L 0.460 for -H, -K, L 0.024 for -h,-k,l 0.045 for h,-h-k,-l 0.000 for -k,-h,-l	Xtriage
Reported twinning fraction	0.023 for H, K, L 0.517 for -H, H+K, -L 0.460 for -H, -K, L	Depositor
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 107016 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9342	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: COA, FAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.42	22/4350 (0.5%)	1.24	29/5865 (0.5%)
1	B	1.44	21/4349 (0.5%)	1.28	29/5863 (0.5%)
All	All	1.43	43/8699 (0.5%)	1.26	58/11728 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 43 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	101	GLU	CB-CG	-8.84	1.35	1.52
1	B	33	GLU	CD-OE1	8.68	1.35	1.25
1	B	331	ALA	CA-CB	7.62	1.68	1.52
1	A	450	TYR	CD1-CE1	7.35	1.50	1.39
1	A	409	PHE	CE2-CZ	7.17	1.50	1.37

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	34	ARG	NE-CZ-NH1	14.21	127.40	120.30
1	A	34	ARG	NE-CZ-NH2	-11.84	114.38	120.30
1	B	22	ARG	NE-CZ-NH2	-10.86	114.87	120.30
1	B	134	ARG	NE-CZ-NH1	-10.66	114.97	120.30
1	B	511	VAL	CG1-CB-CG2	10.17	127.17	110.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	290	MSE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4295	0	4298	77	2
1	B	4294	0	4300	95	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	53	0	31	1	0
3	B	53	0	30	0	0
4	A	48	0	32	0	0
4	B	48	0	32	0	0
5	A	259	0	0	5	0
5	B	290	0	0	11	0
All	All	9342	0	8723	167	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 167 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:290:MSE:CE	1:B:339:GLU:HA	1.70	1.20
1:A:192:ARG:NH2	1:A:564:ALA:O	1.74	1.19
1:A:290:MSE:HE3	1:A:339:GLU:HA	1.16	1.12
1:B:290:MSE:HE1	1:B:339:GLU:CG	1.83	1.09
1:B:290:MSE:HE1	1:B:339:GLU:HG2	1.08	1.08

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:HIS:ND1	1:A:512:ASP:OD2[2_555]	2.09	0.11
1:A:229:ASP:OD2	1:A:387:SER:OG[2_555]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	563/574 (98%)	543 (96%)	18 (3%)	2 (0%)	39	33
1	B	563/574 (98%)	547 (97%)	12 (2%)	4 (1%)	26	19
All	All	1126/1148 (98%)	1090 (97%)	30 (3%)	6 (0%)	34	26

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	482	ASN
1	B	483	LEU
1	A	10	VAL
1	A	456	VAL
1	B	10	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	449/443 (101%)	416 (93%)	33 (7%)	17	11
1	B	449/443 (101%)	408 (91%)	41 (9%)	12	6

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	898/886 (101%)	824 (92%)	74 (8%)	14 9

5 of 74 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	34	ARG
1	B	103	GLN
1	B	494	ASN
1	B	42	ASN
1	B	67	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	62	GLN
1	B	205	GLN
1	B	482	ASN
1	B	131	HIS
1	B	221	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FAD	A	900	-	48,58,58	1.38	7 (14%)	54,89,89	2.55	14 (25%)
4	COA	A	901	-	40,50,50	1.56	2 (5%)	50,75,75	3.01	12 (24%)
3	FAD	B	900	-	48,58,58	1.47	10 (20%)	54,89,89	3.22	13 (24%)
4	COA	B	901	-	40,50,50	1.96	3 (7%)	50,75,75	2.60	15 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	900	-	-	0/30/50/50	0/6/6/6
4	COA	A	901	-	-	0/44/64/64	0/3/3/3
3	FAD	B	900	-	-	0/30/50/50	0/6/6/6
4	COA	B	901	-	-	0/44/64/64	0/3/3/3

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	900	FAD	O2'-C2'	-2.95	1.36	1.43
3	A	900	FAD	O4'-C4'	-2.77	1.37	1.43
3	A	900	FAD	C4X-C10	-2.51	1.36	1.41
3	B	900	FAD	C9A-N10	-2.14	1.35	1.38
3	B	900	FAD	C5'-C4'	2.11	1.54	1.51

The worst 5 of 54 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	900	FAD	N3A-C2A-N1A	-17.81	115.25	128.89
4	A	901	COA	N3A-C2A-N1A	-16.45	116.30	128.89
3	A	900	FAD	N3A-C2A-N1A	-13.75	118.37	128.89
4	B	901	COA	N3A-C2A-N1A	-13.52	118.55	128.89
3	B	900	FAD	O4'-C4'-C5'	-6.22	96.65	110.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	900	FAD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	549/574 (95%)	-0.18	4 (0%)	89 89	25, 34, 53, 70	0
1	B	549/574 (95%)	-0.25	1 (0%)	95 95	22, 33, 50, 65	0
All	All	1098/1148 (95%)	-0.21	5 (0%)	91 92	22, 34, 52, 70	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	565	SER	2.6
1	A	372	GLY	2.5
1	B	444	LEU	2.4
1	A	500	ASN	2.1
1	A	230	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CL	B	575	1/1	1.00	0.17	1.83	28,28,28,28	0
4	COA	A	901	48/48	0.97	0.16	0.36	24,39,66,69	0
4	COA	B	901	48/48	0.98	0.15	0.08	27,41,60,62	0
3	FAD	B	900	53/53	0.97	0.12	0.08	22,29,34,37	0
3	FAD	A	900	53/53	0.98	0.12	0.04	25,31,36,39	0
2	CL	A	575	1/1	1.00	0.12	-0.24	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.