



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:11 PM GMT

PDB ID : 1NW1  
Title : Crystal Structure of Choline Kinase  
Authors : Peisach, D.; Gee, P.; Kent, C.; Xu, Z.  
Deposited on : 2003-02-05  
Resolution : 2.02 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	7858 (2.04-2.00)
Clashscore	102246	9060 (2.04-2.00)
Ramachandran outliers	100387	8952 (2.04-2.00)
Sidechain outliers	100360	8951 (2.04-2.00)
RSRZ outliers	91569	7873 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	429	<div> <div>3%</div> <div>71%</div> <div>13%</div> <div>15%</div> </div>
1	B	429	<div> <div>4%</div> <div>69%</div> <div>14%</div> <div>•</div> <div>17%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CA	A	430	-	-	-	X
2	CA	B	430	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6153 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Choline kinase (49.2 kD).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	365	Total	C	N	O	S	0	0	0
			2945	1904	492	535	14			
1	B	357	Total	C	N	O	S	0	0	0
			2864	1856	480	515	13			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	183	Total	O	0	0
			183	183		
3	B	159	Total	O	0	0
			159	159		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A:
- 
- | Amino Acid | Count | Category |
|------------|-------|----------|
| MET        | 1     | Grey     |
| SER        | 1     | Green    |
| SER        | 1     | Green    |
| ARG        | 1     | Green    |
| LYS        | 1     | Green    |
| VAL        | 1     | Green    |
| SER        | 1     | Green    |
| ARG        | 1     | Green    |
| ALA        | 1     | Green    |
| HIS        | 1     | Green    |
| TYR        | 1     | Green    |
| ASP        | 1     | Green    |
| GLU        | 1     | Green    |
| GLU        | 1     | Green    |
| LEU        | 1     | Green    |
| ALA        | 1     | Green    |
| SER        | 1     | Green    |
| ALA        | 1     | Green    |
| ASN        | 1     | Green    |
| ASN        | 1     | Green    |
| MET        | 1     | Green    |
| SER        | 1     | Green    |
| LEU        | 1     | Green    |
| VAL        | 1     | Green    |
| GLU        | 1     | Green    |
| GLY        | 1     | Green    |
| HIS        | 1     | Green    |
| PHE        | 1     | Green    |
| ARG        | 1     | Green    |
| G32        | 1     | Green    |
| M33        | 1     | Yellow   |
| K34        | 1     | Yellow   |
| L37        | 1     | Green    |
| S38        | 1     | Green    |
| D41        | 1     | Yellow   |
| L42        | 1     | Green    |
| D43        | 1     | Green    |
| M47        | 1     | Yellow   |
| D210       | 1     | Yellow   |
| P50        | 1     | Yellow   |
| K53        | 1     | Yellow   |
| E54        | 1     | Yellow   |
| R62        | 1     | Yellow   |
| T70        | 1     | Green    |
| L73        | 1     | Yellow   |
| R77        | 1     | Yellow   |
| I78        | 1     | Yellow   |
| M85        | 1     | Yellow   |
| S94        | 1     | Yellow   |

- Chain B:
- 
- 4% 69% 14% 17%
- Met Ser Arg Lys Val Ser Arg Ala His Tyr Asp Glu Leu Ala Ser Ala Asn Met Ser Leu Val Ala Glu Gly His Phe Arg Gly Met Lys Glu Leu Ser Thr M40 D41 D45 A46 N47 P50 K53 E54 R55 M58 R62 G66 T70
- E74 H75 L76 R77 I78 T81 G84 H85 S86 M87 T101 R102 H103 L110 P116 E117 T118 L131 L150 S156 L159 S160 C161 I164 S165 L166 A167 H168 R176 K179 E196 A197 L198 Q199 R200 F201 L202 K203 Q204 L205 T206 G207 T208 V209 D210 A211 F212
- A330 Q336 F340 P341 L347 F350 R355 L372 V380 P381 F386 R409 S413 A426 SER HIS GLN

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.37Å 102.14Å 85.37Å 90.00° 90.66° 90.00°	Depositor
Resolution (Å)	37.54 – 2.02 37.54 – 2.02	Depositor EDS
% Data completeness (in resolution range)	94.4 (37.54-2.02) 94.5 (37.54-2.02)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.54 (at 2.01Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.209 , 0.231 0.205 , 0.228	Depositor DCC
$R_{free}$ test set	4367 reflections (7.41%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.9	Xtriage
Anisotropy	0.291	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 58.2	EDS
Estimated twinning fraction	0.042 for h,-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 60767 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6153	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/3023	0.55	0/4103
1	B	0.40	2/2941 (0.1%)	0.55	0/3996
All	All	0.38	2/5964 (0.0%)	0.55	0/8099

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	74	GLU	CB-CG	-5.36	1.42	1.52
1	B	196	GLU	CB-CG	-5.33	1.42	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2945	0	2860	45	0
1	B	2864	0	2769	43	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	183	0	0	7	0
3	B	159	0	0	3	0
All	All	6153	0	5629	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 88 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:200:ARG:HG3	1:B:200:ARG:HH11	1.45	0.82
1:A:101:ILE:HD11	1:A:102:ARG:HH21	1.48	0.78
1:A:159:LEU:CD2	1:A:263:LEU:HD11	2.13	0.77
1:B:200:ARG:CG	1:B:200:ARG:HH11	1.98	0.75
1:B:198:LEU:HD21	1:B:386:PHE:HA	1.70	0.73

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	361/429 (84%)	351 (97%)	10 (3%)	0	100	100
1	B	353/429 (82%)	342 (97%)	11 (3%)	0	100	100
All	All	714/858 (83%)	693 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	317/381 (83%)	312 (98%)	5 (2%)	70	72
1	B	304/381 (80%)	294 (97%)	10 (3%)	45	42
All	All	621/762 (82%)	606 (98%)	15 (2%)	57	56

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	81	ILE
1	B	85	MET
1	B	236	GLU
1	B	55	ARG
1	B	220	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	424	ASN
1	B	182	GLN
1	B	339	ASN
1	A	420	GLN
1	B	336	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	365/429 (85%)	0.19	15 (4%) 41 43	9, 23, 45, 67	0
1	B	357/429 (83%)	0.16	17 (4%) 34 36	9, 23, 46, 73	0
All	All	722/858 (84%)	0.17	32 (4%) 38 40	9, 23, 46, 73	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	426	ALA	4.6
1	B	208	THR	4.4
1	B	101	ILE	3.4
1	A	209	VAL	3.2
1	B	210	ASP	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	B	430	1/1	0.99	0.18	4.11	23,23,23,23	0
2	CA	A	430	1/1	0.98	0.15	2.60	32,32,32,32	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.